Physics 122

Motion in 1D Review

- 1. Compare the distance covered for two cars, one traveling 30.6 m/s and the other 36.1 m/s, as they come to a stop under a constant acceleration of -6.45 m/s^2 ? (Hint: first use the definition of acceleration to find the stopping time for each car)
- 2. A car undergoes a constant acceleration from rest to 28 m/s in 9.5 s. What distance was covered in that time?
- 3. Not noticing a red light a drivers slams on the brakes squeeling to a halt in 3.75 s. Just before hitting the brakes the car was traveling 17 m/s and was 30 m from the light. a) What was the average acceleration of the car? b) Determine if the driver able to stop before reaching the traffic light by finding the distance required to stop.
- 4. An airplane lands with a speed of 70 m/s. After 3.5 s the airplane is traveling 17.5 m/s. a) What was the average acceleration of the airplane? b) What distance does the airplane need to stop?

- 5. During take off a Boeing 747 airplane accelerates at a constant $10.8 \frac{m}{s^2}$. The airplane accelerated, from rest, for 7.8 s before it left the ground. **a**) With what speed did the airplane leave the ground? **b**) What distance was required for take-off?
- 6. A ball is thrown upwards, on the Earth $(a_{gravity} = -9.8 \text{ m/s}^2)$, with an initial speed of +17 m/s. a) How long will the ball be traveling upwards? b) How high up will the ball travel? c) What is the instantaneous velocity after 1.25 s? 2.25 s?
- 7. A loonie dropped from the observation deck on the CN Tower in Toronto takes 8.35 s to hit the ground. **a**) Assuming no air resistance, with what speed is the loonie striking the ground? **b**) How high is the observation deck from the ground?
- 8. During its fall to Earth, hail stones from cumulonimbus clouds reach a terminal velocity (a constant speed) because of air resistance. Assuming no air resistance, calculate at what speed a hail stone would strike the Earth if it continued to accelerate from gravity during its 20.2 s fall to the ground.









- 9. A ball is thrown straight up (assume no air resistance) at the surface of the Earth with $v_o = 25 \text{ m/s}$. How long will the ball be in the air?
- 10. The upward velocity of a cannon ball is initially 142 m/s. How long after the cannon ball is fired will it be 51.0 m above the ground?
- 11. What is the instantaneous velocity of a ball (launched upwards) that has a displacement of 21 m above the ground if the initial velocity was 32 m/s?
- 12. Use $d = v_o t + \frac{1}{2}at^2$ and $a = \frac{v_f v_o}{t}$ to derived $v_f^2 = v_o^2 + 2ad$

Physics 122 Motion in 1D Review Mr. P. MacDonald 1/31/2010

Answer List

- 1. Slow car: d = 72.6 m; Fast car: d = 101 m
- 3. a) $a_{avg} = -4.5 \text{ m/s}^2$, b) No, 32 m is needed to stop.
- 5. **a)** $v_f = 84 \text{ m/s}$, **b)** d = 342 m
- 7. **a)** $v_f = 82 \text{ m/s}, \text{ b)height} = 342 \text{ m}$
- 9. $t = 5.1 \,\mathrm{s}$
- 11. v = +25 m/s or -25m/s

- 2. $d = 133 \,\mathrm{m}$
- 4. **a)** $a_{avg} = -15 \text{ m/s}^2$, **b)** d = 163 m
- 6. **a)** t = 1.73 s, **b)** d = 15 m, **c)** $v_{@1.25} = 4.75 \text{ m/s}$, $v_{@2.25} = -5.05 \text{ m/s}$
- 8. $v_f = 198 \,\mathrm{m/s}$
- 10. t = 0.364 and 28.6 seconds
- 12.