

$$\textcircled{16} \quad 8^{1/4} \cdot \left(\frac{1}{4}\right)^{x/2} = 16^{3/4}$$

$$8^{1/4} \cdot [(4)^{-1}]^{x/2} = 16^{3/4}$$

$$8^{1/4} \cdot 4^{-x/2} = 16^{3/4}$$

$$(2^3)^{1/4} \cdot (2^2)^{-x/2} = (2^4)^{3/4}$$

$$2^{3/4} \cdot 2^{-x} = 2^3$$

$$2^{3/4 - x} = 2^3$$

$$\frac{3}{4} - x = 3$$

$$\frac{3}{4} - 3 = x$$

$$\textcircled{\frac{-9}{4} = x}$$

$$\textcircled{17} \quad \frac{(9^{2x-1}) \cdot (3^{3x})^2}{(27^{x+2})^4} = 1$$

$$\frac{(9^{2x-1}) \cdot 3^{6x}}{27^{4x+8}} = 1$$

$$\frac{(3^2)^{2x-1} \cdot 3^{6x}}{(3^3)^{4x+8}} = 3^0$$

$$\frac{3^{4x-2} \cdot 3^{6x}}{3^{12x+24}} = 3^0$$

$$3^{4x-2+6x-(12x+24)} = 3^0$$

$$4x - 2 + 6x - (12x + 24) = 0$$

$$10x - 2 - 12x - 24 = 0$$

$$-2x - 26 = 0$$

$$-2x = 26$$

$$\textcircled{x = -13}$$

RATIONAL EXPONENTS

In our discussion of exponential functions and equations thus far, we have only considered exponents which were integers, that is, ..., -3, -2, -1, 0, 1, 2, 3, ...

We extend our knowledge of exponents in this section by considering exponents that are **rational numbers**. A *rational number* is any number that can be written as a ratio of integers.

Some examples of rational numbers are: $\frac{2}{3}, \frac{-5}{13}, \frac{0}{12}, \frac{-12}{-87}, \dots$

In the definition, it says that a rational number is any number that *can* be written as a ratio of integers.

$$27^{\frac{2}{3}} = 9$$

Thus, the following are also rational numbers:

$$15 = \frac{15}{1}, 0 = \frac{0}{114}, \sqrt{25} = \frac{5}{1}.$$

There are some real numbers that are **not rational**:

$$\sqrt{13}, \sqrt{43}, \pi$$

And, of course, some numbers are not real, and therefore, **not rational**:

$$\frac{7}{0}, \sqrt{-25}$$

$$27^{\frac{2}{3}} = \left(\sqrt[3]{27} \right)^2 = 9$$

The basic rule we use to evaluate expressions with rational exponents is the following:

$$b^{\frac{m}{n}} = \left(\sqrt[n]{b} \right)^m$$

What this does is change the expression to an equivalent expression in radical form. A

radical is a number in the form $\sqrt{5}, \sqrt[3]{13}, \sqrt[4]{112}$, etc.

A radical, then, is any root of a number. It can be square (2^{nd}) root, cube (3^{rd}) root, 4^{th} -root, etc. **If we want the square root of a number, we do not include the number 2 in our expression.**

For example, $\sqrt{49}$ is taken to mean the square root of $49 = 7$.

If we want a different root, we have to indicate such: $\sqrt[3]{8} = 2$ or $\sqrt[4]{32} = 2$.

Examples:

Write the following as radicals.

1. $13^{\frac{2}{5}} = (\sqrt[5]{13})^2$

$(\sqrt[5]{13})^2$

2. $-123^{\frac{9}{2}} = -(\sqrt[2]{123})^9$

Recall that the exponent only applies to the base directly

below it. $-123^{\frac{9}{2}}$ can be thought of as: $(-1)(123)^{\frac{9}{2}}$.

3. $(-17)^{\frac{1}{5}} = \sqrt[5]{-17}$ The brackets mean that the base is (-17).

4. $\left(\frac{5}{6}\right)^{\frac{4}{5}} = \left(\sqrt[5]{\frac{5}{6}}\right)^4$

5. $\left(\frac{2}{7}\right)^{-\frac{6}{11}} = \left(\frac{7}{2}\right)^{\frac{6}{11}} = \left(\sqrt[11]{\frac{7}{2}}\right)^6$

Recall that a fraction raised to a negative exponent can be written as the reciprocal of the base raised to the positive value of the exponent.

Write each using positive fractional exponents.

$$6. \quad \sqrt[5]{14} = 14^{\frac{1}{5}}$$

$$7. \quad (\sqrt[3]{111})^4 = 111^{\frac{4}{3}}$$

$$8. \quad \frac{1}{\sqrt{13}} = \frac{1}{13^{\frac{1}{2}}} = 13^{-\frac{1}{2}}$$

$$9. \quad \frac{1}{\sqrt[7]{5^{-x}}} = \frac{1}{5^{\frac{-x}{7}}} = 5^{\frac{x}{7}}$$

Evaluate without using a calculator.

10. $25^{\frac{1}{2}} = \sqrt{25} = 5$

11. $-36^{\frac{1}{2}} = -\sqrt{36} = -6$

12. $(-36)^{\frac{1}{2}} = \sqrt{-36} =$ no real solution. There is no real solution when you take an even root of a negative quantity. Thus, $\sqrt[4]{-56}$, $\sqrt[4]{-47}$, have no real solutions.

13. $(-8)^{\frac{-1}{3}} = \sqrt[3]{-8} = -2$, since $(-2)(-2)(-2) = -8$. There is a real solution when you take an odd root of a negative quantity.

14. $\left(\frac{625}{81}\right)^{\frac{1}{4}} = \sqrt[4]{\frac{625}{81}} = \frac{5}{3}$

$$15. \quad \left(\frac{4}{49}\right)^{\frac{-3}{2}} = \left(\frac{49}{4}\right)^{\frac{3}{2}} = \left(\sqrt{\frac{49}{4}}\right)^3 = \left(\frac{7}{2}\right)^3 = \frac{343}{8}$$

$$16. \quad 16^{\frac{3}{2}} + 27^{\frac{-1}{3}} - 4^{\frac{5}{2}}$$
$$(\sqrt{16})^3 + \frac{1}{\sqrt[3]{27}} - (\sqrt{4})^5$$

$$4^3 + \frac{1}{3} - 2^5$$

$$64 + \frac{1}{3} - 32 = 32 + \frac{1}{3} = \frac{97}{3}$$