

## Warm up

$$f(x) = x^3 + 2x$$

$$g(x) = x - 2$$

Find

$$(f \circ g)(x)$$

$$f(g(x))$$

$$f(x-2) = (x-2)^3 + 2(x-2)$$

$$= x^3 - 6x^2 + 12x - 8 + 2x - 4$$

$$= x^3 - 6x^2 + 14x - 12$$

$$g(f(2))$$

$$f(2) = (2)^3 + 2(2)$$

$$= 8 + 4$$

$$= 12$$

$$g(12) = 12 - 2$$

$$= 10$$

## Questions From Homework

$$\textcircled{1} \text{ e) } y = 6x^2 - 7x + 2$$

• Roots ( $y=0$ )

$$0 = 6x^2 - 7x + 2$$

$$0 = (6x^2 - 3x)(4x + 2)$$

$$0 = 3x(2x - 1) - 2(2x - 1)$$

$$0 = (3x - 2)(2x - 1)$$

$$\begin{array}{l|l} 3x - 2 = 0 & 2x - 1 = 0 \\ 3x = 2 & 2x = 1 \\ x = \frac{2}{3} & x = \frac{1}{2} \end{array}$$

Trinomial Decomposition

$$\underline{-4}x\underline{-3} = 12$$

$$\underline{-4} + \underline{-3} = -7$$

Vertex (Complete the square)

$$y = 6x^2 - 7x + 2$$

$$y - 2 = 6x^2 - 7x$$

$$\frac{7}{6} \times \frac{1}{2} = \left(\frac{7}{12}\right)^2 = \frac{49}{144}$$

$$y - 2 = 6\left(x^2 - \frac{7}{6}x\right)$$

$$y - 2 \overset{+\frac{49}{144}}{=} 6\left(x^2 - \frac{7}{6}x + \frac{49}{144}\right)$$

$$y - 2 + \frac{49}{24} = 6\left(x - \frac{7}{12}\right)^2$$

$$y - \frac{48}{24} + \frac{49}{24} = 6\left(x - \frac{7}{12}\right)^2$$

$$y = 6\left(x - \frac{7}{12}\right)^2 - \frac{1}{24}$$

$$V = \left(\frac{7}{12}, \frac{1}{24}\right) \text{ or } (0.58\bar{3}, -0.041\bar{6})$$

## Polynomial Functions

**Polynomial** - an algebraic expression consisting of two or more terms. A polynomial usually contains only one variable. Within each term the variable is raised to a non-negative integer power, and is multiplied by a constant. The simplest types of polynomials are binomials (two terms) and trinomials (three terms)

**Degree of a Polynomial** - the greatest power to which the variable is raised; for example, the degree of the trinomial  $x^4 - 2x + 5$  is 4

A **polynomial** function with real coefficients can be represented by

$$y = f(x) = ax^n + bx^{n-1} + cx^{n-2} + \dots + \square x^0$$



where  $a, b, c, \text{ etc.}$  are real numbers. The shape of the graph of the function is affected by the value of  $n$  (**the Degree of the Polynomial**), the values of the coefficients, and whether the value of  $a$  is positive or negative.

# Cubic Functions


3rd degree Polynomials.  $\longleftrightarrow y = ax^3 + bx^2 + cx + d$

factored form  $y = \underline{\underline{a}}(x - r_1)(x - r_2)(x - r_3)$

$a > 0$  (Positive) Starts in Q3  
Ends in Q1




$a < 0$  (Negative) Starts in Q2  
Ends in Q4



A cubic function has three roots. Either one or three of these roots will be real numbers. Any other roots are complex numbers. The number of *x-intercepts* on the graph of the corresponding cubic function  $y=f(x)$  depends on the nature of the roots.

**Three different real roots**



$$y = (x-1)(x-2)(x+3)$$

↙ a=1


① Roots ( $y=0$ )  
 $0 = (x-1)(x-2)(x+3)$   
 $x = -3, 1, 2$

② y int ( $x=0$ )  
 $y = (0-1)(0-2)(0+3)$   
 $y = (-1)(-2)(3)$   
 $y = 6$

③ Degree  $\rightarrow 3^{\text{rd}}$

④ Stretch:  $a=1$

Three real roots (2 are equal)



$a=1$

$$y = (x+3)(x-2)^2$$

$$y = (x+3)(x-2)(x-2)$$

① Roots ( $y=0$ )

$$0 = (x+3)(x-2)(x-2)$$

$$x = -3, 2, 2 \leftarrow \text{Double Root}$$

② yint ( $x=0$ )

$$y = (0+3)(0-2)^2$$

$$y = (3)(4)$$


$$y = 12$$

③ Degree = 3<sup>rd</sup>

④ Stretch:  $a=1$

**Three equal real roots**

$$y = -(x - 2)^3$$



**Local Maximum** - is the highest point in its immediate region of  $x$ -values.  
This may or may not be the greatest value of the function over its entire domain.

**Local Minimum** - is the lowest point in its immediate region of  $x$ -values.  
This may or may not be the smallest value of the function over its entire domain.



### Calculating Max and Min values on the TI-83



# Homework

