


Homework - Worksheet



Water Equilibrium

Conductivity is due to the presence of ions. For water:



- therefore $K = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]}$ is very small


- slight conductivity shows that equilibrium greatly favors water molecules (less than 2 H⁺ per billion water)

- therefore the concentration of water in pure water and in dilute aqueous solutions is essentially constant and can be combined with the equilibrium constant to produce a new constant called the *Ion Product Constant*

Ionization Constant for water (ion product constant)

$$K_w = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14} \quad \text{at SATP}$$

Since [H⁺] and [OH⁻] are found in 1:1 ratio



[H⁺_(aq)] = [OH⁻_(aq)] = 1.0 x 10⁻⁷ mol/L in **neutral** solutions.

Arrhenius's Theory - acid is a substance that ionizes water produce H⁺ ions.

- additional ions produced by the acid increases the H⁺ concentration in the water. (more acid, more H⁺)

Therefore acids always have a [H⁺] > 10⁻⁷ mol/L

Basic solutions produce a [OH⁻] greater than 10⁻⁷ mol/L

K_w can be used to calculate either [H⁺] or [OH⁻]

$$\text{since } K_w = [\text{H}^+][\text{OH}^-] \text{ then } [\text{H}^+] = K_w / [\text{OH}^-]$$

$$\text{and } [\text{OH}^-] = K_w / [\text{H}^+]$$

pH and pOH

$$\text{pH} + \text{pOH} = 14.00$$

$$\text{pH} = -\log[\text{H}^+_{(\text{aq})}] \quad \text{pOH} = -\log[\text{OH}^-_{(\text{aq})}]$$

$$[\text{H}^+_{(\text{aq})}] = 10^{-\text{pH}} \quad [\text{OH}^-_{(\text{aq})}] = 10^{-\text{pOH}}$$

Ex. Calculate the pH of a solution where $[\text{H}^+_{(\text{aq})}] = 3.24 \times 10^{-4}\text{M}$.

$$\text{pH} = -\log[\text{H}^+_{(\text{aq})}]$$

$$\text{pH} = -\log[3.24 \times 10^{-4}]$$

$$\text{pH} = 3.489$$

Ex. Calculate the concentration of hydroxide ions in a solution with a pOH of 10.14.

$$[\text{OH}^-_{(\text{aq})}] = 10^{-\text{pOH}}$$

$$[\text{OH}^-_{(\text{aq})}] = 10^{-10.14}$$

$$[\text{OH}^-_{(\text{aq})}] = 7.2 \times 10^{-11}\text{M}$$


$$10^x$$

$$10^{(-10.14)}$$

Strong Acids

Calculate the concentration of the hydroxide ions, pH and pOH of a 0.15 mol/L solution of hydrochloric acid at 25°C.

Strong acids will always completely ionize



$$\begin{aligned} [\text{OH}_{(\text{aq})}^{-}] &= ? \\ \checkmark \text{pH} &= ? \\ \text{pOH} &= ? \end{aligned}$$
$$\begin{aligned} \text{pH} &= -\log [\text{H}_{(\text{aq})}^{+}] \\ \text{pH} &= -\log [0.15] \end{aligned}$$

pH = 0.82

$$K_w = [\text{H}_{(\text{aq})}^{+}][\text{OH}_{(\text{aq})}^{-}]$$
$$[\text{OH}_{(\text{aq})}^{-}] = \frac{K_w}{[\text{H}_{(\text{aq})}^{+}]}$$
$$[\text{OH}_{(\text{aq})}^{-}] = \frac{1.0 \times 10^{-14}}{0.15}$$

$[\text{OH}_{(\text{aq})}^{-}] = 6.7 \times 10^{-14} \text{ M}$

$$\begin{aligned} \text{pOH} &= -\log [\text{OH}_{(\text{aq})}^{-}] \\ \text{pOH} &= -\log [6.7 \times 10^{-14}] \end{aligned}$$

pOH = 13.18

