

Correct Homework Sheet

⑥ $\text{d}, \quad G(x) = (x^4 - x + 1)^3 (x^3 - 2)^2$

$$\begin{aligned} G'(x) &= (x^4 - x + 1)^3 (3)(x^3 - 2)^2(3x) + 2(x^4 - x + 1)(4x^3 - 1)(x^3 - 2)^3 \\ &= 6x(x^4 - x + 1)^3 (x^3 - 2)^3 + 2(x^4 - x + 1)(x^3 - 2)^3 (4x^3 - 1) \\ &= 2(x^4 - x + 1)(x^3 - 2)^3 [3x^5 - 3x^3 + 3x + 4x^5 - x^3 - 8x^3 + 2] \\ &= 2(x^4 - x + 1)(x^3 - 2)^3 (7x^5 - 8x^3 + 3x + 2) \end{aligned}$$

⑦ $y - y_1 = m(x - x_1)$

Point: $(2, \frac{1}{2})$ $y = \frac{1}{\sqrt{20-x^4}} = \frac{1}{(20-x^4)^{\frac{1}{2}}} = (20-x^4)^{-\frac{1}{2}}$

$m = \text{slope} = y'(2)$ $y' = -\frac{1}{2}(20-x^4)^{-\frac{3}{2}}(-4x^3)$

$y'(2) = \frac{2(2)}{\sqrt{(20-2^4)^3}} = \frac{2(2)}{\sqrt{64}} = \frac{4}{8} = \frac{1}{2}$

$$y' = \frac{\partial x^3}{(20-x^4)^{\frac{3}{2}}}$$

$$= \frac{16}{\sqrt{64}} \quad y' = \frac{\partial x^3}{\sqrt{(20-x^4)^3}}$$

m = 2

(ii) $y - y_1 = m(x - x_1)$

$$y - \frac{1}{2} = 2(x - 2)$$

$$y - \frac{1}{2} = 2x - 4 \quad \rightarrow 2x - y - \frac{7}{2} = 0$$

$$2y - 1 = 4x - 8$$

0 = 4x - 2y - 7

⑧ $g(2) = 4$

$$g'(2) = 3$$

$$f'(4) = 5$$

$$F'(2) = ?$$

$$F(x) = f(g(x))$$

$$F'(x) = f'(g(x))g'(x)$$

$$F'(2) = f'(g(2))g'(2)$$

$$= [f'(4)][g'(2)]$$

$$= (5)(3)$$

$$= 15$$

Correct Homework Sheet

$$\textcircled{2} \text{ b) } y = \sqrt[3]{\frac{1-x^6}{2+(5x-1)^4}} = \left[\frac{1-x^6}{2+(5x-1)^4} \right]^{\frac{1}{3}} .$$

$$y' = \frac{1}{3} \left[\frac{1-x^6}{2+(5x-1)^4} \right]^{-\frac{2}{3}} \left[\frac{(2+(5x-1)^4)(-6x^5) - (1-x^6)(4)(5x-1)^3(5)}{(2+(5x-1)^4)^2} \right]$$

Correct Homework Sheet


$$③ b) f(x) = \frac{8x^3(12x^3 - 5x)^8}{2 - 3(1 - 32x^{10})^{\frac{1}{5}}}$$

$$\frac{[2 - 3(1 - 32x^{10})^{\frac{1}{5}}] [(8x^2)(8)(12x^3 - 5x)(24x - 5) + (24x^3)(12x^3 - 5x)^8] - [8x^3(12x^3 - 5x)^8] \left[\left(-\frac{3}{5} \right) (1 - 32x^{10})^{-\frac{4}{5}} (-320x^9) \right]}{[2 - 3(1 - 32x^{10})^{\frac{1}{5}}]^2}$$

$$③ c) f(x) = \frac{[x^5 - x(4-x^3)^{\frac{1}{3}}]^6}{12x^{\frac{1}{3}}(5x^3 - 8)}$$

$$f'(x) = \frac{[12x^{\frac{1}{3}}(5x^3 - 8)] [6[x^5 - x(4-x^3)^{\frac{1}{3}}]^5 [5x^4 - x(\frac{1}{3})(4-x^3)^{-\frac{2}{3}}(-2x) + (4-x^3)^{\frac{2}{3}}] - [x^5 - x(4-x^3)^{\frac{1}{3}}]^6 [(12x^{\frac{1}{3}})(7)(5x^3 - 8)^6 (15x^2) + (6x^{-\frac{1}{3}})(5x^3 - 8)]]}{[12x^{\frac{1}{3}}(5x^3 - 8)]^2}$$


To be handed in today
Differentiate the following (do not simplify)



Derivatives of Trigonometric Functions

The Sine Function

- We recall that the derivative $f'(x)$ of a function $f(x)$ gives the slope of the tangent.
- On the next slide we graph $f(x) = \sin x$ together with $f'(x)$, as determined by the slope of the tangent to the sine curve.
 - Note that x is measured in radians.
- The derivative graph resembles the graph of the cosine!



Let's check this using the definition of a derivative...

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x + h) - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\sin x \left(\frac{\cos h - 1}{h} \right) + \cos x \left(\frac{\sin h}{h} \right) \right] \\
 &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h}
 \end{aligned}$$

- Our calculations have brought us to four limits, two of which are easy:
- Since x is constant while $h \rightarrow 0$,

$$\lim_{h \rightarrow 0} \sin x = \sin x \text{ and } \lim_{h \rightarrow 0} \cos x = \cos x$$

- With some work we can also show that

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \text{ and } \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} = 0$$

- Thus our guess is confirmed:

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\
 &= (\sin x) \cdot 0 + (\cos x) \cdot 1 = \cos x
 \end{aligned}$$

Rules to differentiate trigonometric functions:

Given that "u" represents some differentiable function...

$$\frac{d}{du}(\sin u) = \cos u \bullet du$$

$$\frac{d}{du}(\csc u) = -\csc u \cot u \bullet du$$

$$\frac{d}{du}(\cos u) = -\sin u \bullet du$$

$$\frac{d}{du}(\sec u) = \sec u \tan u \bullet du$$

$$\frac{d}{du}(\tan u) = \sec^2 u \bullet du$$

$$\frac{d}{du}(\cot u) = -\csc^2 u \bullet du$$

Let's Practice...

Differentiate the following:

$$y = \sin 3x$$

$$y = \sin(x + 2)$$

$$y = \sin(kx + d)$$

Ex #2.

Differentiate:

a) $y = \sin(x^3)$

b) $y = \sin^3 x$

c) $y = \sin^3(x^2 - 1)$

Ex #3.

Differentiate:

$$y = x^2 \cos x$$

Homework

Worksheet on derivatives of trigonometric functions

Attachments

Derivatives Worksheet.doc