

Questions from Homework

$$\textcircled{4} \quad f(x) = \sqrt{1+x^3} = (1+x^3)^{1/2}$$

$$f'(x) = \frac{1}{2}(1+x^3)^{-1/2} (3x^2)$$

$$= \frac{3x^2}{2(1+x^3)^{1/2}}$$

$$f''(x) = \frac{2(1+x^3)^{1/2}(6x) - 3x^2(1)(1+x^3)^{-1/2}(3x^2)}{[2(1+x^3)^{1/2}]^2}$$

$$= \frac{12x(1+x^3)^{1/2} - 9x^4(1+x^3)^{-1/2}}{4(1+x^3)}$$

$$= \frac{3x(1+x^3)^{-1/2} [4(1+x^3) - 3x^3]}{4(1+x^3)}$$

$$= \frac{3x(1+x^3)^{-1/2}(4+x^3)}{4(1+x^3)}$$

$$= \frac{3x(4+x^3)}{4(1+x^3)(1+x^3)^{1/2}} = \frac{3x(4+x^3)}{4(1+x^3)^{3/2}}$$

$$f''(2) = \frac{3(2)[4+(2)^3]}{4\sqrt{(1+(2)^3)^3}}$$

$$= \frac{6(12)}{4\sqrt{128}}$$

$$= \frac{72}{4(27)}$$

$$= \frac{72}{108}$$

$$= \boxed{\frac{2}{3}}$$

⑧ Quadratic function

$$f(3) = 33$$

$$f(x) = 4x^2 - 2x + 3$$

$$f'(3) = 22$$

$$f'(x) = 8x - 2$$

$$f''(3) = 8$$

$$f''(x) = 8$$

$$⑤ \quad g(x) = \frac{1}{\sqrt{3x+4}} = \frac{1}{(3x+4)^{1/2}} = (3x+4)^{-1/2}$$

$$g'(x) = -\frac{1}{2} (3x+4)^{-3/2} (3) = -\frac{3}{2} (3x+4)^{-3/2}$$

$$g''(x) = \frac{9}{4} (3x+4)^{-5/2} (3) = \frac{27}{4} (3x+4)^{-5/2}$$

$$g'''(x) = -\frac{135}{8} (3x+4)^{-7/2} (3)$$

$$g'''(x) = \frac{-405}{8(3x+4)^{7/2}}$$

$$g'''(x) = \frac{-405}{8\sqrt{(3x+4)^7}}$$

$$g'''(4) = \frac{-405}{8\sqrt{(3(4)+4)^7}}$$

$$= \frac{-405}{8\sqrt{(16)^7}}$$

$$= \frac{-405}{131072}$$

Implicit Differentiation

So far we have described functions by expressing one variable *explicitly* in terms of another variable: for example,

$$y = x^2 \quad \frac{dy}{dx} = 2x$$
$$y' = 2x$$

■ Sometimes an equation only implicitly defines y as a function (or functions) of x .

■ Examples


■ $x^2 + y^2 = 25$

■ $x^3 + y^3 = 6xy$

$$x^2 + y^2 = 25$$
$$y^2 = 25 - x^2$$
$$y = \pm \sqrt{25 - x^2}$$

- The first equation could easily be rearranged for $y = \dots$

$$y = \pm \sqrt{25 - x^2} \quad \leftarrow \text{Actually gives two functions}$$



Implicit Differentiation

- There is a way called *implicit differentiation* to find dy/dx without solving for y :
 - First differentiate both sides of the equation with respect to x ;
 - Then solve the resulting equation for y' or $\frac{dy}{dx}$
- We will always assume that the given equation does indeed define y as a differentiable function of x .

Example

- For the circle $x^2 + y^2 = 25$, find
 - a) dy/dx or y' or (Slope of the tangent)
 - b) an equation of the tangent at the point $(3, 4)$.

Solution:

Start by differentiating both sides of the equation:

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}(25)$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0$$

Remembering that y is a function of x and using the Chain Rule, we have

$$2x + 2yy' = 0$$

Thus...

$$2x + 2y \frac{dy}{dx} = 0$$

Solving for $\frac{dy}{dx}$...

$$2yy' = -2x$$

$$y' = \frac{-2x}{2y}$$

$$y' = \frac{-x}{y} = \frac{-3}{4} = m$$

Therefore at the point $(3, 4)$ the equation of the tangent would be...

$$y - 4 = -\frac{3}{4}(x - 3)$$

or

$$3x + 4y = 25$$

$$y - 4 = -\frac{3}{4}x + \frac{9}{4}$$

$$4y - 16 = -3x + 9$$

$$3x + 4y - 25 = 0$$

Same Example Revisited

- Since it is easy to solve this equation for y , we
 - do so, and then
 - find the equation of the tangent line by earlier methods, and then
 - compare the result with our preceding answer:

Solution

- Solving the equation gives $y = \pm\sqrt{25-x^2}$ as before.
- The point $(3, 4)$ lies on the upper semicircle $y = \sqrt{25-x^2}$ and so we consider the function $f(x) = \sqrt{25-x^2}$

Differentiate f : $y = \sqrt{25-x^2} = (25-x^2)^{1/2}$

$$y' = \frac{1}{2}(25-x^2)^{-1/2}(-2x)$$
$$y' = \frac{-x}{\sqrt{25-x^2}} = \frac{-(3)}{\sqrt{25-(3)^2}} = \frac{-3}{4}$$

Equation:

$$y-4 = \frac{-3}{4}(x-3)$$

$$y-4 = \frac{-3x+9}{4}$$

$$4y-16 = -3x+9$$

$$\boxed{3x+4y+25=0}$$

Solution (cont'd)

- So $f'(3) = -\frac{3}{\sqrt{25-3^2}} = -\frac{3}{4},$

leading to the same equation

$$3x + 4y = 25$$

for the tangent that we obtained earlier.

- Note that although this problem could be done both ways, implicit differentiation was easier!

Sometimes **Implicit Differentiation** is not only the easiest way, it's the *only* way

Example:

Given

$$x^3 + y^3 = 6xy$$

first second

Find $\frac{dy}{dx}$

$$3x^2 + 3y^2 y' = 6xy' + 6y$$

Product Rule

$$3y^2 y' - 6xy' = -3x^2 + 6y$$

Factor

$$y'(3y^2 - 6x) = -3x^2 + 6y$$

$$y' = \frac{-3x^2 + 6y}{3y^2 - 6x}$$

$$y' = \frac{-\cancel{3}(x^2 - 2y)}{\cancel{3}(y^2 - 2x)}$$

$$y' = -\frac{x^2 - 2y}{y^2 - 2x}$$

Find $\frac{dy}{dx}$

Product Rule

$$2x^5 + x^4y + y^5 = 36$$

$$10x^4 + x^4y' + 4x^3y + 5y^4y' = 0$$

$$x^4y' + 5y^4y' = -10x^4 - 4x^3y$$

Factor $y'(x^4 + 5y^4) = -10x^4 - 4x^3y$

$$y' = \frac{-10x^4 - 4x^3y}{x^4 + 5y^4}$$

$$y' = -\frac{10x^4 + 4x^3y}{x^4 + 5y^4}$$

Homework

$$\textcircled{1} \text{ g) } \sqrt{x} + \sqrt{y} = 1$$

$$x^{1/2} + y^{1/2} = 1$$

$$\frac{1}{2}x^{-1/2} + \frac{1}{2}y^{-1/2}y' = 0$$

$$\frac{1}{2\sqrt{x}} + \frac{y'}{2\sqrt{y}} = 0$$

$$\frac{y'}{2\sqrt{y}} = -\frac{1}{2\sqrt{x}}$$

$$y' = -\frac{\sqrt{y}}{\sqrt{x}}$$

$$\boxed{y' = -\frac{\sqrt{y}}{\sqrt{x}}}$$

$$\text{h) } \frac{\partial x}{x+y} = y$$

$$\frac{(x+y)(2) - \partial x(1+y')}{(x+y)^2} = y'$$

$$\frac{2x+2y-\partial x-\partial xy'}{(x+y)^2} = y'$$

$$y'(x+y)^2 = 2y - \partial xy'$$

$$y'(x+y)^2 + \partial xy' = 2y$$

factor $y'[(x+y)^2 + \partial x] = 2y$

$$\boxed{y' = \frac{2y}{(x+y)^2 + \partial x}}$$

#3

$$y = \sqrt[7]{2x^2 + \sqrt{x^7 - 8x\sqrt{3-x^3}}}$$

$$= \left[2x^2 + (x^7 - 8x(3-x^3)^{1/2})^{1/2} \right]^{1/7}$$

$$y' = \frac{1}{7} \left[2x^2 + (x^7 - 8x(3-x^3)^{1/2})^{1/2} \right]^{-6/7} \left[4x + \frac{1}{2}(x^7 - 8x(3-x^3)^{1/2})^{-1/2} (7x^6 - [8x(\frac{1}{2})(3-x^3)^{-1/2}(-3x^2) - 8(3-x^3)^{1/2}]) \right]$$

$$\textcircled{4} \quad y = \frac{(2x+3)^3}{\sqrt{4x-7}} = \frac{(2x+3)^3}{(4x-7)^{1/2}} \quad \begin{array}{l} x=2 \quad (2, 343) \\ y=343 \end{array} \quad \begin{array}{l} \uparrow \\ \text{point} \end{array}$$

$$y = \frac{(2(2)+3)^3}{\sqrt{4(2)-7}} \quad y' = \frac{(4x-7)^{1/2} (3)(2x+3)^2 (2) - (2x+3)^3 (\frac{1}{2})(4x-7)^{-1/2} (4)}{4x-7}$$

$$= \frac{7^3}{\sqrt{1}}$$

$$= 343$$

$$y'(2) = \frac{(1)(3)(49)(2) - (343)(\frac{1}{2})(1)(4)}{1}$$

$$= 294 - 686$$

$$= -392 \quad \leftarrow \text{Slope}$$

$$y - y_1 = m(x - x_1)$$

$$y - 343 = -392(x - 2)$$

$$y - 343 = -392x + 784$$

$$\boxed{392x + y - 1127 = 0}$$