



Homework - Total Energy



$$\Delta E_T = q_l + \Delta H_{\text{vap}} + q_s$$

$$q_l = m C \Delta T$$

$$q_l = (200000 \text{ g}) \left(4.19 \frac{\text{J}}{\text{g} \cdot \text{C}} \right) (80.0 \text{ } ^\circ\text{C})$$

$$q_l = 67040000 \text{ J}$$

$$\Delta H_{\text{vap}} = n H_{\text{vap}}$$

$$\Delta H_{\text{vap}} = \left(\frac{200000 \text{ g}}{18.02 \text{ g/mol}} \right) \left(40.8 \frac{\text{kJ}}{\text{mol}} \right)$$

$$\Delta H_{\text{vap}} = 452830.19 \text{ kJ}$$

$$q_s = m C \Delta T$$

$$q_s = (500000 \text{ g}) \left(0.528 \frac{\text{J}}{\text{g} \cdot \text{C}} \right) (80.0 \text{ } ^\circ\text{C})$$

$$q_s = 21120000 \text{ J}$$

$$\Delta E_T = (67040 \text{ kJ}) + (452830 \text{ kJ}) + (21120 \text{ kJ})$$

$$\boxed{\Delta E_T = 541000 \text{ kJ}}$$

Calorimetry

CALORIMETRY - is the technological process of measuring energy changes using an **isolated system** called a calorimeter.

In the calorimeter the system being studied is surrounded by a known quantity of water. Energy is then transferred between the chemical system and the water. The heat gained by the water can be determined and thus equals the heat lost by the system.


ASSUMPTIONS IN CALORIMETRY

1. no heat is transferred between the calorimeter and the outside environment.
2. any heat absorbed or released by the calorimeter materials is negligible.
3. a dilute aqueous solution has the same density and specific heat capacity as pure water.

Assumption #2 implies

$$\Delta H_{\text{system}} = - q_{\text{calorimeter}}$$

NaCl
3.0g



100mL
19°C

24°C

Example

4.24 g of lithium chloride is dissolved in 100. mL of water at an initial temperature of 16.3°C. The final temperature of the solution is 25.1°C.

Calculate the molar enthalpy of solution, H_s , for lithium chloride.

LiCl

$$\Delta H_s = -q_{\text{H}_2\text{O}}$$
$$nH_s = -vC\Delta T$$

$H_s = ?$

$$\left(\frac{4.24\text{g}}{42.39\text{g/mol}}\right)H_s = -\left(0.100\text{L}\right)\left(4.19\frac{\text{kJ}}{\text{L}\cdot^\circ\text{C}}\right)\left(8.8^\circ\text{C}\right)$$

H₂O

$$H_s = \frac{-\left(0.100\text{L}\right)\left(4.19\frac{\text{kJ}}{\text{L}\cdot^\circ\text{C}}\right)\left(8.8^\circ\text{C}\right)}{\left(\frac{4.24\text{g}}{42.39\text{g/mol}}\right)}$$

$v = 100.\text{mL}$

$T_i = 16.3^\circ\text{C}$

$T_f = 25.1^\circ\text{C}$

$$H_s = -36.9\frac{\text{kJ}}{\text{mol}}$$

A block of aluminum at 95.0°C is placed into $500. \text{ g}$ of water at 21.5°C . The final temperature of the water and aluminum is 28.0°C . Determine the mass of the aluminum.

<u>Al</u>	<u>H₂O</u>
$T_i = 95.0^{\circ}\text{C}$	$m = 500. \text{ g}$
$m = ?$	$T_i = 21.5^{\circ}\text{C}$

$$T_f = 28.0^{\circ}\text{C}$$

$$q_{\text{Al}} = -q_{\text{H}_2\text{O}}$$

$$m_{\text{Al}} C \Delta T = -m_{\text{H}_2\text{O}} C \Delta T$$

$$m_{\text{Al}} \left(0.900 \frac{\text{J}}{\text{g}^{\circ}\text{C}}\right) (28.0^{\circ}\text{C} - 95.0^{\circ}\text{C}) =$$

$$- (500. \text{ g}) \left(4.19 \frac{\text{J}}{\text{g}^{\circ}\text{C}}\right) (28.0^{\circ}\text{C} - 21.5^{\circ}\text{C})$$

$$m_{\text{Al}} = - \frac{(500. \text{ g}) \left(4.19 \frac{\text{J}}{\text{g}^{\circ}\text{C}}\right) (6.5^{\circ}\text{C})}{\left(0.900 \frac{\text{J}}{\text{g}^{\circ}\text{C}}\right) (-67.0^{\circ}\text{C})}$$

$$m_{\text{Al}} = 226 \text{ g}$$

Homework

Quiz

Pre-Lab

Find the molar mass of the following substances:



www.indigo.com
Erlenmeyer Flasks