
# **Chemical Bonding**




#### **Valence electrons**

electrons in the highest occupied energy level of an element's atoms.

- determines the chemical properties of an element
- only electrons used in chemical bonds
- for a representative element, the number of valence electrons corresponds to the group number

#### **Electron dot structure**

diagrams showing the valence electrons as dots





## Table 7.1

Electron Dot Structure of Some Group A Elements								
	Group							
Period	1A	2 <b>A</b>	<b>3A</b>	4A	5 <b>A</b>	6 <b>A</b>	7 <b>A</b>	<b>8</b> A
1	H.							He:
2	Li-	·Be·	·B·	Ċ	N	Ö	· E	Ne
3	Na <sup>.</sup>	·Mg·	Al	Si	.P.	S	CI	Ar
4	K.	Ca	Ga	Ge	As	Se	Br	:Kr

### **Octet Rule**

To form compounds, atoms usually achieve the electron configuration of a noble gas.

At the highest occupied energy level: ns<sup>2</sup>np<sup>6</sup>


Formation of Cations

Cations lose valence electrons to form positively charged ions

Na 
$$\longrightarrow$$
 Na<sup>+</sup> + e<sup>-</sup>

$$Mg \longrightarrow Mg^{2+} + 2e^{-}$$

Transition Metals will attempt to form a pseudo noble-gas configuration.



Ne 15<sup>2</sup>25<sup>2</sup>2p<sup>6</sup>
Ar 15<sup>2</sup>25<sup>2</sup>2p<sup>6</sup>35<sup>2</sup>3p<sup>6</sup>

152252pb3,23p

# Formation of Anions Anions gain electrons to produce a negatively charged ion.

CI 
$$1s^22s^22p^63s^23p^5 \xrightarrow{+e^-}$$
 CI  $1s^22s^22p^63s^23p^6$ 

### Ionization:

$$0 + 2e^{-} \longrightarrow O^{2-}$$

# **Homework**

p. 193 #3-11