

Warm Up

Review of laws of logarithms...

Express the following as a single logarithm:

$$\frac{2}{3}\ln b^6 - \frac{1}{2}\left[\ln b^4 + 8\ln b + 6\ln\sqrt[3]{b}\right]$$

Questions from Homework

③ d) $G(x) = \frac{1 + \log_3 x}{x}$

$$G'(x) = \frac{x \left(\frac{1}{x \ln 3} \right) - (1 + \log_3 x)}{x^2}$$

$$= \frac{\frac{1}{\ln 3} - 1 - \log_3 x}{x^2 \ln 3}$$

$$= \frac{1 - \ln 3 - \ln 3 (\log_3 x)}{x^2 \ln 3}$$

⑦ $f(x) = x \ln x$

i) $f'(x) = x \left(\frac{1}{x} \right) + (1) \ln x$

$f'(x) = 1 + \ln x$


$1 + \ln x = 0$

$\ln x = -1$

$e^{-1} = x$

cv:

$\frac{1}{e} = x$



$f\left(\frac{1}{e}\right) = \left(\frac{1}{e}\right) \ln\left(\frac{1}{e}\right) \rightarrow \ln e^{-1}$

$= \frac{1}{e}(-1)$

$= -\frac{1}{e}$


$\left(\frac{1}{e}, -\frac{1}{e}\right)$ Min

(ii) $f'(x) = 1 + \ln x$

$f''(x) = 0 + \frac{1}{x}$

$f''(x) = \frac{1}{x}$

cv: $x = 0$



CU on $(0, \infty)$
CD on $(-\infty, 0)$

⑧ $f(x) = x (\ln x)^2$

$f'(x) = x (2) (\ln x) \left(\frac{1}{x} \right) + (1) (\ln x)^2$

$f'(x) = 2 \ln x + (\ln x)^2$

$f'(x) = \ln x [2 + \ln x]$

$0 = \ln x [2 + \ln x]$

cv: $\ln x = 0$ | $2 + \ln x = 0$


$e^0 = x$

$1 = x$

$\ln x = -2$

$e^{-2} = x$

$\frac{1}{e^2} = x$



$f\left(\frac{1}{e}\right) = \frac{4}{e^3}$ max

$f(1) = 0$ min

Logarithmic Differentiation

A differentiation process that requires taking the logarithm of both sides before differentiating.

This process will be used in TWO circumstances:

I. Simplifying messy products and quotients

What would it involve to differentiate the following?

$$y = \frac{(x^2 - 1)^5 \sqrt{2x+9} (5x^3 + 2)^8}{(10x-1)\sqrt{5-x^7}}$$

- Quotient rule, multiple product rules and chain rules...

This would be possible but it would be easier to differentiate a group of terms added and subtracted rather than multiplied and divided

Laws of logarithms will do exactly that...turn this mess into a addition and subtraction of terms.

$$y = \frac{(x^2 - 1)^5 \sqrt{2x+9} (5x^3 + 2)^8}{(10x-1)\sqrt{5-x^7}}$$

$$\ln y = \ln \left[\frac{(x^2-1)^5 (2x+9)^{1/2} (5x^3+2)^8}{(10x-1)(5-x^7)^{1/2}} \right]$$

$$\ln y = \ln(x^2-1)^5 + \ln(2x+9)^{1/2} + \ln(5x^3+2)^8 - \ln(10x-1) - \ln(5-x^7)^{1/2}$$

$$\ln y = 5 \ln(x^2-1) + \frac{1}{2} \ln(2x+9) + 8 \ln(5x^3+2) - \ln(10x-1) - \frac{1}{2} \ln(5-x^7)$$

$$\frac{y'}{y} = \left[5 \left(\frac{2x}{x^2-1} \right) + \frac{1}{2} \left(\frac{2}{2x+9} \right) + 8 \left(\frac{15x^2}{5x^3+2} \right) - \left(\frac{10}{10x-1} \right) - \frac{1}{2} \left(\frac{-7x^6}{5-x^7} \right) \right] y$$

$$y' = \left[\frac{10x}{x^2-1} + \frac{1}{2x+9} + \frac{120x^2}{5x^3+2} - \frac{10}{10x-1} + \frac{7x^6}{2(5-x^7)} \right] \left[\frac{(x^2-1)^5 \sqrt{2x+9} (5x^3+2)^8}{(10x-1)\sqrt{5-x^7}} \right]$$

Steps in Logarithmic Differentiation

1. Take logarithms of both sides of an equation.
2. Differentiate implicitly with respect to x .
3. Solve the resulting equation for y'

Use Logarithmic Differentiation to Differentiate the following:

$$y = \frac{e^x \sqrt{x^2 + 1}}{(x^2 + 2)^3}$$

$$\ln y = \ln \left[\frac{e^x (x^2 + 1)^{1/2}}{(x^2 + 2)^3} \right]$$

$$\ln y = \underline{\ln e^x} + \ln(x^2 + 1)^{1/2} - \ln(x^2 + 2)^3$$

$$\ln y = x + \frac{1}{2} \ln(x^2 + 1) - 3 \ln(x^2 + 2)$$

$$y \cdot \frac{y'}{y} = \left[1 + \frac{1}{2} \left(\frac{2x}{x^2 + 1} \right) - 3 \left(\frac{2x}{x^2 + 2} \right) \right] y$$

$$y' = \left[1 + \frac{x}{x^2 + 1} - \frac{6x}{x^2 + 2} \right] \left[\frac{e^x \sqrt{x^2 + 1}}{(x^2 + 2)^3} \right]$$

II. Base and exponent both variables

Have a look at this example:

$$y = x^{x^5}$$

- Does not fit either the power rule or the rules for an exponential function

...What can be done to help this crazy situation??

Of Course...take the logarithm of both sides!!

$$\begin{aligned}y &= x^{x^5} \\ \ln y &= \ln x^{x^5} \\ \ln y &= x^5 \ln x \\ \frac{y'}{y} &= x^5 \left(\frac{1}{x}\right) + 5x^4 \ln x \\ y \cdot \frac{y'}{y} &= [x^4 + 5x^4 \ln x] y \\ y' &= [x^4 + 5x^4 \ln x] x^{x^5}\end{aligned}$$

Example:

Differentiate: $y = (\ln x^5)^{\cos x}$

Practice Questions...

Page 395–396

#1 b, d, e

#2 b, c, e

#3