# **Gravimetric Stoichiometry**

Gravimetric - refers to mass measurement

Stoichiometric - refers to the procedure used to calculate quantities of chemicals.

GRAVIMETRIC STOICHIOMETRY - the procedure for calculating the masses of reactants and products in a chemical reaction.

#### ASSUMPTIONS IN STOICHIOMETRY

- the reaction is spontaneous.
- the reaction is fast
- the reaction is quantitative.
- the reaction is stoichiometric.

(there is a whole number ratio between MOLES of reactant and MOLES of product.)

$$2Hz + Oz \rightarrow 2HzO$$
  
 $m=100g$  ? 9  
 $x?$  50.9  
? 259

Iron can be produced by the reaction of iron (III) oxide, with carbon monoxide to produce iron metal and carbon dioxide. What mass of iron (III) oxide is required to produce 1000. g of iron?

$$Fe_2O_{3(s)} + 3CO_{(g)}$$
  $Fe_{(s)} + 3CO_{2(g)}$   $m = 1000.9$ 

### Step 1: # moles known

#### Step 2: moles unknown

### Step 3: mass unknown

## **Try This**

A)3+ O2-

What mass of aluminum oxide will be produced if 17.0 g of oxygen reacts with excess aluminum?

$$44l_{cs} + 30_{2cg} \longrightarrow 24l_{2}O_{3(5)}$$
 $m = 17.00g$ 
 $m = ?$ 

Step 1: Moles Known  $17.09 O_2 \times \frac{1 \text{ mol } O_2}{32.009 O_2} = 0.53125 \text{ mol } O_2$ 

Step 2! Moles Unknown

0.53125 mol O2 x 2 mol Al2O3 0.35416 mol Al2O3

3 mol O2

Step 3: Mass Un Known

0.35416 mol Al2O3 x 101.969 Al2O3 = 36.19 Al2O3

## Worksheet

$$1) Zn_{(s)} + S_{8(s)} \Rightarrow ZnS_{(s)}$$

$$2) \qquad Al_2O_{3(s)} \quad \Rightarrow \quad Al_{(s)} + O_{2(g)}$$

3) 
$$C_3H_{8(I)} + O_{2(g)} \Rightarrow CO_{2(g)} + H_2O_{(g)}$$

4) 
$$NaCl_{(aq)} + Pb(NO_3)_{2(aq)} \Rightarrow NaNO_{3(aq)} + PbCl_{2(s)}$$

5) 
$$AI_{(s)} + H_2SO_{4(aq)} \Rightarrow H_{2(g)} + AI_2(SO_4)_{3(aq)}$$

6) 
$$KOH_{(aq)} + Cu(NO_3)_{2(aq)} \Rightarrow KNO_{3(aq)} + Cu(OH)_{2(s)}$$