

Questions from Homework

$$\textcircled{3} \text{ c) } \lim_{x \rightarrow 4} \frac{(x-4)(\sqrt{x}+2)}{(\sqrt{x}-2)(\sqrt{x}+2)}$$

$$\lim_{x \rightarrow 4} \frac{\cancel{(x-4)}(\sqrt{x}+2)}{\cancel{(x-4)}} = \boxed{4}$$

$$\text{d) } \lim_{x \rightarrow 0} \frac{\cancel{4(2+x)^2} \cdot \frac{1}{\cancel{(2+x)^2}} - \frac{1}{4} \cdot \cancel{(4)}(2+x)^2}{x(4)(2+x)^2}$$

$$\text{CD.} = \frac{(4)(2+x)^2}{(4)(2+x)^2}$$

$$\lim_{x \rightarrow 0} \frac{4 - (2+x)^2}{4x(2+x)^2} \quad \leftarrow \text{D.f.f. of Squares}$$


$$\lim_{x \rightarrow 0} \frac{(2 - (2+x))(2 + (2+x))}{4x(2+x)^2}$$

$$\lim_{x \rightarrow 0} \frac{(-x)(4+x)}{4x(2+x)^2} = \frac{-4}{16} = \boxed{-\frac{1}{4}}$$

$$\hookrightarrow \frac{(-1)(4+0)}{4(1)(2+0)^2}$$

$$\textcircled{2} \quad f(x) = \begin{cases} |x| & x < 0 \\ 3 & 0 \leq x \leq 3 \\ (x-3)^2 + 1 & x > 3 \end{cases}$$

x	$ x $	x	$f(x)$	x	$f(x)$
0	0	0	3	3	1
-1	1	1	3	4	2
-2	2	2	3	5	5
-3	3	3	3	6	10



Discontinuous at $x = 0, 3$

$$f(3) = 3$$

$$\lim_{x \rightarrow 3} f(x) = \text{DNE}$$

Limits at Infinity



What exactly is infinity?

- It is the *process* of making a value arbitrarily large or small

$+\infty \longrightarrow$ Positive Infinity...process of becoming arbitrarily large

$-\infty \longrightarrow$ Negative Infinity...process of becoming arbitrarily small

4 Definition Let f be a function defined on some interval (a, ∞) . Then

$$\lim_{x \rightarrow \infty} f(x) = L$$

means that the values of $f(x)$ can be made as close to L as we like by taking x sufficiently large.






FIGURE 9


Examples illustrating $\lim_{x \rightarrow \infty} f(x) = L$

Have a look at these limits...

$$\lim_{x \rightarrow \infty} \frac{1}{x} = 0$$

$$\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$$

"As x gets larger
 y gets closer and
closer to 0"



In general...

7 If n is a positive integer, then

$$\lim_{x \rightarrow \infty} \frac{1}{x^n} = 0$$

$$\lim_{x \rightarrow -\infty} \frac{1}{x^n} = 0$$

Calculating limits at infinity without using a graph

• Rational Functions

Note: If every term in a rational expression is divided by the same value, the rational expression will still be equal to its original value

$$\frac{12+8}{6-2} = \frac{20}{4} = 5 \quad \xrightarrow{\text{Divide the numerator and denominator by 2}} \quad \frac{6+4}{3-1} = \frac{10}{2} = 5$$

This will be important when evaluating limits for rational functions approaching infinity...

Look at the following example:


$$\lim_{x \rightarrow \infty} \frac{3x^2 - x - 2}{5x^2 + 4x + 1} = \lim_{x \rightarrow \infty} \frac{\frac{3x^2 - x - 2}{x^2}}{\frac{5x^2 + 4x + 1}{x^2}} = \lim_{x \rightarrow \infty} \frac{3 - \frac{1}{x} - \frac{2}{x^2}}{5 + \frac{4}{x} + \frac{1}{x^2}}$$

$$= \frac{3 - 0 - 0}{5 + 0 + 0} = \frac{3}{5} = 0.6$$

Divide every term by the **HIGHEST** power that is present in either the numerator or denominator of the rational expression once they are expanded

* For limits at infinity compare the degree of the numerator with the degree of the denominator

This graph below validates our solution:



Calculating limits at infinity without using a graph

● Remember

- If the highest degree is in the denominator then the *Limit* will be equal to 0
- If the highest degree is in the numerator then the *Limit* will not exist.
- If the degree is the same in the numerator and denominator then the *Limit* will be equal to the coefficients in front of the highest degree.

Evaluate the following limit:

$$\lim_{n \rightarrow \infty} \frac{n^2 - n}{2n^2 + 1} = \frac{1}{2}$$

$$\lim_{n \rightarrow \infty} \frac{1 - n^5}{1 + 2n^5} = -\frac{1}{2}$$

$$\lim_{n \rightarrow \infty} \frac{4n}{1} = \text{DNE}$$

$$\lim_{x \rightarrow \infty} \frac{-3(x^2 - 4)^2}{3 - 5x^2}$$

$$\lim_{x \rightarrow \infty} \frac{-3(x^4 - 8x^2 + 16)}{3 - 5x^2}$$

$$\text{or } \lim_{x \rightarrow \infty} \frac{-3x^4 + 24x^2 - 48}{3 - 5x^2} = \text{DNE}$$

Homework