

1) 31 200 J

7) 0.0456 J/g °C

2) -31 700 J

8) 424 g

3) 120°C

9) 2.60 J/g °C

4) 28°C

10) 6.21 J

5) 1100 J

11) 42.6 L

6) 14 900 J

Homework - Worksheet

$$\textcircled{3} m = 50.0 \text{ g}$$

$$T_i = 140^\circ\text{C}$$

$$q = -2.5 \text{ kJ}$$

$$C = 2.01 \frac{\text{J}}{\text{g}\cdot^\circ\text{C}}$$

$$T_f = ?$$

$$q = mC\Delta T$$

$$q = mC(T_f - T_i)$$

$$-2500 \text{ J} = (50.0 \text{ g})(2.01 \frac{\text{J}}{\text{g}\cdot^\circ\text{C}})(T_f - 140^\circ\text{C})$$

$$\frac{-2500 \text{ J}}{(50.0 \text{ g})(2.01 \frac{\text{J}}{\text{g}\cdot^\circ\text{C}})} = T_f - 140^\circ\text{C}$$

$$-24.88^\circ\text{C} = T_f - 140^\circ\text{C}$$

$$T_f = 140^\circ\text{C} - 24.88^\circ\text{C}$$

$$T_f = 115.12^\circ\text{C}$$

$$T_f = 120^\circ\text{C}$$

⑪

$$q = v C \Delta T$$

$\frac{\text{kJ}}{\text{L} \cdot ^\circ\text{C}}$

volume
(L)

⑪ $\Delta T = -65.0^\circ\text{C}$

$q = -332 \text{ kJ}$

$V = ?$

$C = 0.0012 \frac{\text{kJ}}{\text{L} \cdot ^\circ\text{C}}$

$$q = v C \Delta T$$

$$-332 \text{ kJ} = v (0.0012 \frac{\text{kJ}}{\text{L} \cdot ^\circ\text{C}}) (-65.0)$$

$$v = \frac{-332 \text{ kJ}}{(0.0012 \frac{\text{kJ}}{\text{L} \cdot ^\circ\text{C}}) (-65.0^\circ\text{C})}$$

$$V = 42.6 \text{ L}$$

PHASE CHANGE AND ENTHALPY

Classifying types of systems:


1. Open system - a system where both matter and energy can flow into or out of the system.
2. Closed system - a system where energy is allowed to be transferred into and out but matter cannot be transferred.
3. Isolated system - a system where neither matter nor energy is allowed to enter or leave the system.

ENTHALPY (H) - The total internal (potential) energy and kinetic energy of a system under constant pressure.

⇒ Enthalpy is usually expressed in kJ.

ENTHALPY CHANGE (ΔH) - A change under constant pressure where the surroundings of a system absorb energy or release it to the system.

PHASE CHANGE - is a change in the state of matter without a change in the chemical composition of the system.




⇒ **always involve a change in energy but never involve a change in temperature.**

Question:

- (i) What is the temperature where water just starts boiling?
- (ii) What is the temperature when water is boiling violently?
- (iii) If energy is still going into the water and the temperature is not increasing, where is the energy going?



Consider melting ice to water and then boiling water to steam:



MOLAR ENTHALPY

For any system:

- an exothermic change involves a decrease in enthalpy

⇒ gives off energy to the surroundings

⇒ ΔH is negative.

→ $(g \rightarrow l)$ $(l \rightarrow s)$
condensation, solidification

- an endothermic change involves an increase in enthalpy.

⇒ takes in energy from the surroundings

⇒ ΔH is positive.

→ $(l \rightarrow g)$ $(s \rightarrow l)$
vaporization, fusion

The enthalpies for substances undergoing phase changes have been measured experimentally. (TABLE 17.3 p. 522)

- enthalpies are reported as molar enthalpies and are expressed as kJ/mol.


Endothermic Phase Changes

- the molar enthalpy of fusion (H_{fus}) represents the quantity of heat that the substance absorbs per mole as it changes state from **solid to liquid**.

- the molar enthalpy of vaporization (H_{vap}) represents the quantity of heat that the substance absorbs per mole as it changes state from **liquid to gas**.

Exothermic Phase Changes

- the molar enthalpy of condensation (H_{cond}) represents the quantity of heat that the substance releases per mole as it changes state from **gas to liquid**
- the molar enthalpy of solidification (H_{solid}) represents the quantity of heat that the substance releases per mole as it changes state from **liquid to solid**.



$$\Delta H_{\text{fus}} = - \Delta H_{\text{solid}}$$

$$\Delta H_{\text{vap}} = - \Delta H_{\text{cond}}$$

Example

If 500. g of $\text{CCl}_2\text{F}_2(l)$ is vaporized at SATP, find the enthalpy change of the system ($H_{\text{vap}} = 34.99 \text{ kJ/mol}$).

$$m = 500. \text{ g}$$



$$H_{\text{vap}} = 34.99 \frac{\text{kJ}}{\text{mol}}$$

$$\Delta H_{\text{vap}} = n H_{\text{vap}}$$

$$\Delta H_{\text{vap}} = \left(\frac{500. \text{ g}}{120.91 \text{ g/mol}} \right) \left(34.99 \frac{\text{kJ}}{\text{mol}} \right)$$

$$\Delta H_{\text{vap}} = 145 \text{ kJ}$$



$$(1 \times 12.01) + (2 \times 35.45) + (2 \times 19.00)$$

$$= 120.91 \text{ g/mol}$$

Worksheet

Heat (q)

- change in kinetic energy
- measures transfer of energy when there are temperature changes (heating or cooling)

Enthalpy (H)

- measures potential energy
- change in energy transfer when system is at constant pressure and same initial and final temperatures