

# Understanding Logarithms

## Focus on...

---


- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$
- determining the characteristics of the graph of  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

$(x,y) \rightarrow (y,x)$   
reflection in  
the line  
 $y=x$

For the exponential function  $y = c^x$ , the inverse is  $x = c^y$ . This inverse is also a function and is called a **logarithmic function**. It is written as  $y = \log_c x$ , where  $c$  is a positive number other than 1.

**Logarithmic Form**

**Exponential Form**



Since our number system is based on powers of 10, **logarithms** with base 10 are widely used and are called **common logarithms**. When you write a common logarithm, you do not need to write the base. For example,  $\log 3$  means  $\log_{10} 3$ .

### logarithmic function

- a function of the form  $y = \log_c x$ , where  $c > 0$  and  $c \neq 1$ , that is the inverse of the exponential function  $y = c^x$

### logarithm

- an exponent
- in  $x = c^y$ ,  $y$  is called the logarithm to base  $c$  of  $x$

### common logarithm

- a logarithm with base 10

Write each of the following in logarithmic form


a)  $32 = 2^5$   
 (ans. ↓, exp. ↙, base ↑)  
 $\log_2 32 = 5$

b)  $2^{-5} = \frac{1}{32}$   
 $\log_2 \left(\frac{1}{32}\right) = -5$

c)  $x = 10^y$   
 $\log_{10} x = y$   
 or  $\log x = y$

Logarithmic Form

Exponential Form



↑  
common logarithm

Write each of the following in exponential form

a)  $\log_4 16 = 2$   
 (ans. ↓, exp. ↙, base ↑)  
 $4^2 = 16$

b)  $\log_2 \left(\frac{1}{32}\right) = -5$   
 $2^{-5} = \frac{1}{32}$

c)  $\log 65 = 1.8129$   
 $10^{1.8129} = 65$

## Example 1

### Evaluating a Logarithm

Evaluate.

a)  $\log_7 49$

Let  $x = \log_7 49$

$$x = \log_7 49$$

$$7^x = 49$$

$$7^x = 7^2$$

$$x = 2$$

$$\log_7 49 = 2$$

b)  $\log_6 1$

b)  $x = \log_6 1$

$$6^x = 1$$

$$6^x = 6^0$$

$$x = 0$$

$$\log_6 1 = 0$$

c)  $\log 0.001$

c)  $x = \log 0.001$

$$10^x = 0.001$$

$$10^x = 10^{-3}$$

$$x = -3$$

$$\log 0.001 = -3$$

d)  $x = \log_2 \sqrt{8}$

$$2^x = \sqrt{8}$$

$$2^x = 8^{1/2}$$

$$2^x = (2^3)^{1/2}$$

$$2^x = 2^{3/2}$$

$$x = 3/2$$

$$\log_2 \sqrt{8} = \frac{3}{2}$$

## Example 2

### Determine an Unknown in an Expression in Logarithmic Form

Determine the value of  $x$ .

a)  $\log_5 x = -3$

b)  $\log_x 36 = 2$

c)  $\log_{64} x = \frac{2}{3}$

a)  $\log_5 x = -3$

$$5^{-3} = x$$

$$\left(\frac{1}{5}\right)^3 = x$$

$$\frac{1}{125} = x$$

b)  $\log_x 36 = 2$

$$x^2 = 36$$

$$x = \pm 6$$

$$x = 6$$

Choose  $x = 6$   
because your base  
has to be greater  
than 0

c)  $\log_{64} x = \frac{2}{3}$

$$64^{\frac{2}{3}} = x$$

$$\left(\sqrt[3]{64}\right)^2 = x$$

$$(4)^2 = x$$

$$16 = x$$

## Questions from Homework

$$\textcircled{4} \text{ c) } \log_4 \sqrt[3]{4}$$
$$\log_4 4^{1/3} = x$$
$$\cancel{4}^x = \cancel{4}^{1/3}$$
$$\boxed{x = \frac{1}{3}}$$

$$\text{d) } \log_{\frac{1}{3}} 27$$
$$\log_{\frac{1}{3}} 27 = x$$
$$\left(\frac{1}{3}\right)^x = 27$$
$$(3^{-1})^x = 3^3$$
$$\cancel{3}^{-x} = \cancel{3}^3$$
$$-x = 3$$
$$\boxed{x = -3}$$

**General Properties of Logarithms:**

If  $C > 0$  and  $C \neq 1$ , then...

$$(i) \log_C 1 = 0$$

$$(ii) \log_C C^x = x$$

$$(iii) C^{\log_C x} = x$$

**Did You Know?**

The input value for a logarithm is called an argument. For example, in the expression  $\log_6 1$ , the argument is 1.

$$(i) \log_5 1 = 0$$

$$(ii) \log_2 2^3 = 3$$

$$(iii) 7^{\log_7 49} = 49$$

$$5^{\log_5 10} = 10$$

### Example 3

#### Graph the Inverse of an Exponential Function

- a) State the inverse of  $f(x) = 3^x$ .
- b) Sketch the graph of the inverse. Identify the following characteristics of the inverse graph:
- the domain and range
  - the  $x$ -intercept, if it exists
  - the  $y$ -intercept, if it exists
  - the equations of any asymptotes

a) Find the inverse.

$$f(x) = 3^x$$

①  $y = 3^x$

②  $x = 3^y$

③  $y = \log_3 x$

④  $f^{-1}(x) = \log_3 x$



**Solution**


a) The inverse of  $y = f(x) = 3^x$  is  $x = 3^y$  or, expressed in logarithmic form,  $y = \log_3 x$ . Since the inverse is a function, it can be written in function notation as  $f^{-1}(x) = \log_3 x$ .

How do you know that  $y = \log_3 x$  is a function?

b) Set up tables of values for both the exponential function,  $f(x)$ , and its inverse,  $f^{-1}(x)$ . Plot the points and join them with a smooth curve.

$(x,y) \rightarrow (y,x)$

$f(x) = 3^x$		$f^{-1}(x) = \log_3 x$	
$x$	$y$	$x$	$y$
-3	$\frac{1}{27}$	$\frac{1}{27}$	-3
-2	$\frac{1}{9}$	$\frac{1}{9}$	-2
-1	$\frac{1}{3}$	$\frac{1}{3}$	-1
0	1	1	0
1	3	3	1
2	9	9	2
3	27	27	3



The graph of the inverse,  $f^{-1}(x) = \log_3 x$ , is a reflection of the graph

of  $f(x) = 3^x$  about the line  $y = x$ . For  $f^{-1}(x) = \log_3 x$ ,

- the domain is  $\{x \mid x > 0, x \in \mathbb{R}\}$  and the range is  $\{y \mid y \in \mathbb{R}\}$
- the x-intercept is 1
- there is no y-intercept
- the vertical asymptote, the y-axis, has equation  $x = 0$ ; there is no horizontal asymptote

How do the characteristics of  $f^{-1}(x) = \log_3 x$  compare to the characteristics of  $f(x) = 3^x$ ?

### Key Ideas


- A logarithm is an exponent.
- Equations in exponential form can be written in logarithmic form and vice versa.

**Exponential Form**      **Logarithmic Form**

$$x = c^y \qquad y = \log_c x$$

- The inverse of the exponential function  $y = c^x$ ,  $c > 0$ ,  $c \neq 1$ , is  $x = c^y$  or, in logarithmic form,  $y = \log_c x$ . Conversely, the inverse of the logarithmic function  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$ , is  $x = \log_c y$  or, in exponential form,  $y = c^x$ .
- The graphs of an exponential function and its inverse logarithmic function are reflections of each other in the line  $y = x$ , as shown.
- For the logarithmic function  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$ ,
  - the domain is  $\{x \mid x > 0, x \in \mathbb{R}\}$
  - the range is  $\{y \mid y \in \mathbb{R}\}$
  - the x-intercept is 1
  - the vertical asymptote is  $x = 0$ , or the y-axis
- A common logarithm has base 10. It is not necessary to write the base for common logarithms:


$$\log_{10} x = \log x$$




## Homework

#1-5, 8, 10, 12, 13, 17 on page 380

8.1 Understanding Logarithms, pages 380 to 382




ii)  $y = \log_2 x$   
 iii) domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ ,  
 range  $\{y \mid y \in \mathbb{R}\}$ ,  
 x-intercept 1, no y-intercept,  
 vertical asymptote  $x = 0$



ii)  $y = \log_3 x$   
 iii) domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ ,  
 range  $\{y \mid y \in \mathbb{R}\}$ ,  
 x-intercept 1, no y-intercept,  
 vertical asymptote  $x = 0$

2. a)  $\log_{12} 144 = 2$       b)  $\log_8 2 = \frac{1}{3}$   
 c)  $\log_{10} 0.000\ 01 = -5$       d)  $\log_7 (y + 3) = 2x$   
 3. a)  $5^2 = 25$       b)  $8^{\frac{2}{3}} = 4$   
 c)  $10^6 = 1\ 000\ 000$       d)  $11^y = x + 3$   
 4. a) 3      b) 0      c)  $\frac{1}{3}$       d) -3  
 5.  $a = 4; b = 5$



domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ ,  
 range  $\{y \mid y \in \mathbb{R}\}$ ,  
 x-intercept 1,  
 no y-intercept,  
 vertical asymptote  $x = 0$

10. They are reflections of each other in the line  $y = x$ .  
 11. a) They have the exact same shape.  
 b) One of them is increasing and the other is decreasing.  
 12. a) 216      b) 81      c) 64      d) 8  
 13. a) 7      b) 6  
 14. a) 0      b) 1  
 15. -1  
 16. 16  
 17. a)  $t = \log_{0.11} N$       b) 145 days  
 18. The larger asteroid had a relative risk that was 1479 times as dangerous.  
 19. 1000 times as great  
 20. 5  
 21.  $m = 14, n = 13$   
 22.  $4n$   
 23.  $y = 3^{2^x}$

### Product Law of Logarithms

The logarithm of a product of numbers can be expressed as the sum of the logarithms of the numbers.

$$\log_c MN = \log_c M + \log_c N$$

*Proof*

Let  $\log_c M = x$  and  $\log_c N = y$ , where  $M$ ,  $N$ , and  $c$  are positive real numbers with  $c \neq 1$ .

Write the equations in exponential form as  $M = c^x$  and  $N = c^y$ :

$$MN = (c^x)(c^y)$$

$$MN = c^{x+y}$$

$$\log_c MN = x + y$$

$$\log_c MN = \log_c M + \log_c N$$

Apply the product law of powers.

Write in logarithmic form.

Substitute for  $x$  and  $y$ .

---

### Quotient Law of Logarithms

The logarithm of a quotient of numbers can be expressed as the difference of the logarithms of the dividend and the divisor.

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

*Proof*

Let  $\log_c M = x$  and  $\log_c N = y$ , where  $M$ ,  $N$ , and  $c$  are positive real numbers with  $c \neq 1$ .

Write the equations in exponential form as  $M = c^x$  and  $N = c^y$ :

$$\frac{M}{N} = \frac{c^x}{c^y}$$

$$\frac{M}{N} = c^{x-y}$$

Apply the quotient law of powers.

$$\log_c \frac{M}{N} = x - y$$

Write in logarithmic form.

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

Substitute for  $x$  and  $y$ .

## Power Law of Logarithms

The logarithm of a power of a number can be expressed as the exponent times the logarithm of the number.

$$\log_c M^P = P \log_c M$$

How could you prove the quotient law using the product law and the power law?

*Proof*

Let  $\log_c M = x$ , where  $M$  and  $c$  are positive real numbers with  $c \neq 1$ .

Write the equation in exponential form as  $M = c^x$ .

Let  $P$  be a real number.

$$M = c^x$$

$$M^P = (c^x)^P$$

$$M^P = c^{xP}$$

Simplify the exponents.

$$\log_c M^P = xP$$

Write in logarithmic form.

$$\log_c M^P = (\log_c M)P$$

Substitute for  $x$ .

$$\log_c M^P = P \log_c M$$

The laws of logarithms can be applied to logarithmic functions, expressions, and equations.

## Example 1

### Use the Laws of Logarithms to Expand Expressions

Write each expression in terms of individual logarithms of  $x$ ,  $y$ , and  $z$ .

a)  $\log_5 \frac{xy}{z}$

b)  $\log_7 \sqrt[3]{x}$

c)  $\log_6 \frac{1}{x^2}$

d)  $\log \frac{x^3}{y\sqrt{z}}$



## Example 2

### Use the Laws of Logarithms to Evaluate Expressions

Use the laws of logarithms to simplify and evaluate each expression.

- a)  $\log_6 8 + \log_6 9 - \log_6 2$
- b)  $\log_7 7\sqrt{7}$
- c)  $2 \log_2 12 - \left( \log_2 6 + \frac{1}{3} \log_2 27 \right)$

### Example 3

#### Use the Laws of Logarithms to Simplify Expressions

Write each expression as a single logarithm in simplest form. State the restrictions on the variable.

a)  $\log_7 x^2 + \log_7 x - \frac{5 \log_7 x}{2}$

b)  $\log_5 (2x - 2) - \log_5 (x^2 + 2x - 3)$

**Key Ideas**

- Let  $P$  be any real number, and  $M$ ,  $N$ , and  $c$  be positive real numbers with  $c \neq 1$ . Then, the following laws of logarithms are valid.

Name	Law	Description
Product	$\log_c MN = \log_c M + \log_c N$	The logarithm of a product of numbers is the sum of the logarithms of the numbers.
Quotient	$\log_c \frac{M}{N} = \log_c M - \log_c N$	The logarithm of a quotient of numbers is the difference of the logarithms of the dividend and divisor.
Power	$\log_c M^P = P \log_c M$	The logarithm of a power of a number is the exponent times the logarithm of the number.

- Many quantities in science are measured using a logarithmic scale. Two commonly used logarithmic scales are the decibel scale and the pH scale.

## Do I really understand??...

a) Express the following as a single logarithm...  $2\log_2 3^2 + \log_2 6 - 3\log_2 3$

b) Evaluate the following...  $\log_2 (32)^{\frac{1}{3}}$

c) Express the following as a single logarithm...  $\frac{1}{2}[(\log_5 a + 2\log_5 b) - 3\log_5 c]$

d) Express as a single logarithm in simplest form...

$$\frac{3}{4} \left[ 12(\log_8 x^2 - 2\log_8 x) + 8\log_8 \sqrt{x} - 4\log_8 \frac{1}{x^7} \right]$$