

Introduction to Trigonometric Equations

trigonometric equation

- an equation involving trigonometric ratios

Focus on...

- algebraically solving first-degree and second-degree trigonometric equations in radians and in degrees
- verifying that a specific value is a solution to a trigonometric equation
- identifying exact and approximate solutions of a trigonometric equation in a restricted domain
- determining the general solution of a trigonometric equation

Did You Know?

In equations, mathematicians often use the notation $\cos^2 \theta$. This means the same as $(\cos \theta)^2$.

Let's start with basic LINEAR trigonometric equations...

...Pre-Calculus 110

Solve: $\sin \theta = 0.9659$, $-360^\circ < \theta < 720^\circ$

- Reference angle?
- Which quadrants?
- Any co-terminal angles acceptable?

- If the domain is in degrees, give solutions in degrees.
- If the domain is in radians, give solutions in radians.

$$\sin \theta = 0.9659, -360^\circ < \theta < 720^\circ$$

Where is $\sin \theta$ positive

$$\bar{\theta} = \sin^{-1}(0.9659)$$

$$\bar{\theta} = 75^\circ$$

 reference angle

Q1	Q2
$\theta = \bar{\theta}$	$\theta = 180^\circ - \bar{\theta}$
$\theta = 75^\circ$	$\theta = 180^\circ - 75^\circ$
	$\theta = 105^\circ$

To find angles between -360° and 720°

$$75^\circ - 360^\circ = -285^\circ$$

$$105^\circ - 360^\circ = -255^\circ$$

$$75^\circ + 360^\circ = 435^\circ$$

$$105^\circ + 360^\circ = 465^\circ$$

Solutions:

$$\theta = -285^\circ, -255^\circ, 75^\circ, 105^\circ, 435^\circ, 465^\circ$$

Solve: $\sec \theta = -1.3054$, $-2\pi \leq \theta \leq 2\pi$ (radians)

$$\cos \theta = \frac{1}{-1.3054}$$

$$\cos \theta = -0.7660$$

$$\bar{\theta} = \cos^{-1}(0.7660)$$

$$\bar{\theta} = 0.698$$

Where is $\cos \theta$ negative

Q2	Q3
$\theta = \pi - \bar{\theta}$	$\theta = \pi + \bar{\theta}$
$\theta = 3.14 - 0.698$	$\theta = 3.14 + 0.698$
$\theta = 2.443$	$\theta = 3.838$

Find angles between $-\pi$ and π

$$2.443 - 6.28 = -3.838$$

$$3.838 - 6.28 = -2.443$$

Solutions:

$$\theta = -3.838, -2.443, 2.443, 3.838$$

Exact Values → No Calculators

Ex. $\sqrt{2} \cos \theta + 1 = 0, -360^\circ \leq \theta \leq 720^\circ$

$$\sqrt{2} \cos \theta = -1$$

$$\cos \theta = \frac{-1}{\sqrt{2}}$$

(Use Special Triangles) $\bar{\theta} = 45^\circ$

Where is $\cos \theta$ negative

Q2	Q3
$\theta = 180^\circ - \bar{\theta}$	$\theta = 180^\circ + \bar{\theta}$
$\theta = 180^\circ - 45^\circ$	$\theta = 180^\circ + 45^\circ$
$\theta = 135^\circ$	$\theta = 225^\circ$

Find angles between -360° and 720°

$$135^\circ - 360^\circ = -225^\circ$$

$$225^\circ - 360^\circ = -135^\circ$$

$$135^\circ + 360^\circ = 495^\circ$$

$$225^\circ + 360^\circ = 585^\circ$$

Solutions.

$$\theta = -225^\circ, -135^\circ, 135^\circ, 225^\circ, 495^\circ, 585^\circ$$

Exact Value → No Calculator

Ex. $\sin x + 1 = 0, -2\pi \leq x \leq 4\pi$

$$\begin{array}{l} \sin x = -1 \\ (\text{Unit Circle}) \quad x = \frac{3\pi}{2} \end{array} \quad \begin{array}{l} \frac{3\pi}{2} - 2\pi \\ \frac{3\pi}{2} - \frac{4\pi}{2} \\ -\frac{\pi}{2} \end{array} \quad \left| \begin{array}{l} \frac{3\pi}{2} + 2\pi \\ \frac{3\pi}{2} + \frac{4\pi}{2} \\ \frac{7\pi}{2} \end{array} \right.$$

Solutions:

$$x = -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}$$

Your Turn

Solve each trigonometric equation in the specified domain.

- a) $3 \cos \theta - 1 = \cos \theta + 1, -2\pi \leq \theta \leq 2\pi$
- b) $4 \sec x + 8 = 0, 0^\circ \leq x < 360^\circ$

Check Up:

$$\cot\theta = 0.7834 \quad 0 < \theta < 2\pi$$

$$\tan\theta = 1.2765$$

$$\bar{\theta} = \tan^{-1}(1.2765)$$

$$\bar{\theta} = 0.9063$$

radians

Where is $\tan\theta$ positive

Q1	Q3
$\theta = \bar{\theta}$	$\theta = \pi + \bar{\theta}$
$\theta = 0.9063$	$\theta = 3.14 + 0.9063$
	$\theta = 4.0463$

Solutions:

$$\theta = 0.9063, 4.0463$$

$$2\cos\theta - 5 = -6$$

$$-360^\circ < \theta < 360^\circ$$

$$2\cos\theta = -1$$

$$\cos\theta = -\frac{1}{2}$$

$$\bar{\theta} = 60^\circ$$

Degrees

Where is $\cos\theta$ negative?

Q2	Q3
$\theta = 180^\circ - \bar{\theta}$	$\theta = 180^\circ + \bar{\theta}$
$\theta = 180^\circ - 60^\circ$	$\theta = 180^\circ + 60^\circ$
$\theta = 120^\circ$	$\theta = 240^\circ$

Find angles between -360° and 360°

$$120^\circ - 360^\circ = -240^\circ$$

$$240^\circ - 360^\circ = -120^\circ$$

Solutions:

$$\theta = -240^\circ, -120^\circ, 120^\circ, 240^\circ$$

$$\textcircled{3} \text{ c) } 5 - \tan^2 \theta = 4, \quad -180^\circ \leq \theta \leq 360^\circ$$

$$-\tan^2 \theta = -1$$

where is $\tan \theta$ (+/-)

$$\tan^2 \theta = 1$$

$$\tan \theta = \pm 1$$

$$\bar{\theta} = 45^\circ$$

Q1	Q2	Q3	Q4
$\theta = 45^\circ$	$\theta = 135^\circ$	$\theta = 225^\circ$	$\theta = 315^\circ$
$\theta = -315^\circ$	$\theta = -225^\circ$	$\theta = -135^\circ$	$\theta = -45^\circ$

Solutions:

$$\theta = -135^\circ, -45^\circ, 45^\circ, 135^\circ, 225^\circ, 315^\circ$$

Questions from Homework

⑪ $\cos \theta = -0.15$

$\bar{\theta} = \cos^{-1}(0.15)$ when finding $\bar{\theta}$ always use positive value

$$\bar{\theta} = 81.4^\circ$$

$\cos^{-1}(-0.15)$
81.37307344

Where is $\cos \theta$ negative

Q2 Q3

$$\theta = 180^\circ - 81.4^\circ \quad \theta = 180^\circ + 81.4^\circ$$

$$\theta = 98.6^\circ \quad \theta = 261.4^\circ$$

$$98.6^\circ + 360^\circ n, n \in \mathbb{I}$$

$$261.4^\circ + 360^\circ n, n \in \mathbb{I}$$

Let's move onto QUADRATIC trigonometric equations...

...Pre-Calculus 110

\nwarrow (2nd Degree) (Factoring!!)

- What strategies can we use to solve quadratic equations?
- Quadratic trigonometric equations will ultimately become TWO linear trigonometric equations.

Solve: $2x^2 + x = 1$

$$\begin{aligned} 2x^2 + x - 1 &= 0 \\ (2x^2 + 2x)(-1x - 1) &= 0 \\ 2x(x+1) - 1(x+1) &= 0 \\ (2x-1)(x+1) &= 0 \\ 2x-1=0 &\quad x+1=0 \\ 2x=1 &\quad x=-1 \\ x=\frac{1}{2} & \end{aligned}$$

$$\boxed{\begin{aligned} 2x-1 &= -2 \\ 2+1 &= 1 \end{aligned}}$$

Solve: $2\sin^2 x + \sin x = 1, 0 \leq x \leq 4\pi$

$$\begin{aligned} 2\sin^2 x + \sin x - 1 &= 0 \\ (2\sin^2 x + 2\sin x)(-\sin x - 1) &= 0 \\ 2\sin x(\sin x + 1) - 1(\sin x + 1) &= 0 \\ (2\sin x - 1)(\sin x + 1) &= 0 \\ 2\sin x - 1 &= 0 \quad \sin x + 1 = 0 \\ \sin x = \frac{1}{2} & \quad \sin x = -1 \\ (\text{Triangles}) \bar{x} &= \frac{\pi}{6} \\ x &= \frac{\pi}{6} \\ x &= \frac{5\pi}{6} \\ x &= \frac{13\pi}{6} \\ x &= \frac{17\pi}{6} \\ x &= \frac{3\pi}{2} \\ x &= \frac{7\pi}{2} \end{aligned}$$

Solutions

$$x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, \frac{17\pi}{6}, \frac{3\pi}{2}, \frac{7\pi}{2}$$

$$\text{Ex. } \cos^2 \theta - \frac{1}{2} \cos \theta = 0, -2\pi \leq \theta \leq 4\pi$$

$$\cos \theta (\cos \theta - \frac{1}{2}) = 0 \quad (\text{Common Factor})$$

$$\cos \theta = 0 \quad (\text{Unit Circle})$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$\theta = \frac{5\pi}{3}, \frac{7\pi}{3}$$

$$\theta = -\frac{5\pi}{3}, -\frac{7\pi}{3}$$

$$\cos \theta - \frac{1}{2} = 0$$

$$\cos \theta = \frac{1}{2} \quad (\text{Triangles})$$

$$\theta = \frac{\pi}{3}$$

Q1

$$\theta = \frac{\pi}{3}$$

$$\theta = \frac{7\pi}{3}$$

$$\theta = -\frac{5\pi}{3}$$

Q4

$$\theta = 2\pi - \frac{\pi}{3} = \frac{6\pi}{3} - \frac{\pi}{3} = \frac{5\pi}{3}$$

$$\theta = \frac{11\pi}{3}$$

$$\theta = -\frac{\pi}{3}$$

Ex. $6\sin^2 x - \sin x = 2, -2\pi \leq \theta \leq 4\pi$ (Decomposition)

General Solution of a Trigonometric Equation

Solve: $3\cos^2 \theta - \cos \theta = 2; \theta \in \mathbb{R}$ *(Decomposition)*

Determine the general solution for $\sin^2 x - 1 = 0$ over the real numbers if x is measured in radians.

$$\begin{aligned} \sin^2 x - 1 &= 0 && \text{(Difference of Squares)} \\ (\sin x + 1)(\sin x - 1) &= 0 \\ \sin x + 1 &= 0 && \sin x - 1 = 0 \\ \sin x &= -1 && \sin x = 1 \\ \text{(unit circle)} & && \text{(unit circle)} \\ \theta = \frac{3\pi}{2} & && \theta = \frac{\pi}{2} \end{aligned}$$

Did You Know?

$2n$, where $n \in \mathbb{I}$, represents all even integers.

$2n + 1$, where $n \in \mathbb{I}$, is an expression for all odd integers.

$$x = \frac{\pi}{2} + 2\pi n, \text{ where } n \in \mathbb{I}$$

$$x = \frac{3\pi}{2} + 2\pi n, \text{ where } n \in \mathbb{I}$$

or

$$x = \frac{\pi}{2} + \pi n, \text{ where } n \in \mathbb{I}$$

or

$$(2n + 1)\left(\frac{\pi}{2}\right), n \in \mathbb{I}$$

Determine the general solution for $\cos^2 x - 1 = 0$, where the domain is real numbers measured in degrees.

Check-Up problem...

(Common Factor)

Solve:

$$\sin x \sec x + 2 \sin x = 0, x \in R \quad (x \text{ is measured in radians})$$

$$\sin x (\sec x + 2) = 0$$

$$\sin x = 0$$

(Unit Circle)

$$x = 0, \pi, 2\pi$$

$$0 + \pi n, n \in \mathbb{Z}$$

$$\sec x + 2 = 0$$

$$\sec x = -2$$

$$\cos x = -\frac{1}{2}$$

(Triangle)

$$x = \frac{\pi}{3}$$

Where is $\cos x$ negative?

Q2

Q3

$$x = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$$

$$x = \pi + \frac{\pi}{3} = \frac{4\pi}{3}$$

$$\frac{2\pi}{3} + 2\pi n, n \in \mathbb{Z}$$

$$\frac{4\pi}{3} + 2\pi n, n \in \mathbb{Z}$$

Unit Review...




What topics have we covered??

- Radian Measure
- Co-terminal angles
- Principal Angles
- Angular Velocity (Open Response)
- The Unit Circle
- Trig Expressions (Open Response)
- Trig Equations (Open Response)

Review...

Without a calculator determine the value of...

$$\frac{2 \cos 180^\circ + \sin 135^\circ}{\cos^2(-330^\circ)}$$



$$\frac{2(-1) + \left(\frac{1}{\sqrt{2}}\right)}{\left(\frac{\sqrt{3}}{2}\right)^2}$$

$$4. -\frac{2}{1} + \frac{\sqrt{2}}{2} \cdot 4$$

$$\frac{3}{4} \cdot 4$$

$\frac{-8 + 2\sqrt{2}}{3}$

Solve: $6 \sin^2 \theta - 3 \sin \theta = 0$, $0^\circ \leq \theta \leq 360^\circ$


- [A] $0^\circ, 30^\circ, 180^\circ, 330^\circ, 360^\circ$
 [C] $30^\circ, 90^\circ, 120^\circ, 270^\circ$

- [B] $0^\circ, 30^\circ, 180^\circ, 150^\circ, 360^\circ$
 [D] $0^\circ, 180^\circ, 210^\circ, 330^\circ, 360^\circ$

$\sin \theta$ is negative + $\tan \theta$ is positive (Quad 3)

If $\csc \theta < 0$ and $\tan \theta > 0$, then which of the following could be a possible measure of angle θ ?

- [A] $\frac{11\pi}{6}$ (Q4) [B] $\frac{4\pi}{3}$ (Q3) [C] $\frac{3\pi}{4}$ (Q2) [D] $\frac{\pi}{2}$



$$-\frac{25\pi}{4} + \frac{8\pi}{1} = -\frac{25\pi}{4} + \frac{32\pi}{4} = \frac{7\pi}{4}$$


What is the principal angle of $-\frac{25\pi}{4}$?

- [A] $\frac{3\pi}{4}$ [B] $\frac{\pi}{4}$

- [C] $-\frac{\pi}{4}$

- [D] $\frac{7\pi}{4}$

If the belt in the pulley system below travels 30 cm, what is the angle of rotation of the smaller pulley?



Small

$$r = 1.5 \text{ cm}$$

$$\alpha = 30 \text{ cm}$$

$$\theta = ?$$

[A] $\frac{\pi}{9}$ radians

[B] 20°

[C] 20 radians

[D] 5°

$$\Theta = \frac{\alpha}{r} = \frac{30 \text{ cm}}{1.5 \text{ cm}} = 20 \text{ rads}$$

Nibbles the hamster is running at 0.02 m/s on an exercise wheel of radius 8 cm. What is the angular velocity of this wheel?

[A] 0.15 rad/minute

[B] 240 rad/minute

[C] 0.25 rad/minute

[D] 15 radians/minute

Given:

$$r = 8 \text{ cm}$$

$$\alpha = 2 \text{ cm/s} \times 1 \text{ s} = 2 \text{ cm}$$

① Find Θ

$$\Theta = \frac{\alpha}{r} = \frac{2}{8} = 0.25 \text{ rads}$$

② Find V_a

$$V_a = \frac{\Theta}{t} = \frac{0.25 \text{ rads}}{\cancel{sec}} \times \frac{\cancel{60 sec}}{1 \text{ min}} = \boxed{\frac{15 \text{ rads}}{\text{min}}}$$

Solve: $2(1 - \sin \theta)^2 + \sin \theta = 2(3 - 4 \sin^2 \theta)$, $-360^\circ \leq \theta \leq 720^\circ$

Chapter 4 Review:

Pages 215 - 217

| Little Johnny has a rock tied to the end of a piece of rope 1.5 m long and he is swinging it around his head in a circular pattern. Mrs. Centripetal, his physics teacher, is watching Johnny out the window of her physics lab and notes that the rock is making 12 revolutions every 48 seconds.

- (a) Determine the angular velocity with which little Johnny is twirling the rope above his head. [2]

- (b) The rock comes flying from the rope 3 minutes after Mrs. Centripetal started to time little Johnny. How far did the rock travel during the 3 minutes? [2]

Attachments

Worksheet - Sketching Angles in Radians.doc