

1) 31 200 J

7) 0.0456 J/g °C

2) -31 700 J

8) 424 g

3) 120°C

9) 2.60 J/g °C

4) 28°C

10) 6.21 J

5) 1100 J

11) 42.6 L

6) 14 900 J

## Homework - Worksheet

$$C = \frac{J}{g \cdot ^\circ C}$$

④  $m = 350g$

$$T_i = 45.0^\circ C$$

$$q = -25.0 kJ$$

$$T_f = ?$$

$$q = mC\Delta T$$

$$-25000 J = (350g)(4.19 \frac{J}{g \cdot ^\circ C})$$

$$(T_f - 45.0^\circ C)$$

$$-17.05^\circ C = T_f - 45.0^\circ C$$

$$T_f = -17.05^\circ C + 45.0^\circ C$$

$$T_f = 28^\circ C$$

⑦  $m = 550g$

$$T_i = 95.0^\circ C$$

$$T_f = 30.0^\circ C$$

$$q = -1.63 \times 10^3 J$$

$$C = ?$$

$$q = mC\Delta T$$

$$-1630 J = (550g)C(-65.0^\circ C)$$

$$C = \frac{-1630 J}{(550g)(-65.0^\circ C)}$$

$$C = 0.0456 \frac{J}{g \cdot ^\circ C}$$

## PHASE CHANGE AND ENTHALPY

Classifying types of systems:


1. Open system - a system where both matter and energy can flow into or out of the system.
2. Closed system - a system where energy is allowed to be transferred into and out but matter cannot be transferred.
3. Isolated system - a system where neither matter nor energy is allowed to enter or leave the system.

ENTHALPY (H) - The total internal (potential) energy and kinetic energy of a system under constant pressure.

⇒ Enthalpy is usually expressed in kJ.

ENTHALPY CHANGE ( $\Delta H$ ) - A change under constant pressure where the surroundings of a system absorb energy or release it to the system.

PHASE CHANGE - is a change in the state of matter without a change in the chemical composition of the system.




⇒ **always involve a change in energy but never involve a change in temperature.**


**Question:**

- (i) What is the temperature where water just starts boiling?
- (ii) What is the temperature when water is boiling violently?
- (iii) If energy is still going into the water and the temperature is not increasing, where is the energy going?



Consider melting ice to water and then boiling water to steam:





## MOLAR ENTHALPY

For any system:

- an exothermic change involves a decrease in enthalpy

⇒ gives off energy to the surroundings

⇒  $\Delta H$  is negative.

- an endothermic change involves an increase in enthalpy.

⇒ takes in energy from the surroundings

⇒  $\Delta H$  is positive.

The enthalpies for substances undergoing phase changes have been measured experimentally. (TABLE 17.3 p. 522)

- enthalpies are reported as molar enthalpies and are expressed as kJ/mol.

	$H_{fus}$	$H_{vap}$
water	$6.01 \frac{\text{kJ}}{\text{mol}}$	$40.7 \frac{\text{kJ}}{\text{mol}}$

## Endothermic Phase Changes


- the molar enthalpy of fusion ( $H_{\text{fus}}$ ) represents the quantity of heat that the substance absorbs per mole as it changes state from **solid to liquid**.

- the molar enthalpy of vaporization ( $H_{\text{vap}}$ ) represents the quantity of heat that the substance absorbs per mole as it changes state from **liquid to gas**.

## Exothermic Phase Changes

- the molar enthalpy of condensation ( $H_{\text{cond}}$ ) represents the quantity of heat that the substance releases per mole as it changes state from **gas to liquid**

- the molar enthalpy of solidification ( $H_{\text{solid}}$ ) represents the quantity of heat that the substance releases per mole as it changes state from **liquid to solid**.




$$\Delta H_{\text{fus}} = - \Delta H_{\text{solid}}$$

$$\Delta H_{\text{vap}} = - \Delta H_{\text{cond}}$$

## Example

If 500. g of  $\text{CCl}_2\text{F}_2(l)$  is vaporized at SATP, find the enthalpy change of the system ( $H_{\text{vap}} = 34.99 \text{ kJ/mol}$ ).

$$m = 500. \text{ g}$$




$$H_{\text{vap}} = 34.99 \frac{\text{kJ}}{\text{mol}}$$

$$\Delta H_{\text{vap}} = ?$$

$$\Delta H_{\text{vap}} = n H_{\text{vap}}$$

$$\Delta H_{\text{vap}} = \left( \frac{500. \text{ g}}{120.91 \text{ g/mol}} \right) \left( 34.99 \frac{\text{kJ}}{\text{mol}} \right)$$

$$\Delta H_{\text{vap}} = 145 \text{ kJ}$$





# Worksheet