

Understanding Logarithms

Focus on...

- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- determining the characteristics of the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

$(x,y) \rightarrow (y,x)$
reflection in
the line
 $y=x$

Questions from Homework

11. Explain how the graph of

$\frac{1}{3}(y + 2) = \log_6(x - 4)$ can be generated by transforming the graph of $y = \log_6 x$.

$y + 2 = 3 \log_6(x - 4)$ Divide $a+k$ by $\frac{1}{3}$
 $y = 3 \log_6(x - 4) - 2$ Subtract 2 from both sides
 $y = \underline{3} \log_6(\underline{x - 4}) - \underline{2}$
 $a = 3 \rightarrow$ A vertical stretch by a factor of 3
 $b = 1 \rightarrow$ No horizontal stretch.
 $h = 4 \rightarrow$ Translated 4 units right.
 $k = -2 \rightarrow$ " " 2 " down.

5. Identify the following characteristics of the graph of each function.

- i) the equation of the asymptote
- ii) the domain and range
- iii) the y-intercept, to one decimal place if necessary
- iv) the x-intercept, to one decimal place if necessary

a) $y = -5 \log_3(x + 3)$

b) $y = \log_6(4(x + 9))$

c) $y = \log_5(\underline{x + 3}) - 2$

d) $y = -3 \log_2(x + 1) - 6$

b) $a = 1$ $b = 4$ $h = -9$ $k = 0$
 $(x, y) \rightarrow \left[\frac{1}{4}x - 9, 1y + 0 \right]$

Base: $y = \log_6 x$
 D: $\{x \mid x > 0, x \in \mathbb{R}\}$
 R: $\{y \mid y \in \mathbb{R}\}$
 x int: (1, 0)
 y int: none
 VA: $x = 0$

For: $y = \log_6(4(x + 9))$
 (i) VA: $x = -9$
 (ii) D: $\{x \mid x > -9, x \in \mathbb{R}\}$
 R: $\{y \mid y \in \mathbb{R}\}$
 (iii) y intercept (Let $x = 0$)
 $y = \log_6(4(x + 9))$
 $y = \log_6(4(0 + 9))$
 $y = \log_6 36$
 $y = 2$ or (0, 2)
 (iv) x-intercept (Let $y = 0$)
 $y = \log_6(4(x + 9))$
 $0 = \log_6(4x + 36) \leftarrow \text{ans.}$
 $6^0 = 4x + 36$

General Properties of Logarithms:

* a logarithm is an exponent!

If $C > 0$ and $C \neq 1$, then...

$$(i) \log_C 1 = 0$$

$$(ii) \log_C C^x = x$$

$$(iii) C^{\log_C x} = x$$

Did You Know?

The input value for a logarithm is called an argument. For example, in the expression $\log_6 1$, the argument is 1.

$$(i) \log_5 1 = 0$$

$$(ii) \log_2 2^3 = 3$$

$$(iii) 7^{\log_7 49} = 49$$

$$5^{\log_5 10} = 10$$

Product Law of Logarithms (Page 394)

The logarithm of a product of numbers can be expressed as the sum of the logarithms of the numbers.

$$\log_c MN = \log_c M + \log_c N$$

Ex: $\log_2 8 + \log_2 3 = \log_2 (8 \times 3) = \log_2 24 \approx 4.59$

Proof

Let $\log_c M = x$ and $\log_c N = y$, where M , N , and c are positive real numbers with $c \neq 1$.

Write the equations in exponential form as $M = c^x$ and $N = c^y$:

$$MN = (c^x)(c^y)$$

$$MN = c^{x+y}$$

$$\log_c MN = x + y$$

$$\log_c MN = \log_c M + \log_c N$$

Apply the product law of powers.

Write in logarithmic form.

Substitute for x and y .

Quotient Law of Logarithms

The logarithm of a quotient of numbers can be expressed as the difference of the logarithms of the dividend and the divisor.

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

Ex: $\log 400 - \log 4 = \log\left(\frac{400}{4}\right) = \log 100 = \boxed{2}$

Proof

Let $\log_c M = x$ and $\log_c N = y$, where M , N , and c are positive real numbers with $c \neq 1$.

Write the equations in exponential form as $M = c^x$ and $N = c^y$:

$$\frac{M}{N} = \frac{c^x}{c^y}$$

$$\frac{M}{N} = c^{x-y}$$

$$\log_c \frac{M}{N} = x - y$$

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

Apply the quotient law of powers.

Write in logarithmic form.

Substitute for x and y .

Power Law of Logarithms

The logarithm of a power of a number can be expressed as the exponent times the logarithm of the number.

$$\log_c M^P = P \log_c M$$

How could you prove the quotient law using the product law and the power law?

Proof

Let $\log_c M = x$, where M and c are positive real numbers with $c \neq 1$.

Write the equation in exponential form as $M = c^x$.

Let P be a real number.

$$M = c^x$$

$$M^P = (c^x)^P$$

$$M^P = c^{xP}$$

$$\log_c M^P = xP$$

$$\log_c M^P = (\log_c M)P$$

$$\log_c M^P = P \log_c M$$

Simplify the exponents.

Write in logarithmic form.

Substitute for x .

The laws of logarithms can be applied to logarithmic functions, expressions, and equations.

Ex: $\log_2 \sqrt{8}$

$\log_2 8^{1/2}$

$\frac{1}{2} \log_2 8$

$\frac{1}{2} (3)$

$\frac{3}{2}$

Homework

Finish Exercise 2

Example 1

Use the Laws of Logarithms to Expand Expressions

Write each expression in terms of individual logarithms of x , y , and z .

a) $\log_5 \frac{xy}{z}$

b) $\log_7 \sqrt[3]{x}$

c) $\log_6 \frac{1}{x^2}$

d) $\log \frac{x^3}{y\sqrt{z}}$

Example 2

Use the Laws of Logarithms to Evaluate Expressions

Use the laws of logarithms to simplify and evaluate each expression.

- a) $\log_6 8 + \log_6 9 - \log_6 2$
- b) $\log_7 7\sqrt{7}$
- c) $2 \log_2 12 - \left(\log_2 6 + \frac{1}{3} \log_2 27 \right)$

Example 3

Use the Laws of Logarithms to Simplify Expressions

Write each expression as a single logarithm in simplest form. State the restrictions on the variable.

a) $\log_7 x^2 + \log_7 x - \frac{5 \log_7 x}{2}$

b) $\log_5 (2x - 2) - \log_5 (x^2 + 2x - 3)$

Key Ideas

- Let P be any real number, and M , N , and c be positive real numbers with $c \neq 1$. Then, the following laws of logarithms are valid.

Name	Law	Description
Product	$\log_c MN = \log_c M + \log_c N$	The logarithm of a product of numbers is the sum of the logarithms of the numbers.
Quotient	$\log_c \frac{M}{N} = \log_c M - \log_c N$	The logarithm of a quotient of numbers is the difference of the logarithms of the dividend and divisor.
Power	$\log_c M^P = P \log_c M$	The logarithm of a power of a number is the exponent times the logarithm of the number.

- Many quantities in science are measured using a logarithmic scale. Two commonly used logarithmic scales are the decibel scale and the pH scale.

Homework

Do I really understand??...

a) Express the following as a single logarithm... $2\log_2 3^2 + \log_2 6 - 3\log_2 3$

b) Evaluate the following... $\log_2 (32)^{\frac{1}{3}}$

c) Express the following as a single logarithm... $\frac{1}{2}[(\log_5 a + 2\log_5 b) - 3\log_5 c]$

d) Express as a single logarithm in simplest form...

$$\frac{3}{4} \left[12(\log_8 x^2 - 2\log_8 x) + 8\log_8 \sqrt{x} - 4\log_8 \frac{1}{x^7} \right]$$