

## Questions from Homework

② c)  $\log_{10} 2 + \log_{10} 5$       d)  $\log_{10} \sqrt{\frac{1}{10}}$

$\log_{10} (2 \times 5)$                        $\log_{10} \left(\frac{1}{10}\right)^{\frac{1}{2}}$

$\log_{10} 10$                                $\log_{10} (10^{-1})^{\frac{1}{2}}$

1     $\log_{10} (10^{-\frac{1}{2}})$

$-\frac{1}{2}$

③ d)  $4 \log_2 x - \frac{1}{3} \log_2 (x^2 + 1) + \log_2 (x - 1)$

$\log_2 x^4 - \log_2 (x^2 + 1)^{\frac{1}{3}} + \log_2 (x - 1)$

$\log_2 \left[ \frac{x^4(x-1)}{(x^2+1)^{\frac{1}{3}}} \right]$

$\log_2 \left( \frac{x^4(x-1)}{\sqrt[3]{x^2+1}} \right)$

Do I really understand??...

- a) Express the following as a single logarithm...  $2\log_2 3^2 + \log_2 6 - 3\log_2 3$
- b) Evaluate the following...  $\log_2(32)^{\frac{1}{3}}$
- c) Express the following as a single logarithm...  $\frac{1}{2}[(\log_5 a + 2\log_5 b) - 3\log_5 c]$
- d) Express as a single logarithm in simplest form...

$$\frac{3}{4} \left[ 12(\log_b x^2 - 2\log_b x) + 8\log_b \sqrt{x} - 4\log_b \frac{1}{x^7} \right]$$

$$\begin{aligned}
 &2\log_2 3^2 + \log_2 6 - 3\log_2 3 \\
 &2\log_2 9 + \log_2 6 - 3\log_2 3 \\
 &\log_2 9 + \log_2 6 - \log_2 3 \\
 &\log_2 81 + \log_2 6 - \log_2 27 \\
 &\log_2 \left( \frac{8 \cdot 6}{27} \right) \\
 &\log_2 18
 \end{aligned}$$

$$\begin{aligned}
 &\text{Let } x = \log_2(32)^{\frac{1}{3}} \\
 &x = \log_2(32)^{\frac{1}{3}} \\
 &2^x = (32)^{\frac{1}{3}} \\
 &2^x = (2^5)^{\frac{1}{3}} \\
 &2^x = 2^{\frac{5}{3}} \\
 &x = \frac{5}{3}
 \end{aligned}$$

$$\begin{aligned}
 &\frac{1}{2}[(\log_5 a + 2\log_5 b) - 3\log_5 c] \\
 &\frac{1}{2}[\log_5 a + \log_5 b^2 - \log_5 c^3] \\
 &\frac{1}{2}[\log_5 \left( \frac{ab^2}{c^3} \right)] \\
 &\frac{1}{2}\log_5 \left( \frac{ab^2}{c^3} \right) \\
 &\log_5 \left( \frac{ab^2}{c^3} \right)^{\frac{1}{2}} \\
 &\log_5 \sqrt{\frac{ab^2}{c^3}} \quad \text{or} \quad \log_5 b \sqrt{\frac{a}{c^3}}
 \end{aligned}$$

$$\begin{aligned}
 &\frac{3}{4} \left[ 12(\log_b x^2 - 2\log_b x) + 8\log_b \sqrt{x} - 4\log_b \frac{1}{x^7} \right] \\
 &\frac{3}{4} [12\log_b x^2 - 24\log_b x + 8\log_b x^{\frac{1}{2}} - 4\log_b x^{-7}] \\
 &9\log_b(x^2) - 8\log_b x + 6\log_b(x^{\frac{1}{2}}) - 3\log_b(x^{-7}) \\
 &\log_b(x^2)^9 - \log_b x^8 + \log_b(x^{\frac{1}{2}})^6 - \log_b(x^{-7})^3 \\
 &\log_b x^{18} - \log_b x^8 + \log_b x^3 - \log_b x^{-21} \\
 &\log_b \left( \frac{x^{18}}{x^{-21}} \right) \\
 &\log_b x^{39}
 \end{aligned}$$

# Logarithmic and Exponential Equations

## Focus on...

---

- solving a logarithmic equation and verifying the solution
- explaining why a value obtained in solving a logarithmic equation may be extraneous
- solving an exponential equation in which the bases are not powers of one another
- solving a problem that involves exponential growth or decay
- solving a problem that involves the application of exponential equations to loans, mortgages, and investments
- solving a problem by modelling a situation with an exponential or logarithmic equation

**General Properties of Logarithms:**

If  $C > 0$  and  $C \neq 1$ , then...

(i)  $\log_C 1 = 0$

(ii)  $\log_C C^x = x$

(iii)  $C^{\log_C x} = x$

**Did You Know?**

The input value for a logarithm is called an argument. For example, in the expression  $\log_6 1$ , the argument is 1.

Example 1

Solve Logarithmic Equations

Solve.

a)  $\log_6 (2x - 1) = \log_6 11$

b)  $\log (8x + 4) = 1 + \log (x + 1)$

c)  $\log_2 (x + 3)^2 = 4$

a)  $\log_6 (2x-1) = \log_6 11$

$2x-1 = 11$

$2x = 12$

$x = 6$

Check  $x=6$

$\log_6 (2(6)-1) = \log_6 11$

$\log_6 11$

$x=6$  is a solution

b)  $\log (8x+4) = 1 + \log (x+1)$

$\log (8x+4) - \log (x+1) = 1$

$\log \left( \frac{8x+4}{x+1} \right) = 1$  logarithmic

$\frac{10^1}{1} = \frac{8x+4}{x+1}$  Exponential

$10(x+1) = 1(8x+4)$

$10x+10 = 8x+4$

$2x = -6$

$x = -3$

Check  $x=-3$

$\log (8(-3)+4) = 1 + \log (-3+1)$

$\log (-24+4)$

$\log (-20)$

undefined

$x=-3$  is an extraneous root (not a solution)

c)  $\log_2 (x+3)^2 = 4$  Logarithmic

$2^4 = (x+3)^2$  Exponential

$16 = (x+3)(x+3)$

$16 = x^2 + 3x + 3x + 9$

$16 = x^2 + 6x + 9$

$0 = x^2 + 6x - 7$  Simple trinomial

$0 = (x-1)(x+7)$

$x-1=0 \quad | \quad x+7=0$

$x=1 \quad | \quad x=-7$

Both  $x=1$  &  $x=-7$  are solutions

Check  $x=-7$

$\log_2 (-7+3)^2 = 4$

$\log_2 (-4)^2$

$\log_2 16$

4

Check  $x=1$

$\log_2 (1+3)^2 = 4$

$\log_2 16$

4

Example 2

Solve Exponential Equations Using Logarithms

Solve. Round your answers to two decimal places.

- a)  $4^x = 605$
- b)  $8(3^{2x}) = 568$
- c)  $4^{2x-1} = 3^{x+2}$

a)  $4^x = 605$  Take the common logarithm of both sides.

$$\log 4^x = \log 605$$

$$x \log 4 = \log 605$$

$$x = \frac{\log 605}{\log 4}$$

$$x \approx 4.62$$

$x \approx 4.62$  is a solution

Check  $x = 4.62$

$4^x$	605
$4^{4.62}$	
605	

b)  $\frac{8(3^{2x})}{8} = \frac{568}{8}$

$$3^{2x} = 71$$

Take the common logarithm of both sides.

$$\log 3^{2x} = \log 71$$

$$\frac{2x \log 3}{2 \log 3} = \frac{\log 71}{2 \log 3}$$

$$x \approx 1.94$$

$x \approx 1.94$  is a solution

Check  $x = 1.94$

$8(3^{2x})$	568
$8(3^{2(1.94)})$	
$8(71)$	
568	

c)  $4^{2x-1} = 3^{x+2}$  Take the common logarithm of both sides.

$$\log 4^{2x-1} = \log 3^{x+2}$$

$$(2x-1) \log 4 = (x+2) \log 3$$

$$2x \log 4 - \log 4 = x \log 3 + 2 \log 3$$

Group like terms

$$2x \log 4 - x \log 3 = 2 \log 3 + \log 4$$

Common factor of x

$$x(2 \log 4 - \log 3) = 2 \log 3 + \log 4$$

$$x = \frac{2 \log 3 + \log 4}{2 \log 4 - \log 3}$$

$x \approx 2.1407$  is a solution

Check  $x = 2.1407$

$4^{2(2.1407)-1}$	$3^{2.1407+2}$
-------------------	----------------

## Questions from Homework

4. Determine whether the possible roots listed are extraneous to the logarithmic equation given.

a)  $\log_7 x + \log_7 (x - 1) = \log_7 4x$   
possible roots:  $x = 0, x = 5$

b)  $\log_6 (x^2 - 24) - \log_6 x = \log_6 5$   
possible roots:  $x = 3, x = -8$

c)  $\log_3 (x + 3) + \log_3 (x + 5) = 1$   
possible roots:  $x = -2, x = -6$

d)  $\log_2 (x - 2) = 2 - \log_2 (x - 5)$   
possible roots:  $x = 1, x = 6$

possible solutions:

test  $x=0$

$$\log_7(0) + \log_7(0-1) \quad | \quad \log_7 4(0)$$

$$\log_7 0 + \log_7(-1)$$

undefined

$x=0$  is an extraneous root

test  $x=5$

$$\log_7(5) + \log_7(5-1) \quad | \quad \log_7 4(5)$$

$$\log_7 5 + \log_7 4$$

$$\log_7(5 \times 4)$$

$$\log_7 20$$

$x=5$  is a solution

## Example 4

### Solve a Problem Involving Exponential Growth and Decay

When an animal dies, the amount of radioactive carbon-14 (C-14) in its bones decreases. Archaeologists use this fact to determine the age of a fossil based on the amount of C-14 remaining.

- The half-life of C-14 is 5730 years.

Head-Smashed-In Buffalo Jump in southwestern Alberta is recognized as the best example of a buffalo jump in North America. The oldest bones unearthed at the site had 49.5% of the C-14 left. How old were the bones when they were found?

(Solve for  $t$ )



Buffalo skull display, Head-Smashed-In buffalo Jump Visitor Centre, near Fort McLeod, Alberta

$c = \frac{1}{2}$  or 0.5  
denominator of exponent



**Solution**

Carbon-14 decays by one half for each 5730-year interval. The mass,  $m$ , remaining at time  $t$  can be found using the relationship  $m(t) = m_0 \left(\frac{1}{2}\right)^{\frac{t}{5730}}$ , where  $m_0$  is the original mass.

Since 49.5% of the C-14 remains after  $t$  years, substitute  $0.495m_0$  for  $m(t)$  in the formula  $m(t) = m_0 \left(\frac{1}{2}\right)^{\frac{t}{5730}}$ .

$$0.495m_0 = m_0 \left(\frac{1}{2}\right)^{\frac{t}{5730}}$$

$$0.495 = 0.5^{\frac{t}{5730}}$$

$$\log 0.495 = \log 0.5^{\frac{t}{5730}}$$

$$\log 0.495 = \frac{t}{5730} \log 0.5$$

$$\frac{5730 \log 0.495}{\log 0.5} = t$$

$$5813 \approx t$$

Instead of taking the common logarithm of both sides, you could have converted from exponential form to logarithmic form. Try this. Which approach do you prefer? Why?

The oldest buffalo bones found at Head-Smashed-In Buffalo Jump date to about 5813 years ago. The site has been used for at least 6000 years.

**Key Ideas**

- When solving a logarithmic equation algebraically, start by applying the laws of logarithms to express one side or both sides of the equation as a single logarithm.
- Some useful properties are listed below, where  $c, L, R > 0$  and  $c \neq 1$ .
  - If  $\log_c L = \log_c R$ , then  $L = R$ .
  - The equation  $\log_c L = R$  can be written with logarithms on both sides of the equation as  $\log_c L = \log_c c^R$ .
  - The equation  $\log_c L = R$  can be written in exponential form as  $L = c^R$ .
  - The logarithm of zero or a negative number is undefined. To identify whether a root is extraneous, substitute the root into the original equation and check whether all of the logarithms are defined.
- You can solve an exponential equation algebraically by taking logarithms of both sides of the equation. If  $L = R$ , then  $\log_c L = \log_c R$ , where  $c, L, R > 0$  and  $c \neq 1$ . Then, apply the power law for logarithms to solve for an unknown.
- You can solve an exponential equation or a logarithmic equation using graphical methods.
- Many real-world situations can be modelled with an exponential or a logarithmic equation. A general model for many problems involving exponential growth or decay is

$$\text{final quantity} = \text{initial quantity} \times (\text{change factor})^{\text{number of changes}}$$

**Key Ideas**

- Let  $P$  be any real number, and  $M$ ,  $N$ , and  $c$  be positive real numbers with  $c \neq 1$ . Then, the following laws of logarithms are valid.

Name	Law	Description
Product	$\log_c MN = \log_c M + \log_c N$	The logarithm of a product of numbers is the sum of the logarithms of the numbers.
Quotient	$\log_c \frac{M}{N} = \log_c M - \log_c N$	The logarithm of a quotient of numbers is the difference of the logarithms of the dividend and divisor.
Power	$\log_c M^P = P \log_c M$	The logarithm of a power of a number is the exponent times the logarithm of the number.

- Many quantities in science are measured using a logarithmic scale. Two commonly used logarithmic scales are the decibel scale and the pH scale.

## Homework

Chapter 7 Review pg. 366-367 (Do all questions)  
 Chapter 8 Review pg. 416-418 (1-14, 18-20, 22)

### • Chapter 7: Exponential Functions

For  $y = c^x$

D:  $\{x | x \in \mathbb{R}\}$

R:  $\{y | y > \underline{0}, y \in \mathbb{R}\}$

x int: none

y int: (0, 1)

HA:  $y = \underline{0}$

For  $y = a c^{b(x-h)} + k$

D:  $\{x | x \in \mathbb{R}\}$

R:  $\{y | y > k, y \in \mathbb{R}\}$    
 if  $a < 0$ , switch inequality

x int: sub 0 in for y

y int: sub 0 in for x

HA:  $y = k$

### • Chapter 8: Logarithmic Functions

For  $y = \log_c x$

D:  $\{x | x > \underline{0}, x \in \mathbb{R}\}$

R:  $\{y | y \in \mathbb{R}\}$

x int: (1, 0)

y int: none

VA:  $x = \underline{0}$

For  $y = a \log_c (b(x-h)) + k$

D:  $\{x | x > h, x \in \mathbb{R}\}$    
 if  $b < 0$ , switch inequality

R:  $\{y | y \in \mathbb{R}\}$

x int: sub 0 in for y

y int: sub 0 in for x

VA:  $x = h$

