


Formation Reactions:

This reactions starts with elements only as reactants.
The reactants will form compounds as products.

elements \Rightarrow compound




The molar enthalpy symbol for a formation reaction is H_f

Simple Decomposition Reactions:

This reaction starts as a compound, which decomposes into its elements.
(opposite of a formation reaction)


compound \Rightarrow elements



The molar enthalpy symbol is H_{sd} .

Combustion Reactions:

The reaction of a substance with excess oxygen to produce an oxide.



The molar enthalpy symbol is H_c .

(b) Molar Enthalpies

H_f - molar enthalpies of formation is the quantity of heat released or absorbed when one mole of a substance forms from its elements.

H_c - molar enthalpies of combustion is the quantity of heat released or absorbed when one mole of a substance reacts with oxygen.

H° - standard molar enthalpy is the quantity of heat released or absorbed when one mole of a substance reacts at SATP

(c) ΔH_r - Enthalpy change is the quantity of heat released or absorbed when a reaction occurs. This may also be called "Heat of Reaction" or "Change in Heat".

\Rightarrow must know the number of moles of a substance reacting to determine the enthalpy change

(d) Molar enthalpy may be determined from the enthalpy change as long as the number of moles (n) are known.

\longrightarrow

$$6.03 \frac{\text{kJ}}{\text{mol}}$$

$$q = -q$$

$$mC\Delta T = -mC\Delta T \quad *T_f \text{ same}$$

$$(170.9)(0.900 \frac{\text{J}}{\text{g}^\circ\text{C}})(T_f - 105.0^\circ\text{C}) =$$


$$-(115.09)(4.19 \frac{\text{J}}{\text{g}^\circ\text{C}})(T_f - 25.0^\circ\text{C})$$

$$T_f = 44.3^\circ\text{C}$$

$$4(2)(x-1) = -1(2)(x-3)$$

Reaction Enthalpies

$$\Delta H_r = nH_r$$



$$H_f^\circ = -98.79 \text{ kJ/mol}$$

How do we find the change in enthalpy of $\text{SO}_{2(g)}$??

$$\Delta H_c = nH_c$$

$$\Delta H_c = (2 \text{ mol})(-98.79 \text{ kJ/mol})$$

$$\Delta H_c = -197.58 \text{ kJ}$$

$$\Delta H_c = nH_c$$


$$\Delta H_c = (1 \text{ mol})(-98.79 \text{ kJ/mol})$$

$$\Delta H_c = -98.79 \text{ kJ}$$

COMMUNICATING ENTHALPY CHANGES


Using ΔH_r notation:

- for chemical reactions not well known, the chemical equation must accompany the enthalpy change. The molar enthalpy of reaction (or change in enthalpy) follows the equation. **For exothermic reactions the $\Delta H_r < 0$.**







② The Enthalpy Change (ΔH_r) may be included as a term in the balanced equation:

(i) In endothermic reactions - energy is reported as a reactant and is transformed in the reaction.



(ii) In exothermic reactions - energy is reported as a product since it is being produced.







POTENTIAL ENERGY DIAGRAMS

- may be used to express enthalpy change (ΔH_r)
- shows the potential energy of the reactants and products of a chemical reaction.
- shows the difference between the initial and final energies as the enthalpy change. (ΔH_r)

Endothermic Rxn



Exothermic Rxn



see Fig 11-8 p 373 (also 11-15,16,17)

For each of the following reactions:

(a) rewrite the equation including the enthalpy change as a term

(b) draw a potential energy diagram

