

# Warm Up

Differentiate:  $e^{xy^2} = 2x - 3xy + e^{\tan x}$

$$e^{xy^2} (x \cdot 2yy' + (1)y^2) = 2 - (3xy' + 3y) + e^{\tan x} \cdot \sec^2 x (1)$$

$$2xyy'e^{xy^2} + y^2e^{xy^2} = 2 - 3xy' - 3y + e^{\tan x} \sec^2 x$$

$$2xyy'e^{xy^2} + 3xy' = 2 - 3y + e^{\tan x} \sec^2 x - y^2e^{xy^2}$$

$$y' (2xye^{xy^2} + 3x) = 2 - 3y + e^{\tan x} \sec^2 x - y^2e^{xy^2}$$

$$y' = \frac{2 - 3y + e^{\tan x} \sec^2 x - y^2e^{xy^2}}{2xye^{xy^2} + 3x}$$

Find the equation of the tangent line to the curve  $y = 1 + xe^{2x}$  at the point where  $x = 0$ .

① Find  $y$ :

$$y = 1 + xe^{2x}$$

$$y = 1 + (0)e^{2(0)}$$

$$y = 1 + 0$$

$$y = 1$$

$$\text{Point } (0, 1)$$

② Find  $y'$ :

$$y = 1 + xe^{2x}$$

$$y' = 0 + x(e^{2x})(2) + (1)e^{2x}$$

$$y' = 2xe^{2x} + e^{2x}$$

$$y'(0) = 2(0)e^{2(0)} + e^{2(0)}$$

$$= 0 + 1$$

$$= 1$$

↑  
 $m = 1$   
slope of the  
tangent

③ Find equation:

$$y - y_1 = m(x - x_1)$$

$$y - 1 = 1(x - 0)$$

$$y - 1 = x$$

$$0 = x - y + 1$$

Questions from Homework

$$\textcircled{c} e^{xy} = 2x + y$$

$$e^{xy}(xy' + 1y) = 2 + y'$$

$$xy'e^{xy} + ye^{xy} = 2 + y'$$

$$xy'e^{xy} - y' = 2 - ye^{xy}$$

$$y'(xe^{xy} - 1) = 2 - ye^{xy}$$

$$y' = \frac{2 - ye^{xy}}{xe^{xy} - 1}$$

$$\textcircled{d} f(x) = xe^x$$


$$\text{a) } f'(x) = x(e^x)(1) + (1)e^x$$

$$= xe^x + e^x$$

$$= e^x(x+1)$$

$$\text{CV: } e^x = 0 \quad | \quad x+1 = 0$$

$$\text{No CV.} \quad | \quad x = -1$$



$$f(-1) = (-1)e^{(-1)}$$

$$= -1 \cdot \frac{1}{e}$$

$$= -\frac{1}{e}$$

$(-1, -\frac{1}{e})$  abs min

$$\text{b) } f'(x) = e^x(x+1)$$

$$f''(x) = e^x(1) + e^x(1)(x+1)$$

$$= e^x + e^x(x+1)$$

$$= e^x[1 + (x+1)]$$

$$= e^x(x+2)$$



CU on  $(-2, \infty)$   
 CD on  $(-\infty, -2)$

$$\text{CV: } x = -2$$

c) Inflection Point when  $x = -2$

$$f(-2) = (-2)e^{(-2)}$$

$$= -2 \cdot \frac{1}{e^2}$$

$$= -\frac{2}{e^2}$$

$(-2, -\frac{2}{e^2})$  Inflection Point

## Derivatives of Logarithmic Functions

Let's work from the known...

- At this point you should know how to differentiate  $y = e^x$ .

$$\log_e y = x$$

$$\ln y = x$$

What other function could this model?

$$\ln y = x$$

Try to differentiate  $\longrightarrow y = \ln x$ .

$$e^y = x$$

Implicit Diff  $\rightarrow e^y \cdot y' = 1$

$$y' = \frac{1}{e^y} = \frac{1}{x}$$

Differentiate:  $y = \ln x^3$

$$e^y = x^3$$

$$e^y \cdot y' = 3x^2$$
$$y' = \frac{3x^2}{e^y} = \frac{3x^2}{x^3} = \frac{3}{x}$$

$$\text{Rule: } d(\ln u) = \frac{1}{u} du$$

$$u = x^7 \quad du = 7x^6$$

Ex:  $y = \ln x^7$

$$y' = \frac{1}{x^7} \cdot 7x^6 = \frac{7}{x}$$

# Practice Problem:

Page 383

#1

## Laws of Logarithms

$$\log_b M + \log_b N = \log_b(MN)$$

$$\log_b M - \log_b N = \log_b\left(\frac{M}{N}\right)$$

$$\log_b(N^p) = p \log_b(N)$$

We have now covered base "e"...both as an exponential and logarithmic function...

## What about other bases??

Will need to know the change of base formula for logarithms:

$$\log_N M = \frac{\log_b M}{\log_b N}$$

Whatever new base you choose

Differentiate:

$$y = \log_6 x^3$$

$$y = \log(5x^4)$$

**Rule:**  $d(\log_b u) = \frac{1}{u \ln b} du$



This leaves one form of exponential function remaining...

- What about a function such as  $y = 3^{9x}$

Try this one...  $y = \pi^{x^5}$

**Rule:**

$$d(b^u) = b^u (\ln b) du, \text{ where } b \in R$$

# Practice Problems:

Page 383 - 384

#1 #2 a #3 #4

#5 #6 #7 #8