

1. [4 points each] Find  $\frac{dy}{dx}$  for each of the following. Do NOT simplify your answers.

(a)  $y = \frac{6 - x^2 + \sqrt[4]{x}}{x^3 + 4}$


$$y' = \frac{(x^3+4)(-\frac{2x}{1} + \frac{1}{4}x^{-3/4}) - (6-x^2+\sqrt[4]{x})(3x^2)}{(x^3+4)^2}$$

(b)  $y = e^x \sin(3x^4 + 5)$

$$y' = 0x^3 e^x \cos(3x^4 + 5) + e^x \sin(3x^4 + 5)$$

(c)  $y = \sec(3x) + \sqrt{\pi x - 5}$

$$y' = 3\sec(3x)\tan(3x) + \frac{1}{2}(\pi x - 5)^{-1/2}(\pi)$$



(g)  $y = \left(x^4 + \frac{1}{x^4} + \ln 4\right)^{15}$   $\rightarrow -4x^{-5}$

$$y' = 15 \left(x^4 + \frac{1}{x^4} + \ln 4\right)^{14} \left(4x^3 - \frac{4}{x^5}\right)$$

$\uparrow$   
constant

(h)  $y = (x^2 + 10)^{\cos x}$

$$\ln y = \ln (x^2 + 10)^{\cos x}$$

$$\ln y = \cos x \ln (x^2 + 10)$$

$$\frac{y'}{y} = \cos x \left(\frac{2x}{x^2 + 10}\right) + \sin x \ln (x^2 + 10)$$

$$y' = \left[\frac{2x \cos x}{x^2 + 10} + \sin x \ln (x^2 + 10)\right] (x^2 + 10)^{\cos x}$$

(i)  $3xy + y^4 = x^8 + \sinh x$

$$3xy' + 3y + 4y^3 y' = 8x^7 + \cosh x$$

$$y'(3x + 4y^3) = 8x^7 - 3y + \cosh x$$

$$y' = \frac{8x^7 - 3y + \cosh x}{3x + 4y^3}$$

2. [4 points] Find the domain of  $f(x) = \sqrt{x^2 - 2x - 3}$

$x^2 - 2x - 3 \geq 0$  ← cannot take the square root of a negative

$y = (x-3)(x+1)$

$D: \{x \mid x \leq -1 \text{ and } x \geq 3, x \in \mathbb{R}\}$

3. [3 points] Answer (a) and (b) for the function  $f$  defined below.

$$f(x) = \begin{cases} x-1 & \text{if } x < 1; \\ (x-1)^2 & \text{if } x > 1; \\ 3 & \text{if } x = 1. \end{cases}$$

$x-1$	$f(x)$
0	-1
-1	-2

$(x-1)^2$	$f(x)$
0	0
3	4

3	$f(x)$
3	3

(a) Find  $\lim_{x \rightarrow 1} f(x)$ .

$\lim_{x \rightarrow 1^-} f(x)$ $\lim_{x \rightarrow 1^-} (x-1) = 0$	$\lim_{x \rightarrow 1^+} f(x)$ $\lim_{x \rightarrow 1^+} (x-1)^2 = 0$	$\lim_{x \rightarrow 1} f(x) = 0$
---	---	-----------------------------------

(b) Is  $f(x)$  continuous at  $x = 1$ ? Justify your answer.

No because  $\lim_{x \rightarrow 1} f(x) = 0$  and  $f(1) = 3$

(a)  $\lim_{x \rightarrow 0} \frac{e^x - 1}{\cos x}$   
 $\lim_{x \rightarrow 0} \frac{e^x}{-\sin x} = \frac{1}{0} = \text{DNE}$

(b)  $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x^2 - x - 2}$   
 $\lim_{x \rightarrow 2} \frac{2x}{2x - 1} = \frac{4}{3}$

(c)  $\lim_{x \rightarrow \infty} \frac{1 - 3x^2}{x^2 - 5x + 7}$   
 $\lim_{x \rightarrow \infty} \frac{\frac{1}{x^2} - 3}{1 - \frac{5}{x} + \frac{7}{x^2}} = \frac{-3}{1} = -3$

(d)  $\lim_{x \rightarrow 1^-} \frac{x^2 + 1}{x^2 - x}$   
 $\lim_{x \rightarrow 1^-} \frac{x^2 + 1}{x(x-1)} = -\infty$   
 $x=1$   
denominator is a very small negative

Use the limit definition of the derivative to find  $f'(x)$ , given  $f(x) = x^2 - 3x + 5$ .

$$f(x+h) = x^2 + 2xh + h^2 - 3x - 3h + 5$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 3x - 3h + 5 - \cancel{x^2} + 3x - 5}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{x(2x+h-3)}{h} = \boxed{2x-3}$$
  

6. Answer (a)-(c) with regard to the function  $f(x) = \sqrt{x-4}$ .  $\rightarrow D: \{x \geq 4, x \in \mathbb{R}\}$   
 $R: \{y \geq 0, y \in \mathbb{R}\}$

(a) [3 points] Find the inverse function  $f^{-1}(x)$ .

$$y = \sqrt{x-4}$$

$$x = \sqrt{y-4}$$

$$x^2 = y-4$$

$$x^2 + 4 = y$$

$$\boxed{f^{-1}(x) = x^2 + 4}$$

x	f(x)
4	0
5	1
8	2
13	3
20	4

 $\rightarrow$ 


x	f'(x)
0	4
1	5
2	8
3	13
4	20

- (b) [2 points] Find the domain and range of  $f^{-1}(x)$  (the inverse function you found in part (a)).

$$\text{Domain } \{x \mid x \geq 0, x \in \mathbb{R}\}$$

$$\text{Range } \{y \mid y \geq 4, y \in \mathbb{R}\}$$

- (c) [2 points] Graph  $y = f(x)$  and  $y = f^{-1}(x)$  on the same coordinate system (axes).



7. [3 points] Suppose that  $h(x) = f(x)g(x)$  where  $f(2) = 3$ ,  $g(2) = 5$ ,  $f'(2) = -2$ , and  $g'(2) = 4$ . Find  $h'(2)$ .

$$h'(2) = f(2)g'(2) + f'(2)g(2)$$

$$= (3)(4) + (-2)(5)$$

$$= 12 - 10$$

$$= 2$$

8. [5 points] Find the equations of both lines through the point  $(2, -3)$  that are tangent to the parabola  $y = x^2 + x$ .

$y = x^2 + x$   
 $y' = \frac{2x+1}{\uparrow m}$

To Find points of tangency use  $y - y_1 = m(x - x_1)$   
 $y - (-3) = (2x+1)(x-2)$   
 $y + 3 = 2x^2 - 3x - 2$   
 $x^2 + x + 3 = 2x^2 - 3x - 2$   
 $0 = x^2 - 4x - 5$   
 $0 = (x-5)(x+1)$   
 $x = 5$  and  $x = -1$

Point is not on curve  
 $x, y_1$

Slope @  $x = 5$   
 $y' = 2(5) + 1 = 11$   
 Equation:  
 $y + 3 = 11(x - 2)$   
 $y = 11x - 22 - 3$   
 $y = 11x - 25$

Slope @  $x = -1$   
 $y' = 2(-1) + 1 = -1$   
 Equation:  
 $y + 3 = -1(x - 2)$   
 $y = -x - 1$


9. [4 points] Find the critical points of the function  $f(x) = \ln(2 + \sin x)$  on the interval  $[0, 2\pi]$ .

$f(x) = \ln(2 + \sin x)$   
 $f'(x) = \frac{1}{2 + \sin x} \cdot \cos x$   
 $f'(x) = \frac{\cos x}{2 + \sin x}$

CV:  $\cos x = 0$  |  $2 + \sin x = 0$   
 $x = \frac{\pi}{2}, \frac{3\pi}{2}$  |  $\sin x = -2$   
 Not possible



10. [6 points] A 15 foot ladder is leaning on a wall of a house. The bottom of the ladder is pulled away from the base of the wall at a constant rate of 2 feet per minute. At what rate is the top of the ladder sliding down the wall when the bottom of the ladder is 5 feet from the wall?



$$y = \sqrt{15^2 - 5^2}$$

$$y = \sqrt{225 - 25}$$

$$y = \sqrt{200}$$

$$y = 10\sqrt{2}$$

$$x^2 + y^2 = 15^2$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$2(5)(2) + 2(10\sqrt{2}) \frac{dy}{dt} = 0$$

$$20\sqrt{2} \frac{dy}{dt} = -20$$

$$\frac{dy}{dt} = \frac{-1}{\sqrt{2}} \text{ ft/min}$$

The top of the ladder is sliding down the wall at a rate of  $\frac{1}{\sqrt{2}}$  ft/min

$$\text{or } \frac{\sqrt{2}}{2}$$

11. [6 points] Find the area of the largest rectangle that can be inscribed in a right triangle with legs of 3cm and 4cm if two sides of the rectangle lie along the legs.

express with a single variable

$$A = xy$$

$$A = x\left(3 - \frac{3x}{4}\right)$$

$$A = 3x - \frac{3x^2}{4}$$

differentiate

$$A' = 3 - \frac{3}{2}x$$

$$\frac{3}{2}x = 3$$


$$3x = 6$$

$$x = 2$$

$$A = xy$$

$$A = 2\left(\frac{3}{2}\right)$$

$$A = 3 \text{ cm}^2$$



maximize Area  
similar triangles:

$$\frac{3-y}{x} = \frac{3}{4}$$

$$3x = 12 - 4y$$

$$4y = 12 - 3x$$

$$y = 3 - \frac{3}{4}x$$

$$y = 3 - \frac{3}{4}(2)$$

$$y = 3 - \frac{3}{2}$$

$$y = \frac{6-3}{2} = \frac{3}{2}$$

The maximum area is  $3 \text{ cm}^2$ .

Dimensions that maximize area are  $2 \text{ cm} \times 1.5 \text{ cm}$

ownership  
enterprise

12. [13 points] Answer (a)-(h) with regard to the function  $f(x) = \frac{x^2 - 2x + 4}{x - 2}$ . The first and second derivatives of  $f$  are given below.

$$f'(x) = \frac{x(x-4)}{(x-2)^2} \quad f''(x) = \frac{8}{(x-2)^3}$$

(a) Find the intercepts, if any.

$y$  int ( $x=0$ )  
 $y = \frac{4}{-2} = -2$   
 $(0, -2)$

$x$  int ( $y=0$ )  
 $x^2 - 2x + 4 = 0$   
 $x = \frac{2 \pm \sqrt{4-16}}{2}$

$x = \frac{2 \pm 2i\sqrt{3}}{2}$   
 $x = 1 \pm i\sqrt{3}$  → Imaginary Roots  
 No  $x$  intercepts

(b) Find the horizontal and vertical asymptotes, if any.

Horizontal None  
 $\lim_{x \rightarrow \infty} \frac{x^2 - 2x + 4}{x - 2} = \text{DNE}$   
 No HA


Vertical  $x=2$   
 $\lim_{x \rightarrow 2^-} \frac{x^2 - 2x + 4}{x - 2} = -\infty$   
 $\lim_{x \rightarrow 2^+} \frac{x^2 - 2x + 4}{x - 2} = +\infty$

Slant  $y=x$   
 $x \rightarrow \frac{x^2 - 2x + 4}{x - 2} = \frac{x^2 - 2x + 4 - (x^2 - 2x)}{x - 2} = \frac{4}{x - 2}$

(c) Find the critical numbers for  $f$ .

$S(x) = \frac{x(x-4)}{(x-2)^2}$       CV:  $x=0, 2, 4$

(d) Determine the intervals where  $f$  is increasing and the intervals where  $f$  is decreasing.



$f(x) = \frac{x^2 - 2x + 4}{x - 2}$

(e) Find all relative (local) maxima and minima for  $f$ .

<p>max <math>(x=0)</math></p> <p><math>f(0) = -2</math></p> <p><math>(0, -2)</math></p>		<p>min <math>(x=4)</math></p> <p><math>f(4) = \frac{16 - 8 + 4}{2} = 6</math></p> <p><math>(4, 6)</math></p>
---	--	--

(f) Determine the intervals where  $f$  is concave up and the intervals where  $f$  is concave down.

$f''(x) = \frac{8}{(x-2)^3}$

CV:  $x=2$

-	+	
←	→	↔
( )	( )	
( )	( )	

CU on  $(2, \infty)$   
CD on  $(-\infty, 2)$

(g) Find the inflection points, if any.

$f(2) = \frac{4 - 4 + 4}{0} = \text{undefined}$

There is no inflection point at  $x=2$  because there is a vertical asymptote here

(h) Sketch the graph of  $y = f(x)$ . Label the intercepts, asymptotes, relative maxima and minima, and inflection points on the graph.