

Example 3

Solve Problems Involving Exponential Equations With Different Bases

Christina plans to buy a car. She has saved \$5000. The car she wants costs \$5900. How long will Christina have to invest her money in a term deposit that pays 6.12% per year, compounded quarterly, before she has enough to buy the car?

Solution

The formula for compound interest is $A = P(1 + i)^n$, where A is the amount of money at the end of the investment; P is the principal amount deposited; i is the interest rate per compounding period, expressed as a decimal; and n is the number of compounding periods.

In this problem:

$$A = \underline{5900}$$

$$P = \underline{5000}$$

$$i = 0.0612 \div 4 \text{ or } \underline{\underline{0.0153}}$$

Divide the interest rate by 4 because interest is paid quarterly or four times a year.

$$\begin{aligned} &\rightarrow 6.12\% \\ &= 0.0612 \div 4 \\ &= 0.0153 \end{aligned}$$

$$A = P(1+i)^n$$

$$\frac{5900}{5000} = \frac{5000}{5000} (1.0153)^n$$

$$1.18 = (1.0153)^n$$

$$(\cancel{1.0153})^{10.9} = (\cancel{1.0153})^n$$

* Express 1.18 with a base of 1.0153 $\rightarrow \frac{\log 1.18}{\log 1.0153} = \underline{\underline{10.9}}$

$$10.9 = n$$

\swarrow n is the number of compound periods ... we want to know the number of years.

$$\hookrightarrow 10.9 \div 4 = \boxed{2.73 \text{ years}}$$

The number of milligrams of a drug remaining in the bloodstream t days after consumption is given by the equation:

$$D = 50(0.9)^t$$

- (a) What percentage of the drug leaves the body each day? 10 %
- (b) The drug can be detected in urine tests when 2 or more mg of the drug remain in the bloodstream. Will there be evidence of this drug in the bloodstream 28 days after consumption? Provide proof! $t=28$

$$D = 50(0.9)^{28}$$
$$D = 2.62 \text{ mg}$$


?

Understanding Logarithms

Focus on...

- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- determining the characteristics of the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

For the exponential function $y = c^x$, the inverse is $x = c^y$. This inverse is also a function and is called a **logarithmic function**. It is written as $y = \log_c x$, where c is a positive number other than 1.



Since our number system is based on powers of 10, **logarithms** with base 10 are widely used and are called **common logarithms**. When you write a common logarithm, you do not need to write the base. For example, $\log 3$ means $\log_{10} 3$.

logarithmic function

- a function of the form $y = \log_c x$, where $c > 0$ and $c \neq 1$, that is the inverse of the exponential function $y = c^x$

logarithm

- an exponent
- in $x = c^y$, y is called the logarithm to base c of x

common logarithm

- a logarithm with base 10

Write each of the following in logarithmic form

a) $32 = 2^5$ ← exp.

↑ ↑
ans. base

$\log_2(32) = 5$

or $5 = \log_2 32$

b) $2^{-5} = \frac{1}{32}$

$\log_2\left(\frac{1}{32}\right) = -5$

c) $x = 10^y$

$\log_{10} x = y$

or $y = \log x$

Logarithmic Form

Exponential Form



Write each of the following in exponential form

a) $\log_4 16 = 2$ ← ans ← exp.

↑
base

$4^2 = 16$

b) $\log_2\left(\frac{1}{32}\right) = -5$

$2^{-5} = \frac{1}{32}$

c) $\log 65 = 1.8129$

$10^{1.8129} = 65$

Example 1

Evaluating a Logarithm

Evaluate.

a) $\log_7 49$

Let $x = \log_7 49$

$$7^x = 49 \text{ (exp. form)}$$

$$7^x = 7^2 \text{ (get common base)}$$

$$x = 2$$

$$\boxed{\log_7 49 = 2}$$

b) $\log_6 1$

Let $x = \log_6 1$

$$6^x = 1$$

$$6^x = 6^0$$

$$x = 0$$

$$\boxed{\log_6 1 = 0}$$

c) $\log 0.001$

Let $x = \log 0.001$

$$10^x = 0.001$$

$$10^x = 10^{-3}$$

$$x = -3$$

$$\boxed{\log 0.001 = -3}$$

d) $\log_2 \sqrt{8}$

Let $x = \log_2 \sqrt{8}$

Example 2

Determine an Unknown in an Expression in Logarithmic Form

Determine the value of x .

a) $\log_5 x = -3$

b) $\log_x 36 = 2$

c) $\log_{64} x = \frac{2}{3}$

a) $5^{-3} = x$

$\left(\frac{1}{5}\right)^3 = x$

$\frac{1}{125} = x$

b) $x^2 = 36$

c) $64^{\frac{2}{3}} = x$

Exponential Function \leftarrow (Inverse) \rightarrow Logarithmic Function

$$y = c^x, c > 0, c \neq 1$$

$$D: \{x \mid x \in \mathbb{R}\}$$

$$R: \{y \mid y > 0, y \in \mathbb{R}\}$$

$$HA: y = 0$$

$$x \text{ int: none}$$

$$y \text{ int: } (0, 1)$$

$$y = \log_c x, c > 0, c \neq 1$$

$$D: \{x \mid x > 0, x \in \mathbb{R}\}$$

$$R: \{y \mid y \in \mathbb{R}\}$$

$$VA: x = 0$$

$$x \text{ int: } (1, 0)$$

$$y \text{ int: none}$$

Example 3



Graph the Inverse of an Exponential Function

- a) State the inverse of $f(x) = 3^x$.
- b) Sketch the graph of the inverse. Identify the following characteristics of the inverse graph:
 - the domain and range
 - the x -intercept, if it exists
 - the y -intercept, if it exists
 - the equations of any asymptotes

Solution


a) The inverse of $y = f(x) = 3^x$ is _____ or, _____ expressed in logarithmic form, _____. Since the inverse is a function, it can be written in function notation as _____

How do you know that $y = \log_3 x$ is a function?

b) Set up tables of values for both the exponential function, $f(x)$, and its inverse, $f^{-1}(x)$. Plot the points and join them with a smooth curve.

$f(x) = 3^x$	
x	y
-3	
-2	
-1	
0	
1	
2	
3	

$f^{-1}(x) = \log_3 x$	
x	y



The graph of the inverse, $f^{-1}(x) = \log_3 x$, is a reflection of the graph

of $f(x) = 3^x$ about the line $y = x$. For $f^{-1}(x) = \log_3 x$,

- the domain is _____ and the range is _____
- the x-intercept is _____
- there is no y-intercept
- the vertical asymptote, the _____ axis, has equation _____ there is no _____ asymptote

How do the characteristics of $f^{-1}(x) = \log_3 x$ compare to the characteristics of $f(x) = 3^x$?

Key Ideas


- A logarithm is an exponent.
- Equations in exponential form can be written in logarithmic form and vice versa.

Exponential Form **Logarithmic Form**

$$x = c^y \qquad y = \log_c x$$

- The inverse of the exponential function $y = c^x$, $c > 0$, $c \neq 1$, is $x = c^y$ or, in logarithmic form, $y = \log_c x$. Conversely, the inverse of the logarithmic function $y = \log_c x$, $c > 0$, $c \neq 1$, is $x = \log_c y$ or, in exponential form, $y = c^x$.
- The graphs of an exponential function and its inverse logarithmic function are reflections of each other in the line $y = x$, as shown.
- For the logarithmic function $y = \log_c x$, $c > 0$, $c \neq 1$,
 - the domain is $\{x \mid x > 0, x \in \mathbb{R}\}$
 - the range is $\{y \mid y \in \mathbb{R}\}$
 - the x-intercept is 1
 - the vertical asymptote is $x = 0$, or the y-axis
- A common logarithm has base 10. It is not necessary to write the base for common logarithms:


$$\log_{10} x = \log x$$




Homework

#1-5, 8, 10, 12, 13, 17 on page 380

8.1 Understanding Logarithms, pages 380 to 382




ii) $y = \log_2 x$
 iii) domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1, no y-intercept,
 vertical asymptote $x = 0$



ii) $y = \log_3 x$
 iii) domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1, no y-intercept,
 vertical asymptote $x = 0$

2. a) $\log_{12} 144 = 2$ b) $\log_8 2 = \frac{1}{3}$
 c) $\log_{10} 0.000\ 01 = -5$ d) $\log_7 (y + 3) = 2x$
3. a) $5^2 = 25$ b) $8^{\frac{2}{3}} = 4$
 c) $10^6 = 1\ 000\ 000$ d) $11^y = x + 3$
4. a) 3 b) 0 c) $\frac{1}{3}$ d) -3
5. $a = 4; b = 5$



domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1,
 no y-intercept,
 vertical asymptote $x = 0$

10. They are reflections of each other in the line $y = x$.
11. a) They have the exact same shape.
 b) One of them is increasing and the other is decreasing.
12. a) 216 b) 81 c) 64 d) 8
13. a) 7 b) 6
14. a) 0 b) 1
15. -1
16. 16
17. a) $t = \log_{0.11} N$ b) 145 days
18. The larger asteroid had a relative risk that was 1479 times as dangerous.
19. 1000 times as great
20. 5
21. $m = 14, n = 13$
22. $4n$
23. $y = 3^{2^x}$

