

Differential and Integral Calculus 120

[Substitution Rule!]

Substitution Rule

It often arises that we need to integrate a function which does not follow one of our basic integral formulas therefore other techniques are necessary.

One such technique is called Substitution that involves a change of variable, which permits us to rewrite an integrand in a form to which we can apply a basic integration rule.

Substitution Rule for Indefinite Integrals

If $u = g(x)$, then $\int f(g(x))g'(x)dx = \int f(u)du$

Substitution Rule for Integration corresponds...

Let's do an example... to the Chain Rule for Differentiation.

$$\text{Find } \int (x^2 - 5)^8 2x dx$$

In using the substitution rule, the idea is to replace a complicated integral by a simpler integral by changing to a new variable u .

In thinking of the appropriate substitution, we try to choose u to be some function in the integrand whose differential du also occurs. (ignoring any constants)

So using the above example we have...

$$\int (x^2 - 5)^8 2x dx$$

Let:

$$u = x^2 - 5 \quad du = 2x dx$$

$$\int u^8 du$$

$$= \frac{u^9}{9} + C$$

$$= \frac{(x^2 - 5)^9}{9} + C$$

Evaluate each of the following:

$$\int \frac{x^2}{\sqrt{1-x^3}} dx$$

$$u = 1-x^3$$

$$du = -3x^2 dx$$

$$-\frac{1}{3} du = x^2 dx$$

$$= \int \frac{x^2}{(1-x^3)^{\frac{1}{2}}} dx$$

$$= -\frac{1}{3} \int (u)^{-\frac{1}{2}} \cdot \left(-\frac{1}{3} du\right)$$

$$= -\frac{1}{3} \cdot 2u^{\frac{1}{2}} + C$$

$$= -\frac{2}{3}(1-x^3)^{\frac{1}{2}} + C$$

$$= -\frac{2}{3}(1-x^3)^{1/2} + C$$

$$\int \sin 4x dx$$

$$u = 4x \quad = \int \sin u \cdot \left(\frac{1}{4} du\right)$$

$$du = 4dx$$

$$\frac{1}{4} du = dx \quad = \frac{1}{4} \int \sin(u) du$$

$$= \frac{1}{4} \cdot -\cos u + C$$

$$= -\frac{1}{4} \cos u + C$$

$$= -\frac{1}{4} \cos 4x + C$$

$$= -\frac{1}{4} \cos 4x + C$$

Evaluate each of the following:

$$\int \frac{\ln x}{x} dx = \int \ln x \cdot \frac{1}{x} dx$$

$$\begin{aligned} u &= \ln x &= \int u du \\ du &= \frac{1}{x} dx &= \frac{u^2}{2} + C \\ &&= \frac{(\ln x)^2}{2} + C \\ &= \frac{1}{2} (\ln x)^2 + C \end{aligned}$$

$$\int (2 + \sin x)^{10} \cos x dx$$

$$\begin{aligned} u &= 2 + \sin x &= \int u^{10} du \\ du &= \cos x dx &= \frac{u^{11}}{11} + C \\ &&= \frac{(2 + \sin x)^{11}}{11} + C \end{aligned}$$

$$= \frac{1}{11} (2 + \sin x)^{11} + C$$

Evaluate each of the following:

$$\int \underline{2x} \sqrt{1+x^2} dx$$

$u = 1+x^2$ $= \int \underline{u}^{\frac{1}{2}} du$
 $du = 2x dx$ $= \frac{2}{3} u^{\frac{3}{2}} + C$
 $= \frac{2}{3} (1+x^2)^{\frac{3}{2}} + C$
 $= \frac{2}{3} (x^2 + 1)^{3/2} + C$

$$\int \underline{x^3} \cos(\underline{x^4 + 2}) dx$$

$u = x^4 + 2$ $= \int \cos(u) \cdot \frac{1}{4} du$
 $du = 4x^3 dx$
 $\frac{1}{4} du = x^3 dx$
 $= \frac{1}{4} \int \cos(u) du$
 $= \frac{1}{4} \sin u + C$
 $= \frac{1}{4} \sin(x^4 + 2) + C$
 $= \frac{1}{4} \sin(x^4 + 2) + C$

Evaluate each of the following:

$$\int \sqrt{2x+1} dx$$

$u = 2x+1$
 $du = 2dx$
 $\frac{1}{2} du = dx$

$$= \int u^{\frac{1}{2}} \cdot \frac{1}{2} du$$

$$= \frac{1}{2} \int u^{\frac{1}{2}} du$$

$$= \frac{1}{2} \cdot \frac{2}{3} u^{\frac{3}{2}} + C$$

$$= \frac{1}{3} u^{\frac{3}{2}} + C$$

$$= \frac{1}{3} (2x+1)^{\frac{3}{2}} + C$$

$$\int \frac{x}{\sqrt{1-4x^2}} dx$$

$u = 1-4x^2$
 $du = -8x dx$
 $-\frac{1}{8} du = x dx$

$$= \int u^{-\frac{1}{2}} \cdot \left(-\frac{1}{8} du\right)$$

$$= -\frac{1}{8} \int u^{-\frac{1}{2}} du$$

$$= -\frac{1}{8} \cdot 2u^{\frac{1}{2}} + C$$

$$= -\frac{1}{4} u^{\frac{1}{2}} + C$$

$$= -\frac{1}{4} \sqrt{1-4x^2} + C$$

$$= -\frac{1}{4} \sqrt{1-4x^2} + C$$

Evaluate the following:

$$\begin{aligned}
 & \int e^{5x} dx \\
 \underline{u=5x} \quad & = \int e^u \cdot \left(\frac{1}{5} du\right) \\
 \underline{du=5dx} \quad & = \frac{1}{5} \int e^u du \\
 \underline{\frac{1}{5} du = dx} \quad & = \frac{1}{5} e^u + C \\
 & = \frac{1}{5} e^{5x} + C \\
 & = \frac{1}{5} e^{5x} + C
 \end{aligned}$$

Evaluate:

$$\int \sqrt{1+x^2} x^5 dx$$

appropriate substitution becomes more obvious if we factor x^5 as $x^4 \cdot x$

$$\begin{aligned}
 & \begin{array}{l} u = 1+x^2 \\ du = 2x dx \\ \frac{1}{2} du = x dx \end{array} & \begin{aligned} & \int \sqrt{1+x^2} \cdot \underline{\underline{x^4}} \cdot \underline{\underline{x dx}} \\ &= \int u^{\frac{1}{2}} \cdot (u-1)^3 \cdot \left(\frac{1}{2} du\right) \\ &= \frac{1}{2} \int u^{\frac{1}{2}} \cdot (u-1)^3 du \\ &= \frac{1}{2} \int u^{\frac{1}{2}} (u^3 - 3u^2 + 3u - 1) du \\ &= \frac{1}{2} \int u^{\frac{5}{2}} - 3u^{\frac{7}{2}} + 3u^{\frac{9}{2}} - u^{\frac{11}{2}} du \\ &= \frac{1}{2} \left(\frac{2}{7} u^{\frac{7}{2}} - \frac{3}{5} u^{\frac{9}{2}} + \frac{3}{3} u^{\frac{11}{2}} \right) + C \\ &= \frac{1}{7} u^{\frac{7}{2}} - \frac{3}{5} u^{\frac{9}{2}} + \frac{1}{3} u^{\frac{11}{2}} + C \\ &= \frac{1}{7} (1+x^2)^{7/2} - \frac{3}{5} (1+x^2)^{5/2} + \frac{1}{3} (1+x^2)^{3/2} + C \end{aligned}
 \end{aligned}$$

Homework - Exercise 11.3 - pp. 511-512 - Q. 1 - 3

RED BOOK

Let's do one where the higher degree is not under the radical, but outside... how do we handle this????

$$\int x^2 \sqrt{8x+5} dx = \frac{(8x+5)^{\frac{3}{2}}}{672} [24x^2 - 12x + 5] + C$$

So, $\int x^2 \sqrt{8x+5} dx$ well let $u = \sqrt{8x+5}$ therefore
 $u^2 = 8x+5$ and $\frac{u^2-5}{8} = x$
 So

$2udu = 8dx$
 And

$$\frac{1}{4}udu = dx$$

Now let's start the integration:

$$\begin{aligned} &= \int x^2 \sqrt{8x+5} dx \\ &= \int \left(\frac{u^2-5}{8} \right)^2 u \left(\frac{1}{4}udu \right) \\ &= \int \left(\frac{u^4 - 10u^2 + 25}{64} \right) \left(\frac{u^2}{4} du \right) \\ &= \int \frac{u^6 - 10u^4 + 25u^2}{256} du \\ &= \frac{1}{256} \int u^6 - 10u^4 + 25u^2 du \end{aligned}$$

Now integrate, I brought out the 1/256 (makes it funnier lol.....)

$$= \frac{1}{256} \left(\frac{u^7}{7} - 2u^5 + \frac{25u^3}{3} \right) + C$$

So, now fill in for u and simplify.....

$$\begin{aligned} &= \frac{1}{256} \left(\frac{(8x+5)^{\frac{7}{2}}}{7} - 2(8x+5)^{\frac{5}{2}} + \frac{25(8x+5)^{\frac{3}{2}}}{3} \right) + C \\ &\quad \text{a } 1/21 \text{ as well as } (8x+5)^{3/2} \\ &= \frac{(8x+5)^{\frac{3}{2}}}{5376} [3(8x+5)^2 - 42(8x+5) + 175] + C \\ &\quad \text{and get.....} \\ &= \frac{(8x+5)^{\frac{3}{2}}}{5376} [192x^2 - 96x + 40] + C \\ &\quad \text{reduce} \\ &= \frac{(8x+5)^{\frac{3}{2}}}{672} [24x^2 - 12x + 5] + C \end{aligned}$$

maybe now factor out
mulphy this out
and then you can
amazing!

Questions From Homework

$$\begin{aligned}
 & \textcircled{2} \quad \int \frac{x+1}{x^2+2x-6} dx \\
 & \qquad u = x^2 + 2x - 6 \\
 & \qquad du = 2x + 2 dx \\
 & \qquad \frac{1}{2} du = x + 1 dx \\
 & = \int \frac{1}{u} \cdot \left(\frac{1}{2} du \right) \\
 & = \frac{1}{2} \int \frac{1}{u} du \\
 & = \frac{1}{2} \ln|u| + C
 \end{aligned}$$

When performing a definite integration using substitution, we have to change the limits of integration so that they are the appropriate values of u.

Substitution Rule for Definite Integrals

$$\text{If } u = g(x), \text{ then } \int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du$$

Example...

$$\begin{aligned}
 \int_1^9 \frac{e^{\sqrt{x}}}{\sqrt{x}} dx &= \int_1^3 e^u \cdot 2du \\
 u = \sqrt{x} &= x^{\frac{1}{2}} \\
 du = \frac{1}{2}x^{-\frac{1}{2}}dx &= \frac{1}{2}\frac{dx}{dx} \\
 2du = \frac{1}{\sqrt{x}}dx &= 2e^u \Big|_1^3 \\
 \frac{x}{u} \Big|_1^9 &= 2e^3 - 2e^1 = 2(e^3 - e) \\
 1 \Big| \sqrt{1} = 1 &= 2e(e^2 - 1) \\
 &= 2e(e+1)(e-1)
 \end{aligned}$$

Find the area under the curve....

$$\begin{aligned}
 A &= \int_0^1 \frac{1}{2x+1} dx = \int_1^3 \frac{1}{u} \cdot \left(\frac{1}{2}du\right) \\
 u = 2x+1 &= \frac{1}{2} \int_1^3 \frac{1}{u} du \\
 du = 2dx &= \frac{1}{2} \ln|u| \Big|_1^3 \\
 \frac{1}{2}du = dx &= \frac{1}{2} \ln 3 - \frac{1}{2} \ln 1 \\
 \frac{x}{u} \Big|_0^1 &= \frac{1}{2} \ln 3 - \frac{1}{2} \ln(0) \\
 1 \Big| \frac{1}{2(0)+1} = 1 &= \boxed{\frac{1}{2} \ln 3}
 \end{aligned}$$

$$= \frac{1}{2} \ln 3$$

Evaluate:

$$= \frac{26}{3}$$

$$\int_0^4 \sqrt{2x+1} dx$$

$$u = 2x+1$$

$$du = 2dx$$

$$\frac{1}{2}du = dx$$

$$\begin{array}{c|cc} x & u \\ \hline \frac{1}{4} & 9 \\ 0 & 1 \end{array}$$

$$= \int_1^9 u^{\frac{1}{2}} \cdot \left(\frac{1}{2}du\right)$$

$$= \frac{1}{2} \int_1^9 u^{\frac{1}{2}} du$$

$$= \frac{1}{2} \cdot \frac{2}{3} u^{\frac{3}{2}} \Big|_1^9$$

$$= \frac{1}{3} u^{\frac{3}{2}} \Big|_1^9$$

$$= \frac{1}{3} (9)^{\frac{3}{2}} - \frac{1}{3} (1)^{\frac{3}{2}}$$

$$= \frac{27}{3} - \frac{1}{3}$$

$$= \boxed{\frac{26}{3}}$$

$$= \frac{1}{14}$$

$$\int_1^2 \frac{1}{(3-5x)^2} dx$$

$$u = 3-5x$$

$$du = -5dx$$

$$\frac{-1}{5}du = dx$$

$$\begin{array}{c|cc} x & u \\ \hline \frac{1}{2} & -7 \\ 1 & -2 \end{array}$$

$$= \int_{-2}^{-7} u^{-2} \cdot \left(-\frac{1}{5}du\right)$$

$$= -\frac{1}{5} \int_{-2}^{-7} u^{-2} du$$

$$= -\frac{1}{5} \cdot -1 u^{-1} \Big|_{-2}^{-7}$$

$$= \frac{1}{5u} \Big|_{-2}^{-7}$$

$$= \frac{1}{5(-7)} - \frac{1}{5(-2)}$$

$$= -\frac{1}{35} + \frac{1}{10}$$

$$= \frac{-2 + 7}{70}$$

$$= \frac{5}{70}$$

$$= \boxed{\frac{1}{14}}$$

Evaluate:

$$\int_1^e \frac{\ln x}{x} dx = \int_0^1 u du = \left[\frac{u^2}{2} \right]_0^1$$

$$u = \ln x$$

$$du = \frac{1}{x} dx$$

$$\begin{array}{c} x \\ \cancel{x} \\ e \\ \hline 1 \end{array} \begin{array}{c} u \\ \cancel{u} \\ 1 \\ \hline 0 \end{array}$$

$$= \frac{(1)^2}{2} - \frac{(0)^2}{2}$$

$$= \frac{1}{2} - 0$$

$$= \boxed{\frac{1}{2}}$$

$$= \frac{1}{2}$$

Homework - Exercise 11.3 - pp. 511-512 - Q. 4 and 6

RED BOOK