

### Questions from Quiz

$$\textcircled{1} \quad f(x) = \sqrt{3x+1} \quad f(x+h) = \sqrt{3(x+h)+1} = \sqrt{3x+3h+1}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\sqrt{3x+3h+1} - \sqrt{3x+1}}{h} \cdot \frac{(\sqrt{3x+3h+1} + \sqrt{3x+1})}{(\sqrt{3x+3h+1} + \sqrt{3x+1})}$$

$$= \lim_{h \rightarrow 0} \frac{3h+1 - (3x+1)}{h(\sqrt{3x+3h+1} + \sqrt{3x+1})}$$

$$= \lim_{h \rightarrow 0} \frac{3h}{h(\sqrt{3x+3h+1} + \sqrt{3x+1})} = \frac{3}{2\sqrt{3x+1}}$$

$$\textcircled{2} \text{ b) } f'(x) = \frac{1}{2x^{1/2}} + \frac{1}{3x^{2/3}} + \frac{1}{4x^{3/4}} \cdot (2x^3 + 5x)^3$$

$$\textcircled{4} \quad f(x) = \frac{x^2}{\sqrt{x-3}}$$

$$f'(x) = \frac{2x(\sqrt{x-3}) - x^2 \left( \frac{1}{2\sqrt{x}} \right)}{(\sqrt{x-3})^2}$$

$$f'(x) = \frac{2x^{\frac{3}{2}} - 6x^{\frac{1}{2}} - \frac{x^2}{2\sqrt{x}} \cdot 2\sqrt{x}}{2\sqrt{x}(\sqrt{x-3})^2}$$

$$f'(x) = \frac{4x^{\frac{3}{2}} - 12x^{\frac{1}{2}} - x^2}{2\sqrt{x}(\sqrt{x-3})^2}$$

$$f'(x) = \frac{3x^{\frac{3}{2}} - 12x^{\frac{1}{2}}}{2\sqrt{x}(\sqrt{x-3})^2} = \frac{3x^{\frac{3}{2}}(x^{\frac{1}{2}} - 4)}{2x^{\frac{1}{2}}(x^{\frac{1}{2}} - 3)^2} = \frac{3x(\sqrt{x} - 4)}{2(\sqrt{x} - 3)^2}$$

$$\textcircled{5} \quad y = (3x+4)^3 \quad \text{Find equation of tangent when } x=-1$$

(i) when  $x = -1$   
 $y = (2(-1)^2 + 5(-1))^3 (3(-1) + 4)^3$

(ii)  $y = (2x^2 + 5x)^3 (3x + 4)^3$

$y = (-27)(1)$

$y = -27$

Point  $\rightarrow (-1, -27)$

(iii)  $y'(-1)$

$y'(-1) = 3(-3)(1)(1)^2 + (-3)^2(2)(1)(3)$

$m = y'(-1) = 27 + (-162) = -135$   $\rightarrow$  Slope

(iv)  $y - y_1 = m(x - x_1)$

$y + 27 = -135(x + 1)$

$y + 27 = -135x - 135$

$135x + y + 162 = 0$

**Questions from Quiz**

- 6.) Find the points on the curve  $y = \frac{x}{x-1}$  where the tangent line is parallel to the line  $x + 4y = 1$ . (6)

$$\begin{array}{lll} \textcircled{1} \quad 4y = -x + 1 & \textcircled{2} \quad y = \frac{x}{x-1} & \textcircled{3} \quad \frac{-1}{(x-1)^2} = -\frac{1}{4} \\ y = -\frac{1}{4}x + \frac{1}{4} & m = y' = \frac{1(x-1) - 1(x)}{(x-1)^2} & (x-1)^2 = 4 \\ m = -\frac{1}{4} & m = y' = \frac{x-1-x}{(x-1)^2} & x^2 - 2x + 1 = 4 \\ & m = y' = \frac{-1}{(x-1)^2} & x^2 - 2x - 3 = 0 \\ & & (x-3)(x+1) = 0 \\ & & x-3=0 \quad | \quad x+1=0 \\ & & x=3 \quad | \quad x=-1 \end{array}$$

\textcircled{4} when  $x=3$     \textcircled{5} when  $x=-1$

$$\begin{array}{ll} y = \frac{3}{3-1} = \frac{3}{2} & y = \frac{-1}{-1-1} = \frac{1}{2} \\ (3, \frac{3}{2}) & (-1, \frac{1}{2}) \end{array}$$

$$\begin{array}{l} \textcircled{6} \quad f(x) = \frac{x^3}{\sqrt{x}-3} \\ f'(x) = \frac{\cancel{2x}(\sqrt{x}-3) - x^2 \left( \frac{1}{\cancel{2\sqrt{x}}} \right)}{(\sqrt{x}-3)^2} \end{array}$$

$$f'(x) = \frac{3x^2 \cancel{\sqrt{x}} - 6x \cancel{\sqrt{x}} - x^2 \cdot \cancel{\sqrt{x}}}{2\cancel{\sqrt{x}}(\sqrt{x}-3)^2}$$

$$f'(x) = \frac{3x^2 - 12x - x^2}{2\sqrt{x}(\sqrt{x}-3)^2}$$

$$f'(x) = \frac{3x^2 - 12x^{\frac{1}{2}}}{2\sqrt{x}(\sqrt{x}-3)^2}$$

$$f'(x) = \frac{3x^{\frac{3}{2}}(x^{\frac{1}{2}} - 4)}{2x^{\frac{1}{2}}(x^{\frac{1}{2}} - 3)^2} = \frac{3x(\sqrt{x} - 4)}{2(\sqrt{x} - 3)^2}$$

$$\textcircled{7b} \quad f(x) = x^{\frac{1}{2}} + x^{\frac{1}{3}} + x^{\frac{1}{4}}$$

$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}} + \frac{1}{3}x^{-\frac{2}{3}} + \frac{1}{4}x^{-\frac{3}{4}}$$

$$f'(x) = \frac{1}{2x^{\frac{1}{2}}} + \frac{1}{3x^{\frac{2}{3}}} + \frac{1}{4x^{\frac{3}{4}}}$$

## Questions from Homework

④ Find  $\frac{dy}{dt} \Big|_{t=1}$  if  $y = \sqrt{1+r^3}$  and  $r = \frac{t+1}{2t+1}$

when  $t=1$ 

$$y = (1+r^3)^{\frac{1}{3}}$$

$$r = \frac{2}{3}$$

$$\frac{dy}{dr} = \frac{1}{3}(1+r^3)^{-\frac{2}{3}}(2r) \quad \frac{dr}{dt} = \frac{1(2t+1) - 2(t+1)}{(2t+1)^2}$$

$$\frac{dy}{dt} = \frac{\cancel{2r}}{\cancel{3}(1+r^3)^{\frac{2}{3}}} \quad \frac{dr}{dt} = \frac{2t+1 - 2t - 2}{(2t+1)^2}$$

$$\frac{dy}{dt} = \frac{r}{\sqrt{1+r^3}} \quad \frac{dr}{dt} = \frac{-1}{(2t+1)^2}$$

$$\frac{dy}{dt} \Big|_{t=1} = \left[ \frac{r}{\sqrt{1+r^3}} \right] \left[ \frac{-1}{(2t+1)^2} \right]$$

$$= \left[ \frac{\left(\frac{2}{3}\right)}{\sqrt{1+\left(\frac{2}{3}\right)^3}} \right] \left[ \frac{-1}{9} \right]$$

$$= \left[ \frac{\frac{2}{3}}{\sqrt{\frac{13}{9}}} \right] \left[ \frac{-1}{9} \right]$$

$$= \left[ \frac{\frac{2}{3}}{\frac{\sqrt{13}}{3}} \right] \left[ \frac{-1}{9} \right]$$

$$= \left[ \frac{\frac{2}{3} \cdot \frac{3}{\sqrt{13}}}{\frac{1}{9}} \right] \left[ -\frac{1}{9} \right]$$

$$= -\frac{2}{9\sqrt{13}}$$

⑥ b)  $f(x) = (2x+1)(4x-1)^5$

$$f'(x) = 2(4x-1)^5 + (2x+1)(5)(4x-1)^4(4)$$

$$f'(x) = 2(4x-1)^5 + 20(2x+1)(4x-1)^4$$

$$f'(x) = 2(4x-1)^4 [4x-1 + 10(2x+1)]$$

$$f'(x) = 2(4x-1)^4 (4x-1 + 20x+10)$$

$$f'(x) = 2(4x-1)^4 (24x+9)$$

$$f'(x) = 6(4x-1)^4 (8x+3)$$

## Questions from Homework

$$\begin{aligned}
 & \text{If } t=4 \\
 & v = 3(4) - \sqrt{4} = 12 - 2 = \underline{\underline{10}} \\
 \textcircled{5} \text{ Find } \frac{ds}{dt} \Big|_{t=4} & \text{ if } s = v + \frac{50}{v} \quad \text{and } v = 3t - \sqrt{t} \\
 & s = v + 50v^{-1}
 \end{aligned}$$

## Differentiation Rules

### Product Rule:

**The Product Rule** If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx} [f(x)g(x)] = f(x) \frac{d}{dx} [g(x)] + g(x) \frac{d}{dx} [f(x)]$$

Express the product rule verbally if you are considering a function of the form...

$$f(x) = (\text{First}) \times (\text{Second})$$

" The derivative of the product of two functions is the the first multiplied by the derivative of second, plus the derivative of first multiplied by the second"

*Get in the habit of verbalizing the rule as you differentiate...it will help when the functions get more complicated.*

## Quotient Rule:

**The Quotient Rule** If  $f$  and  $g$  are differentiable, then

$$\frac{d}{dx} \left[ \frac{f(x)}{g(x)} \right] = \frac{g(x) \frac{d}{dx} [f(x)] - f(x) \frac{d}{dx} [g(x)]}{[g(x)]^2}$$

Express the quotient rule verbally ...

"The denominator multiplied by the derivative of the numerator, minus the numerator multiplied by the derivative of the denominator, all over the denominator squared"

## Combining the Chain Rule With the Product and Quotient Rule:

**The Chain Rule** If  $f$  and  $g$  are both differentiable and  $F = f \circ g$  is the composite function defined by  $F(x) = f(g(x))$ , then  $F$  is differentiable and  $F'$  is given by

$$\text{the product} \quad F'(x) = f'(g(x))g'(x)$$

Differentiate the following function and simplify your answer:

$$y = (x^2 + 1)^3(2 - 3x)^4$$

$$\begin{aligned} y' &= (x^2 + 1)^3(4)(2 - 3x)^3(-3) + 3(x^2 + 1)^2(2x)(2 - 3x)^4 \\ &= -12(x^2 + 1)^3(2 - 3x)^3 + 6x(x^2 + 1)^2(2 - 3x)^4 \\ &= -6(x^2 + 1)^2(2 - 3x)^3 \left[ 2\cancel{(x^2 + 1)} - x\cancel{(2 - 3x)} \right] \\ &= -6(x^2 + 1)^2(2 - 3x)^3 [2x^2 + 2 - 2x + 3x^2] \\ &= \boxed{-6(x^2 + 1)^2(2 - 3x)^3 (5x^2 - 2x + 2)} \end{aligned}$$

$$g(x) = \frac{(3x+2)^2}{2x}$$

$$\begin{aligned} g'(x) &= \frac{2x(2)(3x+2)(3) - (3x+2)^2(2)}{(2x)^3} \\ &= \frac{12x(3x+2) - 2(3x+2)^2}{4x^3} \\ &= \frac{2(3x+2)[6x - \cancel{(3x+2)}]}{4x^3} \\ &= \frac{2(3x+2)(3x-2)}{4x^3} \\ &= \boxed{\frac{(3x+2)(3x-2)}{2x^3}} \quad \text{or} \quad \frac{9x^2 - 4}{2x^3} \end{aligned}$$

Differentiate the following functions and simplify your answers:

$$s = \left( \frac{2t-1}{t+2} \right)^6$$

$$\frac{ds}{dt} = 6 \left[ \frac{2t-1}{t+2} \right]^5 \left[ \frac{(t+2)(2) - (2t-1)(1)}{(t+2)^6} \right]$$

$$\frac{ds}{dt} = 6 \left[ \frac{(2t-1)^5}{(t+2)^5} \right] \left[ \frac{5}{(t+2)^6} \right]$$

$$\frac{ds}{dt} = \frac{30(2t-1)^5}{(t+2)^7}$$

$$g(x) = (9x^{-3})(5x^3 - 1)^5$$

$$g'(x) = (9x^{-3})[6(5x^3 - 1)^5(15x^2)] - 27x^{-4}(5x^3 - 1)^6$$

$$g'(x) = 810x^{-1}(5x^3 - 1)^5 - 27x^{-4}(5x^3 - 1)^6$$

$$g'(x) = 27x^{-4}(5x^3 - 1)^5 \left[ \frac{30x^3 - 5x^3 + 1}{30x^3 - 5x^3 + 1} \right]$$

$$g'(x) = 27x^{-4}(5x^3 - 1)^5(25x^3 + 1)$$

$$g'(x) = \frac{27(5x^3 - 1)^5(25x^3 + 1)}{x^4}$$

**Example 1**

Let  $F(x) = f(g(x))$        $F'(x) = f'(g(x)) \cdot g'(x)$   
 If  $f(2) = 3$ ,  $f'(2) = 5$ ,  $g(1) = 2$  and  $g'(1) = 4$  find  $F'(1)$ .

$$\begin{aligned} F'(1) &= f'(\underline{g(1)}) \cdot g'(1) \\ &= \underline{f'(2)} \cdot \underline{g'(1)} \\ &= 5 \cdot 4 \\ &= 20 \end{aligned}$$

Question #6 ex. 2.4

Given      Find  $(fg)'(2) = f'(2)g(2) + f(2)g'(2)$

$$\begin{aligned} f(2) &= 3 & f'(2) &= 5 \\ f'(2) &= 5 & g(2) &= -1 \\ g(2) &= -1 & g'(2) &= -4 \\ g'(2) &= -4 & & \end{aligned}$$

$$\begin{aligned} &= (5)(-1) + (3)(-4) \\ &= -5 - 12 \\ &= -17 \end{aligned}$$

$$\begin{aligned} \text{Find } \left(\frac{f}{g}\right)'(2) &= \frac{f'(2)g(2) - f(2)g'(2)}{(g(2))^2} \\ &= \frac{(5)(-1) - (3)(-4)}{(-1)^2} \\ &= \frac{-5 + 12}{1} \\ &= 7 \end{aligned}$$

**Example 2**

If  $y = u^{10} + u^5 + 2$ , where  $u = 1 - 3x^2$ , find  $\frac{dy}{dx} \Big|_{x=1}$

$$\frac{dy}{du} = 10u^9 + 5u^4 \quad \frac{du}{dx} = -6x \quad \text{when } x=1 \\ u = 1 - 3(1)^2 = -2$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$\begin{aligned} \frac{dy}{dx} \Big|_{x=1} &= (10u^9 + 5u^4)(-6x) \\ &= (10(-2)^9 + 5(-2)^4)(-6(1)) \\ &= (-5120 + 80)(-6) \\ &= (-5040)(-6) \\ &= 30240 \end{aligned}$$

If  $y = u^{10} + u^5 + 2$ , where  $u = 1 - 3x^2$ , find  $\frac{dy}{dx} \Big|_{x=1}$

$$y = (1-3x^2)^{10} + (1-3x^2)^5 + 2$$

$$\frac{dy}{dx} = 10(1-3x^2)^9(-6x) + 5(1-3x^2)^4(-6x)$$

$$\frac{dy}{dx} \Big|_{x=1} = 10(-2)^9(-6) + 5(-2)^4(-6)$$

$$= 30720 - 480$$

$$= 30240$$

## Differentiation Rules Worksheet

$$\textcircled{3} \text{ b) } f(x) = \frac{8x^3(12x^2-5x)^8}{2-3\sqrt[3]{1-32x^{10}}} = \frac{8x^3(12x^2-5x)^8}{2-3(1-32x^{10})^{\frac{1}{3}}}$$

$$\begin{aligned} f'(x) &= \frac{[24x^2(12x^2-5x)^8 + 8x^3(8)(12x^2-5x)(24-5)] [2-3(1-32x^{10})^{\frac{1}{3}}]}{[2-3(1-32x^{10})^{\frac{1}{3}}]^2} \\ &\quad - \frac{[8x^3(12x^2-5x)^8] \left[ -\frac{3}{5}(1-32x^{10})^{-\frac{2}{5}}(-320x^9) \right]}{[2-3(1-32x^{10})^{\frac{1}{3}}]^2} \end{aligned}$$

$$\textcircled{6} \text{ b) } y = \frac{16}{\sqrt{x-1}}$$

$$y' = \frac{0(\cancel{\sqrt{x-1}}) - 16\left(\frac{1}{2}\right)(x-1)^{-\frac{1}{2}}(1)}{x-1}$$

$$y' = \frac{-8}{(x-1)^{\frac{3}{2}}} \cdot \frac{1}{(x-1)}$$

$$y' = \frac{-8}{(x-1)^{\frac{3}{2}}}$$


---

$$y = \frac{16}{\sqrt{x-1}} = 16(x-1)^{-\frac{1}{2}}$$

$$y' = -8(x-1)^{-\frac{3}{2}}(1)$$

$$y' = \frac{-8}{(x-1)^{\frac{3}{2}}}$$

# Homework

③  $y = u^2 - 2u^5$  and  $u = x - \frac{1}{x^2}$  find  $\frac{dy}{dx} \Big|_{x=4}$

$$y = \left(x - \frac{1}{x^2}\right)^2 - 2\left(x - \frac{1}{x^2}\right)^5$$

$$\frac{dy}{dx} = 2\left(x - \frac{1}{x^2}\right)\left(1 - \frac{1}{2x^3}\right) - 10\left(x - \frac{1}{x^2}\right)^4\left(1 - \frac{1}{2x^3}\right)$$

Page 103

# 3-10

$$\textcircled{3} \text{ Find } \left. \frac{dy}{dx} \right|_{x=4}$$

$$\text{if } y = u^3 - 2u^5$$

$$\frac{dy}{du} = 3u^2 - 10u^4$$

$$\text{and } u = x - \sqrt[5]{x}$$

$$\frac{du}{dx} = 1 - \frac{1}{5x^{\frac{4}{5}}}$$

$$\frac{du}{dx} = 1 - \frac{1}{2\sqrt[5]{x}}$$

$$\left. \frac{dy}{dx} \right|_{x=4} = \left[ \frac{dy}{du} \right] \left[ \frac{du}{dx} \right]$$

when  $x=4 \quad u=2$

$$= [2u - 10u^4] \left[ 1 - \frac{1}{2\sqrt[5]{x}} \right]$$

$$= [2(2) - 10(2)^4] \left[ 1 - \frac{1}{2\sqrt[5]{4}} \right]$$

$$= (-156) \left( \frac{3}{4} \right)$$

$$= \boxed{-117}$$

Page 103

# 3-10

$$\textcircled{4} \quad \left. \frac{dy}{dt} \right|_{t=1}$$

$$y = \sqrt{1+r^3}$$

$$\frac{dy}{dr} = \frac{1}{2}(1+r^3)^{-\frac{1}{2}}(3r^2)$$

$$\frac{dy}{dr} = \frac{r}{\sqrt{1+r^3}}$$

$$r = \frac{t+1}{\partial t+1}$$

$$\frac{dr}{dt} = \frac{1(\partial t+1) - \partial(t+1)}{(\partial t+1)^2}$$

$$\frac{dr}{dt} = \frac{-1}{(\partial t+1)^2}$$

$$\left. \frac{dy}{dt} \right|_{t=1} = \left[ \frac{dy}{dr} \right] \left[ \frac{dr}{dt} \right]$$

$$= \left[ \frac{r}{1+r^3} \right] \left[ \frac{-1}{(\partial t+1)^2} \right]$$

$$= \left[ \frac{\frac{2}{3}}{\sqrt{1+(\frac{2}{3})^3}} \right] \left[ \frac{-1}{(\partial(1)+1)^2} \right]$$

$$= \left[ \frac{\frac{2}{3}}{\frac{\sqrt{13}}{9}} \right] \left[ -\frac{1}{9} \right]$$

$$= \left[ \frac{\frac{2}{3} \cdot \frac{3}{\sqrt{13}}}{9} \right] \left[ -\frac{1}{9} \right]$$

$$= \boxed{\frac{-2}{9\sqrt{13}}}$$

when  $t=1 \quad r = \frac{2}{3}$