



$$q = mC\Delta T$$
$$q = mC(T_f - T_i)$$
$$(-24.0 - T_i)$$



Formation Reactions:

This reactions starts with elements only as reactants.
The reactants will form compounds as products.

elements \Rightarrow compound




The molar enthalpy symbol for a formation reaction is H_f

Simple Decomposition Reactions:

This reaction starts as a compound, which decomposes into its elements.
(opposite of a formation reaction)

compound \Rightarrow elements



The molar enthalpy symbol is H_{SD} .

Combustion Reactions:

The reaction of a substance with excess oxygen to produce an oxide.



The molar enthalpy symbol is H_c . NO_2, SO_2

(b) Molar Enthalpies

H_f - molar enthalpies of formation is the quantity of heat released or absorbed when one mole of a substance forms from its elements.


H_c - molar enthalpies of combustion is the quantity of heat released or absorbed when one mole of a substance reacts with oxygen.

H° - standard molar enthalpy is the quantity of heat released or absorbed when one mole of a substance reacts at SATP

(c) ΔH_r - Enthalpy change is the quantity of heat released or absorbed when a reaction occurs. This may also be called "Heat of Reaction" or "Change in Heat".
 $q = mC\Delta T$
 \Rightarrow must know the number of moles of a substance reacting to determine the enthalpy change

(d) Molar enthalpy may be determined from the enthalpy change as long as the number of moles (n) are known.

$$\Delta H_r = nH_r$$



$$H^\circ = -98.79 \text{ kJ/mol}$$

How do we find the change in enthalpy of $\text{SO}_{2(g)}$??

$$\Delta H_r = nH_r$$

$$\Delta H_r = (2 \text{ mol}) \left(-98.79 \frac{\text{kJ}}{\text{mol}} \right)$$


$$\Delta H_r = -197.58 \text{ kJ}$$

$$H_r = \Delta H_r / n$$

COMMUNICATING ENTHALPY CHANGES

Using ΔH_r notation:

- for chemical reactions not well known, the chemical equation must accompany the enthalpy change. The molar enthalpy of reaction (or change in enthalpy) follows the equation. **For exothermic reactions the $\Delta H_r < 0$.**




The Enthalpy Change (ΔH_r) may be included as a term in the balanced equation:

(i) In endothermic reactions - energy is reported as a reactant and is transformed in the reaction.




(ii) In exothermic reactions - energy is reported as a product since it is being produced.




POTENTIAL ENERGY DIAGRAMS

- may be used to express enthalpy change (ΔH_r)
- shows the potential energy of the reactants and products of a chemical reaction.
- shows the difference between the initial and final energies as the enthalpy change. (ΔH_r)

Endothermic Rxn



Exothermic Rxn



see Fig 11-8 p 373 (also 11-15,16,17)

For each of the following reactions:

- (a) rewrite the equation including the enthalpy change as a term
(b) draw a potential energy diagram

