

Questions from Homework

① If $h(x) = f(g(x))$, determine $g(x)$

a) $h(x) = (2x-5)^3$ and $f(x) = x^3$

$$g(x) = 2x-5$$

b) $h(x) = (5x+1)^3 - (5x+1)$ and $f(x) = x^3 - x$

$$g(x) = 5x+1$$

⑨ $j(x) = x^2$ and $k(x) = x^3$

$$k(j(x)) = k(x^2) = (x^2)^3 = x^6$$

$$j(k(x)) = j(x^3) = (x^3)^2 = x^6$$
Yes

⑩ $s(x) = x^2 + 1$ and $t(x) = x - 3$

$$s(t(x)) = s(x-3) = (x-3)^2 + 1 = \underline{\underline{x^2 - 6x + 10}}$$

$$t(s(x)) = t(x^2 + 1) = (x^2 + 1) - 3 = \underline{\underline{x^2 - 2}}$$
No

Function Operations

To combine two functions, $f(x)$ and $g(x)$, add or subtract as follows:

Sum of Functions


$$\begin{aligned} h(x) &= f(x) + g(x) \\ h(x) &= (f + g)(x) \end{aligned}$$

Difference of Functions

$$\begin{aligned} h(x) &= f(x) - g(x) \\ h(x) &= (f - g)(x) \end{aligned}$$

Key Ideas


- You can add two functions, $f(x)$ and $g(x)$, to form the combined function $h(x) = (f + g)(x)$.
- You can subtract two functions, $f(x)$ and $g(x)$, to form the combined function $h(x) = (f - g)(x)$.
- * The domain of the combined function formed by the sum or difference of two functions is the domain common to the individual functions. For example,
 - Domain of $f(x)$: $\{x \mid x \leq 3, x \in \mathbb{R}\}$
 - Domain of $g(x)$: $\{x \mid x \geq -5, x \in \mathbb{R}\}$
 - Domain of $h(x)$: $\{x \mid -5 \leq x \leq 3, x \in \mathbb{R}\}$
- The range of a combined function can be determined using its graph.
- To sketch the graph of a sum or difference of two functions given their graphs, add or subtract the y -coordinates at each point.



1. Consider the graphs of the functions $f(x)$, $g(x)$, and $h(x)$.

- a) Copy the table and use the graph of each function to complete the columns.

x	$f(x)$	$g(x)$	$h(x)$
-2			
-1			
0			
1			
2			
3			
4			



- b) What do you notice about the relationship between each value of $h(x)$ and the corresponding values of $f(x)$ and $g(x)$?

Example 1**Determine the Sum of Two Functions**

Consider the functions $f(x) = 2x + 1$ and $g(x) = x^2$.

- Determine the equation of the function $h(x) = f(x) + g(x)$.
- Sketch the graphs of $f(x)$, $g(x)$, and $h(x)$ on the same set of coordinate axes.
- State the domain and range of $h(x)$.
- Determine the values of $f(x)$, $g(x)$, and $h(x)$ when $x = 4$.

$$\begin{aligned} a) h(x) &= f(x) + g(x) \\ &= 2x + 1 + x^2 \\ &= x^2 + 2x + 1 \end{aligned}$$

$$f(x) = 2x + 1$$

$$D: \{x | x \in \mathbb{R}\}$$

$$g(x) = x^2$$

$$D: \{x | x \in \mathbb{R}\}$$


$$h(x) = x^2 + 2x + 1$$

$$D: \{x | x \in \mathbb{R}\}$$


- Method 1: Use Paper and Pencil

x	$f(x) = 2x + 1$	$g(x) = x^2$	$h(x) = x^2 + 2x + 1$
-2	-3	4	1
-1	-1	1	0
0	1	0	1
1	3	1	4
2	5	4	9

How could you use the values in the columns for $f(x)$ and $g(x)$ to determine the values in the column for $h(x)$?



How are the y -coordinates of points on the graph of $h(x)$ related to those on the graphs of $f(x)$ and $g(x)$?



How are the y -coordinates of points on the graph of $h(x)$ related to those on the graphs of $f(x)$ and $g(x)$?

- c) The function $f(x) = 2x + 1$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $g(x) = x^2$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $h(x) = (f + g)(x)$ has domain $\{x \mid x \in \mathbb{R}\}$, which consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$.
 The range of $h(x)$ is $\{y \mid y \geq 0, y \in \mathbb{R}\}$.

- d) Substitute $x = 4$ into $f(x)$, $g(x)$, and $h(x)$.

$$\begin{array}{lll} f(x) = 2x + 1 & g(x) = x^2 & h(x) = x^2 + 2x + 1 \\ f(4) = 2(4) + 1 & g(4) = 4^2 & h(4) = 4^2 + 2(4) + 1 \\ f(4) = 8 + 1 & g(4) = 16 & h(4) = 16 + 8 + 1 \\ f(4) = 9 & & h(4) = 25 \end{array}$$

Example 2**Determine the Difference of Two Functions**

Consider the functions $f(x) = \sqrt{x-1}$ and $g(x) = x - 2$.

- Determine the equation of the function $h(x) = (f - g)(x)$.
- Sketch the graphs of $f(x)$, $g(x)$, and $h(x)$ on the same set of coordinate axes.
- State the domain of $h(x)$.
- Use the graph to approximate the range of $h(x)$.

$$\begin{aligned} a) h(x) &= f(x) - g(x) \\ &= \sqrt{x-1} - (x-2) \\ &= \sqrt{x-1} - x + 2 \end{aligned}$$

$$\begin{array}{l|l|l} f(x) = \sqrt{x-1} & g(x) = x-2 & h(x) = \sqrt{x-1} - x + 2 \\ \rightarrow x-1 \geq 0 \rightarrow x \geq 1 & D: \{x | x \in \mathbb{R}\} & D: \{x \geq 1, x \in \mathbb{R}\} \\ D: \{x | x \geq 1, x \in \mathbb{R}\} & & \end{array}$$


b) Method 1: Use Paper and Pencil

For the function $f(x) = \sqrt{x-1}$, the value of the radicand must be greater than or equal to zero: $x - 1 \geq 0$ or $x \geq 1$.


x	$f(x) = \sqrt{x-1}$	$g(x) = x - 2$	$h(x) = \sqrt{x-1} - x + 2$
-2	undefined	-4	undefined
-1	undefined	-3	undefined
0	undefined	-2	undefined
1	0	-1	1
2	1	0	1
5	2	3	-1
10	3	8	-5

Why is the function $h(x)$ undefined when $x < 1$?

How could you use the values in the columns for $f(x)$ and $g(x)$ to determine the values in the column for $h(x)$?




How could you use the y-coordinates of points on the graphs of $f(x)$ and $g(x)$ to create the graph of $h(x)$?



How could you use the y-coordinates of points on the graphs of $f(x)$ and $g(x)$ to create the graph of $h(x)$?

- c) The function $f(x) = \sqrt{x-1}$ has domain $\{x \mid x \geq 1, x \in \mathbb{R}\}$.
 The function $g(x) = x-2$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $h(x) = (f-g)(x)$ has domain $\{x \mid x \geq 1, x \in \mathbb{R}\}$, which consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$.



What values of x belong to the domains of both $f(x)$ and $g(x)$?

- d) From the graph, the range of $h(x)$ appears to be approximately $\{y \mid y \leq 1.2, y \in \mathbb{R}\}$.




How can you use a graphing calculator to verify the range?

Homework



finish #1-11 on page 483-484

10.1 Sums and Differences of Functions, pages 483 to 487

1. a) $h(x) = |x - 3| + 4$ b) $h(x) = 2x - 3$
 c) $h(x) = 2x^2 + 3x + 2$ d) $h(x) = x^2 + 5x + 4$
2. a) $h(x) = 5x + 2$ b) $h(x) = -3x^2 - 4x + 9$
 c) $h(x) = -x^2 - 3x + 12$ d) $h(x) = \cos x - 4$
3. a) $h(x) = x^2 - 6x + 1$; $h(2) = -7$
 b) $m(x) = -x^2 - 6x + 1$; $m(1) = -6$
 c) $p(x) = x^2 + 6x - 1$; $p(1) = 6$
4. a) $y = 3x^2 + 2 + \sqrt{x+4}$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 b) $y = 4x - 2 - \sqrt{x+4}$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 c) $y = \sqrt{x+4} - 4x + 2$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 d) $y = 3x^2 + 4x$; domain $\{x \mid x \in \mathbb{R}\}$



6. a) 8 b) 6 c) 7
 d) not in the domain
7. a) B b) C c) A



9. a) $y = 3x^2 + 11x + 1$
c) $y = 3x^2 + 3x + 1$

b) $y = 3x^2 - 3x + 3$
d) $y = 3x^2 - 11x + 3$

10. a) $g(x) = x^2$
c) $g(x) = -3x + 1$
11. a) $g(x) = x^2 - 1$
c) $g(x) = 8x - 9$

- b) $g(x) = \sqrt{x+7}$
d) $g(x) = 3x^2 - x - 4$
- b) $g(x) = -\sqrt{x-4}$
d) $g(x) = 2x^2 - 11x - 6$