

Questions from Homework

11. If $h(x) = (f - g)(x)$ and $f(x) = 5x + 2$, determine $g(x)$.
- a) $h(x) = -x^2 + 5x + 3$
 - b) $h(x) = \sqrt{x - 4} + 5x + 2$
 - c) $h(x) = -3x + 11$
 - d) $h(x) = -2x^2 + 16x + 8$

Function Operations

To combine two functions, $f(x)$ and $g(x)$, multiply or divide as follows:

Product of Functions

$$h(x) = f(x)g(x)$$

$$h(x) = (f \cdot g)(x)$$

Quotient of Functions

$$h(x) = \frac{f(x)}{g(x)}$$



$$h(x) = \left(\frac{f}{g}\right)(x)$$

The domain of a product of functions is the domain common to the original functions. However, the domain of a quotient of functions must take into consideration that division by zero is undefined. The domain of a quotient, $h(x) = \frac{f(x)}{g(x)}$, is further restricted for values of x where $g(x) = 0$.

Example

Consider $f(x) = \sqrt{x-1}$ and $g(x) = x-2$.

The domain of $f(x)$ is $\{x \mid x \geq 1, x \in \mathbb{R}\}$, and the domain of $g(x)$ is $\{x \mid x \in \mathbb{R}\}$. So, the domain of $(f \cdot g)(x)$ is $\{x \mid x \geq 1, x \in \mathbb{R}\}$, while the domain of $\left(\frac{f}{g}\right)(x)$ is $\{x \mid x \geq 1, x \neq 2, x \in \mathbb{R}\}$



Key Ideas

- The combined function $h(x) = (f \cdot g)(x)$ represents the product of two functions, $f(x)$ and $g(x)$.
- The combined function $h(x) = \left(\frac{f}{g}\right)(x)$ represents the quotient of two functions, $f(x)$ and $g(x)$, where $g(x) \neq 0$.
- The domain of a product or quotient of functions is the domain common to both $f(x)$ and $g(x)$. The domain of the quotient $\left(\frac{f}{g}\right)(x)$ is further restricted by excluding values where $g(x) = 0$.
- The range of a combined function can be determined using its graph.

Example 1

Determine the Product of Functions

Given $f(x) = (x + 2)^2 - 5$ and $g(x) = 3x - 4$, determine $h(x) = (f \cdot g)(x)$. State the domain and range of $h(x)$.

Solution

To determine $h(x) = (f \cdot g)(x)$, multiply the two functions.

$$h(x) = (f \cdot g)(x)$$

$$h(x) = f(x)g(x)$$


$$h(x) = ((x + 2)^2 - 5)(3x - 4)$$

$$h(x) = (x^2 + 4x - 1)(3x - 4)$$

$$h(x) = 3x^3 - 4x^2 + 12x^2 - 16x - 3x + 4$$

$$h(x) = 3x^3 + 8x^2 - 19x + 4$$

cubic functions



How can you tell from the original functions that the product is a cubic function?

The function $f(x) = (x + 2)^2 - 5$ is quadratic with domain $\{x \mid x \in \mathbb{R}\}$.

The function $g(x) = 3x - 4$ is linear with domain $\{x \mid x \in \mathbb{R}\}$.

The domain of $h(x) = (f \cdot g)(x)$ consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$.

Therefore, the cubic function $h(x) = 3x^3 + 8x^2 - 19x + 4$ has domain $\{x \mid x \in \mathbb{R}\}$ and range $\{y \mid y \in \mathbb{R}\}$.

Example 2

Determine the Quotient of Functions

Consider the functions $f(x) = x^2 + x - 6$ and $g(x) = 2x + 6$.

- Determine the equation of the function $h(x) = \left(\frac{g}{f}\right)(x)$.
- Sketch the graphs of $f(x)$, $g(x)$, and $h(x)$ on the same set of coordinate axes.
- State the domain and range of $h(x)$.

Solution

- To determine $h(x) = \left(\frac{g}{f}\right)(x)$, divide the two functions.

$$h(x) = \left(\frac{g}{f}\right)(x)$$

$$h(x) = \frac{g(x)}{f(x)}$$

$$h(x) = \frac{2x + 6}{x^2 + x - 6}$$

$$h(x) = \frac{2(x + 3)}{(x + 3)(x - 2)}$$

Factor.

$$h(x) = \frac{2\cancel{(x + 3)}}{\cancel{(x + 3)}(x - 2)}$$

$$h(x) = \frac{2}{x - 2}, x \neq -3, 2 \quad \text{Identify any non-permissible values.}$$


VA \uparrow
 $x - 2 = 0$
 $x = 2$

Hole: $(x = -3)$
 $h(-3) = \frac{2}{-3 - 2} = -\frac{2}{5} \quad (-3, -\frac{2}{5})$

HA: $\lim_{x \rightarrow \infty} \frac{2x + 6}{x^2 + x - 6} = 0 \rightarrow y = 0$

b) Method 1: Use Paper and Pencil

x	$f(x) = x^2 + x - 6$	$g(x) = 2x + 6$	$h(x) = \frac{2}{x-2}, x \neq -3, 2$
-3	0	0	does not exist
-2	-4	2	$-\frac{1}{2}$
-1	-6	4	$-\frac{2}{3}$
0	-6	6	-1
1	-4	8	-2
2	0	10	undefined
3	6	12	2
4	14	14	1




How are the y-coordinates of points on the graph of $h(x)$ related to those on the graphs of $f(x)$ and $g(x)$?

- c) The function $f(x) = x^2 + x - 6$ is quadratic with domain $\{x \mid x \in \mathbb{R}\}$.
 The function $g(x) = 2x + 6$ is linear with domain $\{x \mid x \in \mathbb{R}\}$.
 The domain of $h(x) = \left(\frac{g}{f}\right)(x)$ consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$, excluding values of x where $f(x) = 0$.

Since the function $h(x)$ does not exist at $\left(-3, -\frac{2}{5}\right)$ and is undefined at $x = 2$, the domain is $\{x \mid x \neq -3, x \neq 2, x \in \mathbb{R}\}$. This is shown in the graph by the point of discontinuity at $\left(-3, -\frac{2}{5}\right)$ and the vertical asymptote that appears at $x = 2$.

How do you know there is a point of discontinuity and an asymptote?



The range of $h(x)$ is $\{y \mid y \neq 0, -\frac{2}{5}, y \in \mathbb{R}\}$.

H.A. hole

(Red arrows point from 'H.A.' to the value 0 and from 'hole' to the value $-\frac{2}{5}$ in the range set above.)



Homework

finish #1-9 on page 496-497


10.2 Products and Quotients of Functions, pages 496 to 498

1. a) $h(x) = x^2 - 49, k(x) = \frac{x+7}{x-7}, x \neq 7$
- b) $h(x) = 6x^2 + 5x - 4, k(x) = \frac{2x-1}{3x+4}, x \neq -\frac{4}{3}$
- c) $h(x) = (x+2)\sqrt{x+5}, k(x) = \frac{\sqrt{x+5}}{x+2}, x \geq -5, x \neq -2$
- d) $h(x) = \sqrt{-x^2 + 7x - 6}, k(x) = \frac{\sqrt{x-1}}{\sqrt{6-x}}, 1 \leq x < 6$

2. a) -3 b) 0 c) -1 d) 0




4. a) $h(x) = x^3 + 7x^2 + 16x + 12$




domain $\{x \mid x \in \mathbb{R}\}$, range $\{y \mid y \in \mathbb{R}\}$

- b) $h(x) = x^3 - 3x^2 - 9x + 27$




domain $\{x \mid x \in \mathbb{R}\}$, range $\{y \mid y \in \mathbb{R}\}$

- c) $h(x) = \frac{1}{x^2 + x}$




domain $\{x \mid x \neq 0, -1, x \in \mathbb{R}\}$,
range $\{y \mid y \leq -4 \text{ or } y > 0, y \in \mathbb{R}\}$

5. a) $h(x) = x + 3, x \neq -2$




domain $\{x \mid x \neq -2, x \in \mathbb{R}\}$,
range $\{y \mid y \neq 1, y \in \mathbb{R}\}$

b) $h(x) = \frac{1}{x + 3}, x \neq \pm 3$



domain $\{x \mid x \neq \pm 3, x \in \mathbb{R}\}$,
range $\{y \mid y \neq 0, \frac{1}{6}, y \in \mathbb{R}\}$

c) $h(x) = \frac{x}{x + 1}, x \neq -1, 0$




domain $\{x \mid x \neq -1, 0, x \in \mathbb{R}\}$,
range $\{y \mid y \neq 0, 1, y \in \mathbb{R}\}$


6. a) $y = x^3 + 3x^2 - 10x - 24$


b) $y = \frac{x^2 - x - 6}{x + 4}, x \neq -4$ c) $y = \frac{2x - 1}{x + 4}, x \neq -4$

d) $y = \frac{x^2 - x - 6}{x^2 + 8x + 16}, x \neq -4$

7. a) $g(x) = 3$ b) $g(x) = -x$
 c) $g(x) = \sqrt{x}$ d) $g(x) = 5x - 6$
 8. a) $g(x) = x + 7$ b) $g(x) = \sqrt{x + 6}$
 c) $g(x) = 2$ d) $g(x) = 3x^2 + 26x - 9$

9. a)  $f(x)$:
 domain $\{x \mid x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$
 $g(x)$: domain $\{x \mid x \in \mathbb{R}\}$,
 range $\{y \mid -1 \leq y \leq 1, y \in \mathbb{R}\}$

b)  $y = (2x + 5)\cos(x)$
 domain $\{x \mid x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$



11. a) $y = \frac{f(x)}{g(x)}$ b) $y = f(x)f(x)$
 c) The graphs of $y = \frac{\sin x}{\cos x}$ and $y = \tan x$ appear to be the same. The graphs of $y = 1 - \cos^2 x$ and $y = \sin^2 x$ appear to be the same.