


Exponential Functions

The graph of an **exponential function**, such as $y = c^x$, is increasing for $c > 1$, decreasing for $0 < c < 1$, and neither increasing nor decreasing for $c = 1$. From the graph, you can determine characteristics such as domain and range, any intercepts, and any asymptotes.



exponential function

- a function of the form $y = c^x$, where c is a constant ($c > 0$) and x is a variable



Why is the definition of an exponential function restricted to positive values of c ?

Did You Know?

Any letter can be used to represent the base in an exponential function. Some other common forms are $y = a^x$ and $y = b^x$. In this chapter, you will use the letter c . This is to avoid any confusion with the transformation parameters, a , b , h , and k , that you will apply in Section 7.2.

Key Ideas

- An exponential function of the form $y = c^x, c > 0,$
 - is increasing for $c > 1$
 - is decreasing for $0 < c < 1$
 - is neither increasing nor decreasing for $c = 1$
- * has a domain of $\{x \mid x \in \mathbb{R}\}$
- * has a range of $\{y \mid y > 0, y \in \mathbb{R}\}$
- has a y-intercept of 1 $(0,1)$
- has no x-intercept
- has a horizontal asymptote at $y = 0$



Example 1

Analyse the Graph of an Exponential Function

Graph each exponential function. Then identify the following:

- the domain and range
- the x -intercept and y -intercept, if they exist
- whether the graph represents an increasing or a decreasing function
- the equation of the horizontal asymptote

a) $y = 4^x$

b) $f(x) = \left(\frac{1}{2}\right)^x$

Solution

a) Method 1: Use Paper and Pencil


Use a table of values to graph the function.

Select integral values of x that make it easy to calculate the corresponding values of y for $y = 4^x$.

$$y = 4^x$$

x	y
-2	$\frac{1}{16}$
-1	$\frac{1}{4}$
0	1
1	4
2	16

- $D: \{x | x \in \mathbb{R}\}$
- $R: \{y | y > 0, y \in \mathbb{R}\}$
- x-int: none
- y-int: $(0, 1)$
- HA: $y = 0$




b) Method 1: Use Paper and Pencil

Use a table of values to graph the function.

Select integral values of x that make it easy to calculate the corresponding values of y for $f(x) = \left(\frac{1}{2}\right)^x$.

x	$f(x)$
-3	8
-2	4
-1	2
0	1
1	$\frac{1}{2}$
2	$\frac{1}{4}$


- $D: \{x | x \in \mathbb{R}\}$
- $R: \{y | y > 0, y \in \mathbb{R}\}$
- x -int: none
- y -int: $(0, 1)$
- HA: $y = 0$



Example 2

Write the Exponential Function Given Its Graph

What function of the form $y = c^x$ can be used to describe the graph shown?



decreasing:
base is $0 < c < 1$
Find c

Solution

Look for a pattern in the ordered pairs from the graph.

x	y
-2	16
-1	4
0	1

$> \frac{1}{4}$
 $> \frac{1}{4}$

As x values increase by 1
the values decrease by a
factor of $\frac{1}{4}$

$$c = \frac{1}{4}$$

Choose a point other than (0, 1) to substitute into the function $y = \left(\frac{1}{4}\right)^x$ to verify that the function is correct. Try the point $(-2, 16)$.

Why should you not use the point (0, 1) to verify that the function is correct?

Check:

Left Side

Right Side

$$y = \left(\frac{1}{4}\right)^x$$

$$16 \quad \left(\frac{1}{4}\right)^{-2}$$


$$(4)^2$$

$$16$$

Example 3

Application of an Exponential Function

A radioactive sample of radium (Ra-225) has a **half-life** of 15 days. The mass, m , in grams, of Ra-225 remaining over time, t , in 15-day intervals, can be modelled using the exponential graph shown.



- What is the initial mass of Ra-225 in the sample? What value does the mass of Ra-225 remaining approach as time passes? *1.0g*
- What are the domain and range of this function?
- Write the exponential decay model that relates the mass of Ra-225 remaining to time, in 15-day intervals.
- Estimate how many days it would take for Ra-225 to decay to $\frac{1}{30}$ of its original mass.

b) D: $\{x | x \geq 0, x \in \mathbb{R}\}$
 R: $\{y | 0 < y \leq 1, y \in \mathbb{R}\}$

c) $y = (1) \left(\frac{1}{2}\right)^{\frac{x}{15}}$

Initial Amount (points to 1)
Base ($\frac{1}{2}$) half life (points to $\frac{1}{2}$)
Time it takes to $\frac{1}{2}$ (points to $\frac{x}{15}$)

d) $\frac{1}{30} = 1 \left(\frac{1}{2}\right)^{\frac{x}{15}}$

$\frac{1}{30} = \left(\frac{1}{2}\right)^{\frac{x}{15}}$ $\star \frac{\log(\frac{1}{30})}{\log(\frac{1}{2})}$

$\left(\frac{1}{2}\right)^{4.91} = \left(\frac{1}{2}\right)^{\frac{x}{15}}$

15 \cdot 4.91 = $\frac{x}{15}$ \cdot 15

73.6 = x

Homework

#1-8 on page 343

$${}^4C_1 = 4$$

$${}^4C_2 = 6$$

$${}^4C_3 = 4$$





$${}^4C_4 = 1$$

1, 5, 10, 25, 6, 11, 26, 15, 30, 35


16, 31, 40, 36, 41

7.1 Characteristics of Exponential Functions, pages 342 to 345

1. a) No, the variable is not the exponent.
 b) Yes, the base is greater than 0 and the variable is the exponent.
 c) No, the variable is not the exponent.
 d) Yes, the base is greater than 0 and the variable is the exponent.
2. a) $f(x) = 4^x$ b) $g(x) = \left(\frac{1}{4}\right)^x$
 c) $x = 0$, which is the y -intercept
3. a) B b) C c) A
4. a) $f(x) = 3^x$ b) $f(x) = \left(\frac{1}{5}\right)^x$




- 6. a) $c > 1$; number of bacteria increases over time
- b) $0 < c < 1$; amount of actinium-225 decreases over time
- c) $0 < c < 1$; amount of light decreases with depth
- d) $c > 1$; number of insects increases over time




- b) i) 1 person ii) 2 people
- iii) 16 people iv) 1024 people

- 8. a) If the population increases by 10% each year, the population becomes 110% of the previous year's population. So, the growth rate is 110% or 1.1 written as a decimal.



domain $\{t \mid t \geq 0, t \in \mathbb{R}\}$ and range $\{P \mid P \geq 100, P \in \mathbb{R}\}$

- c) The base of the exponent would become $100\% - 5\%$ or 95%, written as 0.95 in decimal form.



domain $\{t \mid t \geq 0, t \in \mathbb{R}\}$ and range $\{P \mid 0 < P \leq 100, P \in \mathbb{R}\}$