

Understanding Logarithms

Focus on...

- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- determining the characteristics of the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

Questions from Homework

$$\textcircled{3} \quad \text{b) } \log_2 32 = 5 \quad \log_3 \left(\frac{1}{27}\right) = -3$$

$$\frac{\log 32}{\log 2} = 5 \quad \frac{\log \left(\frac{1}{27}\right)}{\log 3} = -3$$

$$\text{j) } \log_9 \sqrt{3} = 0.25$$

$$\frac{\log(\sqrt{3})}{\log 9} = 0.25 \text{ or } \frac{1}{4}$$

$$\textcircled{4} \quad \text{f) } 3^{2x-1} = 5 \quad (\text{exp form})$$

\uparrow base
 \uparrow ans

$$\log_3(5) = 2x - 1$$

$$\frac{\log_3(5) + 1}{2} = \frac{2x}{2}$$

$$\frac{\log_3(5) + 1}{2} = x$$

$$\frac{1}{2}(\log_3(5) + 1) = x$$

$$\text{g) } \log_2(\log_3 x) = 4 \quad (\text{log form})$$

\uparrow Base
 \uparrow ans
 \uparrow exp

$$2^4 = \log_3 x \quad (\text{Exp. form})$$

$$16 = \log_3 x \quad (\text{log form})$$

\uparrow exp
 \uparrow Base
 \uparrow ans

$$3^{16} = x \quad (\text{Exp. form})$$

$$43\,046\,721 = x$$

$$\text{h) } 10^{5^x} = 3 \quad (\text{Exp. form})$$

\uparrow Base
 \uparrow ans

$$\log_{10}(3) = 5^x \quad (\text{Exp form})$$

\uparrow ans
 \uparrow Base

$$\log_5(\log_{10} 3) = x$$

General Properties of Logarithms:

If $c > 0$ and $c \neq 1$, then...

$$(i) \log_c 1 = 0$$

$$(ii) \log_c c^x = x$$

$$(iii) c^{\log_c x} = x$$

Did You Know?

The input value for a logarithm is called an argument. For example, in the expression $\log_6 1$, the argument is 1.

$$(i) \log_5 1 = 0 \quad (ii) \log_2 2^3 = 3 \quad (iii) 7^{\log_7 49} = 49$$

$$5^{\log_5 10} = 10$$

Product Law of Logarithms

The logarithm of a product of numbers can be expressed as the sum of the logarithms of the numbers.

$$\log_c MN = \log_c M + \log_c N$$

$$\begin{aligned} \log_2 16 + \log_2 2 &= \log_2 (16 \cdot 2) \\ &= \log_2 32 \\ &= 5 \end{aligned}$$

Proof

Let $\log_c M = x$ and $\log_c N = y$, where M , N , and c are positive real numbers with $c \neq 1$.

Write the equations in exponential form as $M = c^x$ and $N = c^y$:

$$MN = (c^x)(c^y)$$

$$MN = c^{x+y}$$

$$\log_c MN = x + y$$

$$\log_c MN = \log_c M + \log_c N$$

Apply the product law of powers.

Write in logarithmic form.

Substitute for x and y .

Quotient Law of Logarithms

The logarithm of a quotient of numbers can be expressed as the difference of the logarithms of the dividend and the divisor.

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

ex: $\log_2 80 - \log_2 5 = \log_2 \left(\frac{80}{5}\right)$
 $= \log_2 16$
 $= 4$

Proof

Let $\log_c M = x$ and $\log_c N = y$, where M , N , and c are positive real numbers with $c \neq 1$.

Write the equations in exponential form as $M = c^x$ and $N = c^y$:

$$\frac{M}{N} = \frac{c^x}{c^y}$$

$$\frac{M}{N} = c^{x-y}$$

$$\log_c \frac{M}{N} = x - y$$

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

Apply the quotient law of powers.

Write in logarithmic form.

Substitute for x and y .

Power Law of Logarithms

The logarithm of a power of a number can be expressed as the exponent times the logarithm of the number.

$$\log_c M^P = P \log_c M$$

How could you prove the quotient law using the product law and the power law?

Proof

Let $\log_c M = x$, where M and c are positive real numbers with $c \neq 1$.

Write the equation in exponential form as $M = c^x$.

Let P be a real number.

$$M = c^x$$

$$M^P = (c^x)^P$$

$$M^P = c^{xP}$$

Simplify the exponents.

$$\log_c M^P = xP$$

Write in logarithmic form.

$$\log_c M^P = (\log_c M)P$$

Substitute for x .

$$\log_c M^P = P \log_c M$$

The laws of logarithms can be applied to logarithmic functions, expressions, and equations.

$$\begin{aligned} \text{ex: } & \log_3 \sqrt[4]{27} \\ &= \log_3 (27)^{\frac{1}{4}} \\ &= \frac{1}{4} \log_3 27 \\ &= \frac{1}{4} \frac{(3)}{1} \\ &= \frac{3}{4} \end{aligned}$$

Product Law of Logarithms

The logarithm of a product of numbers can be expressed as the sum of the logarithms of the numbers.

$$\log_c MN = \log_c M + \log_c N$$

Quotient Law of Logarithms

The logarithm of a quotient of numbers can be expressed as the difference of the logarithms of the dividend and the divisor.

$$\log_c \frac{M}{N} = \log_c M - \log_c N$$

Power Law of Logarithms

The logarithm of a power of a number can be expressed as the exponent times the logarithm of the number.

$$\log_c M^P = P \log_c M$$

How could you prove the quotient law using the product law and the power law?

Example 1

Use the Laws of Logarithms to Expand Expressions

Write each expression in terms of individual logarithms of x , y , and z .

a) $\log_5 \frac{xy}{z}$

b) $\log_7 \sqrt[3]{x}$

c) $\log_6 \frac{1}{x^2}$

d) $\log \frac{x^3}{y\sqrt{z}}$

$$\text{a) } \log_5 \frac{xy}{z} = \log_5 x + \log_5 y - \log_5 z$$

$$\text{b) } \log_7 \sqrt[3]{x} = \log_7 x^{\frac{1}{3}} = \frac{1}{3} \log_7 x$$

$$\begin{aligned} \text{c) } \log_6 \frac{1}{x^2} &= \log_6 1 - \log_6 x^2 \\ &= 0 - 2 \log_6 x \\ &= -2 \log_6 x \end{aligned}$$

$$\begin{aligned} \text{d) } \log \left(\frac{x^3}{y\sqrt{z}} \right) \\ &= \log \left(\frac{x^3}{y z^{\frac{1}{2}}} \right) \\ &= \log x^3 - \log y - \log z^{\frac{1}{2}} \\ &= 3 \log x - \log y - \frac{1}{2} \log z \end{aligned}$$

Example 2

Use the Laws of Logarithms to Evaluate Expressions

Use the laws of logarithms to simplify and evaluate each expression.

a) $\log_6 8 + \log_6 9 - \log_6 2$

b) $\log_7 7\sqrt{7}$

c) $2 \log_2 12 - (\log_2 6 + \frac{1}{3} \log_2 27)$

$$a) \log_6 8 + \log_6 9 - \log_6 2 = \log_6 \left(\frac{8 \cdot 9}{2} \right) = \log_6 (36) = 2$$

$$b) \log_7 7\sqrt{7} = \log_7 7(7)^{\frac{1}{2}} = \log_7 7^{\frac{3}{2}} = \frac{3}{2}$$

$$\hookrightarrow \frac{3}{2} (\log_7 7)$$

$$\frac{3}{2} (1)$$

$$\frac{3}{2}$$

$$c) 2 \log_2 12 - (\log_2 6 + \frac{1}{3} \log_2 27)$$

$$\log_2 12^2 - (\log_2 6 + \log_2 27^{\frac{1}{3}})$$

$$= \log_2 144 - (\log_2 6 + \log_2 3)$$

$$= \log_2 144 - \log_2 6 - \log_2 3$$

$$= \log_2 \left(\frac{144}{6 \cdot 3} \right)$$

$$= \log_2 8$$

$$= 3$$

Example 3

Use the Laws of Logarithms to Simplify Expressions

Write each expression as a single logarithm in simplest form. State the restrictions on the variable.

a) $\log_7 x^2 + \log_7 x - \frac{5 \log_7 x}{2}$

b) $\log_5 (2x - 2) - \log_5 (x^2 + 2x - 3)$

a) $\log_7 x^2 + \log_7 x - \frac{5}{2} \log_7 x$

$\log_7 x^2 + \log_7 x - \log_7 x^{5/2}$

$\log_7 \left(\frac{x^2 \cdot x}{x^{5/2}} \right) \rightarrow 3 - \frac{5}{2}$

$\log_7 x^{1/2} \leftarrow \frac{6}{2} - \frac{5}{2}$

b) $\log_5 (2x - 2) - \log_5 (x^2 + 2x - 3)$

$\log_5 \left(\frac{2x - 2}{x^2 + 2x - 3} \right)$

$\log_5 \left(\frac{2(x-1)}{(x-1)(x+3)} \right)$

$\log_5 \left(\frac{2}{x+3} \right)$

For the original expression to be defined, both logarithmic terms must be defined.

$$2x - 2 > 0$$

$$2x > 2$$

$$x > 1$$

$$x^2 + 2x - 3 > 0$$

$$(x + 3)(x - 1) > 0$$

$$x < -3 \text{ or } x > 1$$

What other methods could you have used to solve this quadratic inequality?

The conditions $x > 1$ and $x < -3$ or $x > 1$ are both satisfied when $x > 1$.

Hence, the variable x needs to be restricted to $x > 1$ for the original expression to be defined and then written as a single logarithm.

Therefore, $\log_5 (2x - 2) - \log_5 (x^2 + 2x - 3) = \log_5 \frac{2}{x+3}$, $x > 1$.

Key Ideas

- Let P be any real number, and M , N , and c be positive real numbers with $c \neq 1$. Then, the following laws of logarithms are valid.

Name	Law	Description
Product	$\log_c MN = \log_c M + \log_c N$	The logarithm of a product of numbers is the sum of the logarithms of the numbers.
Quotient	$\log_c \frac{M}{N} = \log_c M - \log_c N$	The logarithm of a quotient of numbers is the difference of the logarithms of the dividend and divisor.
Power	$\log_c M^P = P \log_c M$	The logarithm of a power of a number is the exponent times the logarithm of the number.

- Many quantities in science are measured using a logarithmic scale. Two commonly used logarithmic scales are the decibel scale and the pH scale.

Homework

Exercise 3

Do I really understand??...

a) Express the following as a single logarithm... $2\log_2 3^2 + \log_2 6 - 3\log_2 3$

b) Evaluate the following... $\log_2 (32)^{\frac{1}{3}}$

c) Express the following as a single logarithm... $\frac{1}{2}[(\log_5 a + 2\log_5 b) - 3\log_5 c]$

d) Express as a single logarithm in simplest form...

$$\frac{3}{4} \left[12(\log_8 x^2 - 2\log_8 x) + 8\log_8 \sqrt{x} - 4\log_8 \frac{1}{x^7} \right]$$