


Understanding Logarithms

Focus on...

- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- determining the characteristics of the graph of $y = \log_c x$, $c > 0$, $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

For the exponential function $y = c^x$, the inverse is $x = c^y$. This inverse is also a function and is called a **logarithmic function**. It is written as $y = \log_c x$, where c is a positive number other than 1.

Logarithmic Form Exponential Form



Since our number system is based on powers of 10, **logarithms** with base 10 are widely used and are called **common logarithms**. When you write a common logarithm, you do not need to write the base. For example, $\log 3$ means $\log_{10} 3$.

$$\text{or } \log_{10} 150 = \log 150$$

logarithmic function

- a function of the form $y = \log_c x$, where $c > 0$ and $c \neq 1$, that is the inverse of the exponential function $y = c^x$

logarithm

- an exponent
- in $x = c^y$, y is called the logarithm to base c of x

common logarithm

- a logarithm with base 10

Write each of the following in logarithmic form

a) $32 = 2^5$
 (Handwritten: "ans" points to 5, "exp" points to 2, "Base" points to 2)
 $\log_2(32) = 5$

b) $2^{-5} = \frac{1}{32}$
 $\log_2\left(\frac{1}{32}\right) = -5$

c) $x = 10^y$
 $\log_{10}(x) = y$
 $\log x = y$

Logarithmic Form Exponential Form



Write each of the following in exponential form

a) $\log_4 16 = 2$
 (Handwritten: "ans" points to 2, "exp" points to 16, "Base" points to 4)
 $4^2 = 16$

b) $\log_2\left(\frac{1}{32}\right) = -5$
 $2^{-5} = \frac{1}{32}$

c) $\log 65 = 1.8129$
 $10^{1.8129} = 65$

Example 1

Evaluating a Logarithm

Evaluate.

a) $\log_7 49 = ?$

$$\frac{\log 49}{\log 7} = ?$$

$$x = \log_7 49$$

$$7^x = 49$$

~~$$7^x = (7)^2$$~~

$$x = 2$$

$$\boxed{\log_7 49 = 2}$$

b) $\log_6 1 = ?$

$$\frac{\log 1}{\log 6} = ?$$

$$x = \log_6 1$$

$$6^x = 1$$

~~$$6^x = (6)^0$$~~

$$x = 0$$

$$\boxed{\log_6 1 = 0}$$

c) $\log 0.001 = ?$

$$\frac{\log 0.001}{\log 10} = ?$$

$$x = \log 0.001$$

$$10^x = 0.001$$

~~$$10^x = (10)^{-3}$$~~

$$x = -3$$

$$\boxed{\log 0.001 = -3}$$

d) $\log_2 \sqrt{8} = ?$

$$\frac{\log \sqrt{8}}{\log 2} = ?$$

$$x = \log_2 \sqrt{8}$$

$$2^x = \sqrt{8}$$

$$2^x = (8)^{1/2}$$

$$2^x = (2^3)^{1/2}$$

~~$$2^x = 2^{3/2}$$~~

$$x = \frac{3}{2}$$

$$\boxed{\log_2 \sqrt{8} = \frac{3}{2}}$$

Example 2

Determine an Unknown in an Expression in Logarithmic Form

Determine the value of x . (convert to exponential form)

a) $\log_5 x = -3$

b) $\log_x 36 = 2$

c) $\log_{64} x = \frac{2}{3}$

a) $\log_5 x = -3$ (log. form)

$$5^{-3} = x \text{ (exp. form)}$$

$$\left(\frac{1}{5}\right)^3 = x$$

$$\boxed{\frac{1}{125} = x}$$

b) $\log_x 36 = 2$ (log. form)

$$x^2 = 36 \text{ (exp. form)}$$

$$x = \pm 6$$

$$\boxed{x = 6} \quad c > 0$$

c) $\log_{64} x = \frac{2}{3}$ (log. form)

$$64^{\frac{2}{3}} = x \text{ (exp. form)}$$

$$16 = x$$

$$2^4 = 16 \quad (\text{exponential form})$$

↑ Base ↑ ans.

← exp.

$$\log_2(16) = 4 \quad (\text{logarithmic form})$$

Chapter 7	Inverse	Chapter 8
Exponential		Logarithmic
$y = c^x, c > 0, c \neq 1$		$y = \log_c x, c > 0, c \neq 1$
D: $\{x x \in \mathbb{R}\}$		D: $\{x x > 0, x \in \mathbb{R}\}$ or $(0, \infty)$
R: $\{y y > 0, y \in \mathbb{R}\}$		R: $\{y y \in \mathbb{R}\}$ or $(-\infty, \infty)$
x-int: none		x-int: $x = 1$ (1, 0)
y-int: $y = 1$ or (0, 1)		y-int: none
HA: $y = 0$		VA: $x = 0$

Example 3

Graph the Inverse of an Exponential Function

- a) State the inverse of $f(x) = 3^x$.
- b) Sketch the graph of the inverse. Identify the following characteristics of the inverse graph: $(x, y) \rightarrow (y, x)$
- the domain and range
 - the x-intercept, if it exists
 - the y-intercept, if it exists
 - the equations of any asymptotes

$$a) f(x) = 3^x$$

$$y = 3^x$$

$$x = 3^y \quad (\text{exp. form})$$

$$\log_3 x = y \quad (\text{log form})$$

$$y = \log_3 x$$

$$f^{-1}(x) = \log_3 x$$

$y = 3^x$ passes the HLT

$$b) y = 3^x \quad (x, y) \rightarrow (y, x)$$

x	y
-2	1/9
-1	1/3
0	1
1	3
2	9

Inverse

$$y = \log_3 x$$

x	y
1/9	-2
1/3	-1
1	0
3	1
9	2

Solution


a) The inverse of $y = f(x) = 3^x$ is $x = 3^y$ or, expressed in logarithmic form, $y = \log_3 x$. Since the inverse is a function, it can be written in function notation as

How do you know that $y = \log_3 x$ is a function?

b) Set up tables of values for both the exponential function, $f(x)$, and its inverse, $f^{-1}(x)$. Plot the points and join them with a smooth curve.

$y = 3^x$ $f(x) = 3^x$	
x	y
-3	$\frac{1}{27}$
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9
3	27

$y = \log_3 x$ $f^{-1}(x) = \log_3 x$	
x	y
$\frac{1}{27}$	-3
$\frac{1}{9}$	-2
$\frac{1}{3}$	-1
1	0
3	1
9	2
27	3



The graph of the inverse, $f^{-1}(x) = \log_3 x$, is a reflection of the graph

of $f(x) = 3^x$ about the line $y = x$. For $f^{-1}(x) = \log_3 x$,

- the domain is $\{x \mid x > 0, x \in \mathbb{R}\}$ and the range is $\{y \mid y \in \mathbb{R}\}$ or $(-\infty, \infty)$
- the x-intercept is 1 or $(1, 0)$
- there is no y-intercept
- the vertical asymptote, the y-axis, has equation $x = 0$; there is no horizontal asymptote

How do the characteristics of $f^{-1}(x) = \log_3 x$ compare to the characteristics of $f(x) = 3^x$?

Key Ideas


- A logarithm is an exponent.
- Equations in exponential form can be written in logarithmic form and vice versa.

Exponential Form **Logarithmic Form**

$$x = c^y \qquad y = \log_c x$$

- The inverse of the exponential function $y = c^x$, $c > 0$, $c \neq 1$, is $x = c^y$ or, in logarithmic form, $y = \log_c x$. Conversely, the inverse of the logarithmic function $y = \log_c x$, $c > 0$, $c \neq 1$, is $x = \log_c y$ or, in exponential form, $y = c^x$.
- The graphs of an exponential function and its inverse logarithmic function are reflections of each other in the line $y = x$, as shown.
- For the logarithmic function $y = \log_c x$, $c > 0$, $c \neq 1$,
 - the domain is $\{x \mid x > 0, x \in \mathbb{R}\}$
 - the range is $\{y \mid y \in \mathbb{R}\}$
 - the x-intercept is 1
 - the vertical asymptote is $x = 0$, or the y-axis
- A common logarithm has base 10. It is not necessary to write the base for common logarithms:

$$\log_{10} x = \log x$$



Questions from Homework

$$\textcircled{1} \text{ a) } y = 2^x$$

$$x = 2^y$$

$$\log_2(x) = y$$

$$y = \log_2 x$$

$$D: \{x \mid x > 0, x \in \mathbb{R}\}$$

$$R: \{y \mid y \in \mathbb{R}\}$$

$$x\text{-int: } x = 1$$

$$y\text{-int: none}$$

$$VA: x = 0$$

$$\text{b) } y = \left(\frac{1}{3}\right)^x$$

$$x = \left(\frac{1}{3}\right)^y$$

$$\log_{\frac{1}{3}}(x) = y$$

$$y = \log_{\frac{1}{3}} x$$

$$D: \{x \mid x > 0, x \in \mathbb{R}\}$$

$$R: \{y \mid y \in \mathbb{R}\}$$

$$x\text{-int: } x = 1$$

$$y\text{-int: none}$$

$$VA: x = 0$$

17. The growth of a new social networking site can be modelled by the exponential function $N(t) = 1.1^t$, where N is the number of users after t days.

- a) Write the equation of the inverse.
- b) How long will it take, to the nearest day, for the number of users to exceed 1 000 000?

$$\textcircled{2} \text{ d) } 7^{\overset{\text{exp}}{2x}} = \underbrace{y+3}_{\text{ans}} \rightarrow \log_7(y+3) = 2x$$

↑ Base

$$\textcircled{3} \text{ c) } \log_{10}(1000000) = 6 \rightarrow 10^6 = 1000000$$

↑ Base
 ↑ ans
 ↑ exp.

$$\textcircled{4} \text{ c) } \log_4 \sqrt[3]{4} = \frac{1}{3}$$

$$\frac{\log(4^{1/3})}{\log 4} = \frac{1}{3}$$

$$x = \log_4(4)^{1/3} \text{ (log form)}$$

~~$$4^x = 4^{1/3} \text{ (exp. form)}$$~~

$$x = \frac{1}{3}$$

$$\textcircled{5} \quad a < \log_p 28 < b$$

↑

$$4 < \log_p 28 < 5$$

$2^4 = 16$
 $2^5 = 32$
 $\log_p 28 = 4.8$

$$\frac{\log 28}{\log 2} = 4.8$$


$$\textcircled{1a} \text{ c) } \log_{1/4} x = -3 \rightarrow \left(\frac{1}{4}\right)^{-3} = x$$

↑ Base
 ↑ ans
 ↑ exp


$$4^3 = x$$

$$\boxed{64 = x}$$

8.1 Understanding Logarithms, pages 380 to 382




ii) $y = \log_2 x$
 iii) domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1, no y-intercept,
 vertical asymptote $x = 0$



ii) $y = \log_3 x$
 iii) domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1, no y-intercept,
 vertical asymptote $x = 0$

2. a) $\log_{12} 144 = 2$ b) $\log_8 2 = \frac{1}{3}$
 c) $\log_{10} 0.000\ 01 = -5$ d) $\log_7 (y + 3) = 2x$
3. a) $5^2 = 25$ b) $8^{\frac{2}{3}} = 4$
 c) $10^6 = 1\ 000\ 000$ d) $11^y = x + 3$
4. a) 3 b) 0 c) $\frac{1}{3}$ d) -3
5. $a = 4; b = 5$



domain $\{x \mid x > 0, x \in \mathbb{R}\}$,
 range $\{y \mid y \in \mathbb{R}\}$,
 x-intercept 1,
 no y-intercept,
 vertical asymptote $x = 0$

10. They are reflections of each other in the line $y = x$.
11. a) They have the exact same shape.
 b) One of them is increasing and the other is decreasing.
12. a) 216 b) 81 c) 64 d) 8
13. a) 7 b) 6
14. a) 0 b) 1
15. -1
16. 16
17. a) $t = \log_{0.11} N$ b) 145 days
18. The larger asteroid had a relative risk that was 1479 times as dangerous.
19. 1000 times as great
20. 5
21. $m = 14, n = 13$
22. $4n$
23. $y = 3^{2^x}$

Transformations of Logarithmic Functions

$$y = a \log_c (b(x-h)) + k$$

Focus on...

- explaining the effects of the parameters a , b , h , and k in $y = a \log_c (b(x-h)) + k$ on the graph of $y = \log_c x$, where $c > 1$
- sketching the graph of a logarithmic function by applying a set of transformations to the graph of $y = \log_c x$, where $c > 1$, and stating the characteristics of the graph

Remember:

Parameter	Transformation
a	$(x, y) \rightarrow (x, ay)$
b	$(x, y) \rightarrow \left(\frac{x}{b}, y\right)$
h	$(x, y) \rightarrow (x + h, y)$
k	$(x, y) \rightarrow (x, y + k)$

Example 1

Translations of a Logarithmic Function

- a) Use transformations to sketch the graph of the function
 $y = \log_3(x + 9) + 2$.
- b) Identify the following characteristics of the graph of the function.
 - i) the equation of the asymptote
 - ii) the domain and range
 - iii) the y-intercept, if it exists
 - iv) the x-intercept, if it exists

a) $y = 1 \log_3(x+9) + 2$ $c = 3 \rightarrow$ base

$a = 1 \rightarrow$ no vertical stretch or reflection

$b = 1 \rightarrow$ no horizontal stretch or reflection

$h = -9 \rightarrow$ 9 units left

$k = 2 \rightarrow$ 2 units up

$(x, y) \rightarrow \left(\frac{1}{1}x + (-9), 1y + 2\right)$

$(x, y) \rightarrow (x - 9, y + 2)$

$y = 3^x$


x	y
-2	1/9
-1	1/3
0	1
1	3
2	9

$y = \log_3 x$

x	y
1/9	-2
1/3	-1
1	0
3	1
9	2

$(x, y) \rightarrow (x - 9, y + 2)$

x	y
-8.8 or -80/9	0
-8.6 or -26/3	1
-8	2
-6	3
0	4



$$b) \text{ VA: } x = -9$$

$$i) \text{ D: } \{x \mid x > -9, x \in \mathbb{R}\} \text{ or } (-9, \infty)$$

$$\text{R: } \{y \mid y \in \mathbb{R}\}$$

$$iii) \text{ y int (x=0)}$$

$$y = \log_3(x+9) + 2$$

$$y = \log_3(0+9) + 2$$

$$y = \log_3(9) + 2$$

$$y = 2 + 2$$

$$y = 4$$

$$(0, 4)$$

$$\frac{\log 9}{\log 3} = 2$$

$$iv) \text{ x int (y=0)}$$

$$y = \log_3(x+9) + 2$$

$$0 = \log_3(x+9) + 2$$

$$-2 = \log_3(x+9) \text{ (log form)}$$

$$\begin{array}{ccc} \uparrow & \uparrow & \uparrow \\ \text{exp} & \text{Base} & \text{ans} \end{array}$$

$$3^{-2} = x+9$$

$$\left(\frac{1}{3}\right)^2 = x+9$$

$$\frac{1}{9} = x+9$$

$$\frac{1}{9} - \frac{9}{9} = x$$

$$\frac{1}{9} - \frac{81}{9} = x$$

$$-\frac{80}{9} = x$$

$$(-8.\bar{8}, 0)$$


Example 2

Reflections, Stretches, and Translations of a Logarithmic Function

- a) Use transformations to sketch the graph of the function
 $y = -\log_2(2x + 6)$. $\rightarrow y = -\log_2(2(x+3))$
- b) Identify the following characteristics of the graph of the function.
- the equation of the asymptote
 - the domain and range
 - the y -intercept, if it exists
 - the x -intercept, if it exists

Key Ideas

- To represent real-life situations, you may need to transform the basic logarithmic function $y = \log_b x$ by applying reflections, stretches, and translations. These transformations should be performed in the same manner as those applied to any other function.
- The effects of the parameters a , b , h , and k in $y = a \log_c (b(x - h)) + k$ on the graph of the logarithmic function $y = \log_c x$ are shown below.



- Only parameter h changes the vertical asymptote and the domain. None of the parameters change the range.

Homework

Questions #1, 2, 4, 5, 8, 11 on
page 389 - 391