

# Understanding Logarithms

## Focus on...

---

- demonstrating that a logarithmic function is the inverse of an exponential function
- sketching the graph of  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$
- determining the characteristics of the graph of  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$
- explaining the relationship between logarithms and exponents
- expressing a logarithmic function as an exponential function and vice versa
- evaluating logarithms using a variety of methods

**exponential**

For the exponential function  $y = c^x$ , the inverse is  $x = c^y$ . This inverse is also a function and is called a **logarithmic function**. It is written as  $y = \log_c x$ , where  $c$  is a positive number other than 1.

**logarithmic****Logarithmic Form****Exponential Form**

Since our number system is based on powers of 10, **logarithms** with base 10 are widely used and are called **common logarithms**. When you write a common logarithm, you do not need to write the base.

For example,  $\log 3$  means  $\log_{10} 3$ . or  $\log_{10} 150 = \log 150$

**logarithmic  
function**

- a function of the form  $y = \log_c x$ , where  $c > 0$  and  $c \neq 1$ , that is the inverse of the exponential function  $y = c^x$

**logarithm**

- an exponent
- in  $x = c^y$ ,  $y$  is called the logarithm to base  $c$  of  $x$

**common logarithm**

- a logarithm with base 10

Write each of the following in logarithmic form

$$a) \underset{\text{Base}}{32} = \underset{\text{exp}}{2^5}$$

$$\log_2(32) = 5$$


$$b) 2^{-5} = \frac{1}{32}$$

$$\log_2\left(\frac{1}{32}\right) = -5$$

$$c) x = 10^y$$

$$\log_{10}(x) = y$$

$$\log x = y$$



Write each of the following in exponential form

$$a) \underset{\text{Base}}{\log_4 16} = \underset{\text{exp}}{2}$$

$$4^2 = 16$$

$$b) \log_2\left(\frac{1}{32}\right) = -5$$

$$2^{-5} = \frac{1}{32}$$

$$c) \log 65 = 1.8129$$

$$10^{1.8129} = 65$$

**Example 1****Evaluating a Logarithm**

Evaluate.

a)  $\log_7 49 = 2$       b)  $\log_6 1 = 0$       c)  $\log 0.001 = -3$       d)  $\log_2 \sqrt{8} = 1.5$

$$\frac{\log 49}{\log 7} = 2$$

$$\frac{\log 1}{\log 6} = 0$$

$$\frac{\log 0.001}{\log 10} = -3$$

$$\frac{\log \sqrt{8}}{\log 2} = 1.5$$

$$x = \log_7 49$$

$$7^x = 49$$

$$\cancel{7^x} = (\cancel{7})^2$$

$$x = 2$$

$$\boxed{\log_7 49 = 2}$$

$$x = \log_6 1$$

$$6^x = 1$$

$$\cancel{6^x} = (\cancel{6})^0$$

$$x = 0$$

$$\boxed{\log_6 1 = 0}$$

$$x = \log 0.001$$

$$10^x = 0.001$$

$$\cancel{10^x} = (\cancel{10})^{-3}$$

$$x = -3$$

$$\boxed{\log 0.001 = -3}$$

$$\cancel{2^x} = (\cancel{2})^{3/2}$$

$$x = \frac{3}{2}$$

$$\boxed{\log_2 \sqrt{8} = \frac{3}{2}}$$

**Example 2****Determine an Unknown in an Expression in Logarithmic Form**

Determine the value of  $x$ . (Convert to exponential form)

a)  $\log_5 x = -3$

b)  $\log_x 36 = 2$

c)  $\log_{64} x = \frac{2}{3}$

a)  $\log_5 x = -3$  (log. form)

$5^{-3} = x$  (exp. form)

$\left(\frac{1}{5}\right)^3 = x$

b)  $\log_x 36 = 2$  (log. form)

$x^2 = 36$  (exp. form)

$x = \pm 6$

$x = 6$

$c > 0$

$$\boxed{\frac{1}{125} = x}$$

c)  $\log_{64} x = \frac{2}{3}$  (log. form)


$64^{\frac{2}{3}} = x$  (exp. form)

$16 = x$

$$2^4 = 16 \quad (\text{exponential form})$$

$\uparrow$                      $\uparrow$   
Base                  ans.

$$\log_2(16) = 4 \quad (\text{logarithmic form})$$



**Example 3****Graph the Inverse of an Exponential Function**

- a) State the inverse of  $f(x) = 3^x$ .
- b) Sketch the graph of the inverse. Identify the following characteristics of the inverse graph:  $(x,y) \rightarrow (y,x)$
- the domain and range
  - the x-intercept, if it exists
  - the y-intercept, if it exists
  - the equations of any asymptotes

$$\text{a) } f(x) = 3^x$$

$y = 3^x$  passes the HLT

$$y = 3^x$$

$$x = 3^y \quad (\text{exp. form})$$

$$\log_3 x = y \quad (\text{log form})$$

$$y = \log_3 x$$

$$\boxed{f^{-1}(x) = \log_3 x}$$

Inverse

$$\text{b) } y = 3^x \quad (x,y) \rightarrow (y,x) \quad y = \log_3 x$$

x	y
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9

x	y
$\frac{1}{9}$	-2
$\frac{1}{3}$	-1
1	0
3	1
9	2

**Solution**

a) The inverse of  $y = f(x) = 3^x$  is  $x = 3^y$  or,


expressed in logarithmic form,  $y = \log_3 x$ . Since the inverse is a function, it can be written in function notation as

How do you know that  $y = \log_3 x$  is a function?

b) Set up tables of values for both the exponential function,  $f(x)$ , and its inverse,  $f^{-1}(x)$ . Plot the points and join them with a smooth curve.

$y = 3^x$	
$x$	$y$
-3	$\frac{1}{27}$
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9
3	27

$y = \log_3 x$	
$x$	$y$
$\frac{1}{27}$	-3
$\frac{1}{9}$	-2
$\frac{1}{3}$	-1
1	0
3	1
9	2
27	3



The graph of the inverse,  $f^{-1}(x) = \log_3 x$ , is a reflection of the graph

of  $f(x) = 3^x$  about the line  $y = x$ . For  $f^{-1}(x) = \log_3 x$ ,

- the domain is  $\{x | x > 0, x \in \mathbb{R}\}$  and the range is  $\{y | y \in \mathbb{R}\}$  or  $(-\infty, \infty)$
- the  $x$ -intercept is 1 or  $(1, 0)$
- there is no  $y$ -intercept
- the vertical asymptote, the  $y$ -axis, has equation  $x = 0$ ; there is no horizontal asymptote

How do the characteristics of  $f^{-1}(x) = \log_3 x$  compare to the characteristics of  $f(x) = 3^x$ ?

**Key Ideas**


- A logarithm is an exponent.
- Equations in exponential form can be written in logarithmic form and vice versa.

**Exponential Form      Logarithmic Form**

$$x = c^y \qquad y = \log_c x$$

- The inverse of the exponential function  $y = c^x$ ,  $c > 0$ ,  $c \neq 1$ , is  $x = c^y$  or, in logarithmic form,  $y = \log_c x$ . Conversely, the inverse of the logarithmic function  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$ , is  $x = \log_c y$  or, in exponential form,  $y = c^x$ .
- The graphs of an exponential function and its inverse logarithmic function are reflections of each other in the line  $y = x$ , as shown.
- For the logarithmic function  $y = \log_c x$ ,  $c > 0$ ,  $c \neq 1$ ,
  - the domain is  $\{x \mid x > 0, x \in \mathbb{R}\}$
  - the range is  $\{y \mid y \in \mathbb{R}\}$
  - the  $x$ -intercept is 1
  - the vertical asymptote is  $x = 0$ , or the  $y$ -axis
- A common logarithm has base 10. It is not necessary to write the base for common logarithms:

$$\log_{10} x = \log x$$



## Questions from Homework

a)  $y = 2^x$

$$x = 2^y$$

$$\log_2(x) = y$$

$$y = \log_2 x$$

D:  $\{x | x > 0, x \in \mathbb{R}\}$

R:  $\{y | y \in \mathbb{R}\}$

x-int:  $x=1$

y-int: none

VA:  $x=0$

b)  $y = \left(\frac{1}{3}\right)^x$

$$x = \left(\frac{1}{3}\right)^y$$

$$\log_{\frac{1}{3}}(x) = y$$

$$y = \log_{\frac{1}{3}} x$$

D:  $\{x | x > 0, x \in \mathbb{R}\}$

R:  $\{y | y \in \mathbb{R}\}$

x-int:  $x=1$

y-int: none

VA:  $x=0$

17. The growth of a new social networking site can be modelled by the exponential function  $N(t) = 1.1^t$ , where  $N$  is the number of users after  $t$  days.

- Write the equation of the inverse.
- How long will it take, to the nearest day, for the number of users to exceed 1 000 000?

$$\textcircled{2} \text{ If } 7^{2x} = y+3 \rightarrow \log_7(y+3) = 2x$$

↑  
 Base

↙  
 exp

↘  
 ans

$$\textcircled{3} \leftarrow \log_{10}(1000000) = 6 \longrightarrow 10^6 = 1000000$$

↑      ↑      ↑  
 Base    ans    exp.

$$\textcircled{4} \text{ c) } \log_4 \sqrt[3]{4} = \frac{1}{3}$$

$$\frac{\log(4^{\frac{1}{3}})}{\log 4} = \frac{1}{3}$$

$$x = \log_4(4)^{\frac{1}{3}} \quad (\text{log form})$$

$$\cancel{4^x} = 4^{\frac{1}{3}} \quad (\text{exp. form})$$

$$x = \frac{1}{3}$$

$$\textcircled{5} \quad a < \log_2 x < b \quad \begin{array}{l} 2^4 = 16 \\ 2^5 = 32 \end{array}$$

$$4 < \log_2 x < 5$$

$$\log_{10} 28 = 4.8$$

$$\frac{\log \bar{x}}{\log \sigma} = 4.8$$


$$\textcircled{12} \quad c) \quad \log_{\frac{1}{4}} x = -3 \quad \rightarrow \quad \left(\frac{1}{4}\right)^{-3} = x$$

Base ans exp


$$4^3 = x$$

$$64 = x$$

## 8.1 Understanding Logarithms, pages 380 to 382



- ii)  $y = \log_2 x$   
 iii) domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ , range  $\{y \mid y \in \mathbb{R}\}$ , x-intercept 1, no y-intercept, vertical asymptote  $x = 0$



- ii)  $y = \log_{\frac{1}{3}} x$   
 iii) domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ , range  $\{y \mid y \in \mathbb{R}\}$ , x-intercept 1, no y-intercept, vertical asymptote  $x = 0$

2. a)  $\log_{12} 144 = 2$

c)  $\log_{10} 0.00001 = -5$

3. a)  $5^2 = 25$

c)  $10^6 = 1\ 000\ 000$

4. a) 3

b) 0

5. a) 4; b) 5

b)  $\log_8 2 = \frac{1}{3}$


d)  $\log_7 (y + 3) = 2x$

b)  $8^{\frac{2}{3}} = 4$

d)  $11^y = x + 3$

c)  $\frac{1}{3}$

d) -3



domain  $\{x \mid x > 0, x \in \mathbb{R}\}$ , range  $\{y \mid y \in \mathbb{R}\}$ , x-intercept 1, no y-intercept, vertical asymptote  $x = 0$

10. They are reflections of each other in the line  $y = x$ .

11. a) They have the exact same shape.

b) One of them is increasing and the other is decreasing.

12. a) 216      b) 81      c) 64      d) 8

13. a) 7      b) 6

14. a) 0      b) 1

15. -1

16. 16

17. a)  $t = \log_{1.1} N$       b) 145 days

18. The larger asteroid had a relative risk that was 1479 times as dangerous.

19. 1000 times as great

20. 5

21.  $m = 14$ ,  $n = 13$

22.  $4n$

23.  $y = 3^{2^x}$

# Transformations of Logarithmic Functions

$$y = a \log_c(b(x-h)) + k$$

Focus on...

- explaining the effects of the parameters  $a$ ,  $b$ ,  $h$ , and  $k$  in  $y = a \log_c(b(x-h)) + k$  on the graph of  $y = \log_c x$ , where  $c > 1$
- sketching the graph of a logarithmic function by applying a set of transformations to the graph of  $y = \log_c x$ , where  $c > 1$ , and stating the characteristics of the graph

## Remember:

Parameter	Transformation
$a$	$(x, y) \rightarrow (x, ay)$
$b$	$(x, y) \rightarrow \left(\frac{x}{b}, y\right)$
$h$	$(x, y) \rightarrow (x + h, y)$
$k$	$(x, y) \rightarrow (x, y + k)$

**Example 1****Translations of a Logarithmic Function**

- a) Use transformations to sketch the graph of the function

$$y = \log_3(x + 9) + 2.$$

- b) Identify the following characteristics of the graph of the function.

- i) the equation of the asymptote
- ii) the domain and range
- iii) the y-intercept, if it exists
- iv) the x-intercept, if it exists

a)  $y = \log_3(x + 9) + 2$        $c = 3 \rightarrow \text{base}$

$a = 1 \rightarrow \text{no vertical stretch or reflection}$

$b = 1 \rightarrow \text{no horizontal stretch or reflection}$

$h = -9 \rightarrow 9 \text{ units left}$

$k = 2 \rightarrow 2 \text{ units up}$

$$(x, y) \rightarrow \left( \frac{1}{3}x + (-9), 1y + 2 \right)$$

$$(x, y) \rightarrow (x - 9, y + 2)$$

$$y = 3^x$$


x	y
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9

$$y = \log_3 x$$

x	y
$\frac{1}{9}$	-2
$\frac{1}{3}$	-1
1	0
3	1
9	2

$$(x, y) \rightarrow (x - 9, y + 2)$$

x	y
-8.8 or $-\frac{80}{9}$	0
-8.6 or $-\frac{86}{9}$	1
-8	2
-6	3
0	4



b) (0) VA:  $x = -9$

(iii) D:  $\{x | x > -9, x \in \mathbb{R}\}$  or  $(-9, \infty)$

R:  $\{y | y \in \mathbb{R}\}$

(iii) y int ( $x=0$ )

$$y = \log_3(x+9) + 2$$

$$y = \log_3(0+9) + 2$$

$$y = \boxed{\log_3(9)} + 2$$

$$y = \underline{2} + 2 \quad \frac{\log 9}{\log 3} = \underline{2}$$

$$y = 4$$

$$(0, 4)$$

(iv) x int ( $y=0$ )

$$y = \log_3(x+9) + 2$$

$$\textcircled{0} = \log_3(x+9) + 2$$

$$-2 = \log_3(x+9) \text{ (log form)}$$

$\begin{matrix} \uparrow \\ \text{exp} \end{matrix}$        $\begin{matrix} \uparrow \\ \text{base} \end{matrix}$        $\begin{matrix} \nearrow \\ \text{ans} \end{matrix}$

$$3^{-2} = x+9$$

$$\left(\frac{1}{3}\right)^2 = x+9$$

$$\frac{1}{9} = x+9$$

$$\frac{1}{9} - \frac{81}{9} = x$$

$$\frac{1}{9} - \frac{81}{9} = x$$

$$-\frac{80}{9} = x$$

$$(-8.8, 0)$$

**Example 2****Reflections, Stretches, and Translations of a Logarithmic Function**

- a) Use transformations to sketch the graph of the function

$$y = -\log_2(2(x+3))$$

- b) Identify the following characteristics of the graph of the function.

- i) the equation of the asymptote
- ii) the domain and range
- iii) the y-intercept, if it exists
- iv) the x-intercept, if it exists

a)  $c=2$  (Base)

$a=-1 \rightarrow$  reflected vertically in the x-axis  
and no vertical stretch

$b=2 \rightarrow$  no horizontal reflection in the y-axis.  
and horizontally stretched by a factor of  $\frac{1}{2}$

$h=-3 \rightarrow$  horizontal translation 3 units left

$k=0 \rightarrow$  no vertical translation


$$(x, y) \rightarrow \left(\frac{1}{2}x + (-3), -1y + 0\right)$$

$$(x, y) \rightarrow \left(\frac{1}{2}x - 3, -y\right)$$

$$\begin{array}{|c|c|} \hline y & = 2^x \\ \hline -2 & \frac{1}{4} \\ -1 & \frac{1}{2} \\ 0 & 1 \\ 1 & 2 \\ 2 & 4 \\ \hline \end{array}$$

$$\begin{array}{|c|c|} \hline y & = \log_2 x \\ \hline 1/4 & -2 \\ 1/2 & -1 \\ 1 & 0 \\ 2 & 1 \\ 4 & 2 \\ \hline \end{array}$$

$$\begin{array}{|c|c|} \hline (x, y) \rightarrow \left(\frac{1}{2}x - 3, -y\right) & \\ \hline -2.875 & -2\frac{3}{8} \\ -2.75 & -1\frac{1}{4} \\ -2.5 & -\frac{5}{2} \\ -2 & -2 \\ -1 & -1 \\ \hline \end{array}$$



b) (i) VA:  $x = -3$

(ii) D:  $\{x | x > -3, x \in \mathbb{R}\}$  or  $(-3, \infty)$

R:  $\{y | y \in \mathbb{R}\}$  or  $(-\infty, \infty)$

(iii) y-int ( $x=0$ )

$$y = -\log_2(2x+6)$$

$$y = -\log_2(2(0)+6)$$

$$y = -\boxed{\log_2(6)} \quad \frac{\log_2 6}{\log_2 2} = 2.58$$

$$y = -2.58$$

$$\boxed{y = -2.58}$$

(iv) x-int ( $y=0$ )

$$y = -\log_2(2x+6)$$

$$\frac{0}{-1} = \frac{-\log_2(2x+6)}{-1}$$

$$0 = \log_2(2x+6) \quad (\text{log})$$

$$2^0 = 2x+6 \quad (\text{exp})$$

$$1 = 2x+6$$

$$\frac{-5}{2} = \frac{2x}{2}$$

$$\boxed{-2.5 = x}$$

### Key Ideas

- To represent real-life situations, you may need to transform the basic logarithmic function  $y = \log_b x$  by applying reflections, stretches, and translations. These transformations should be performed in the same manner as those applied to any other function.
- The effects of the parameters  $a$ ,  $b$ ,  $h$ , and  $k$  in  $y = a \log_c(b(x - h)) + k$  on the graph of the logarithmic function  $y = \log_c x$  are shown below.

Vertically stretch by a factor of  $|a|$  about the  $x$ -axis. Reflect in the  $x$ -axis if  $a < 0$ .

Horizontally stretch by a factor of  $\frac{1}{|b|}$  about the  $y$ -axis. Reflect in the  $y$ -axis if  $b < 0$ .

Horizontally translate  $h$  units.

Vertically translate  $k$  units.

$$y = a \log_c(b(x - h)) + k$$

- Only parameter  $h$  changes the vertical asymptote and the domain. None of the parameters change the range.

## Homework

**Questions #1, 2, 4, 5, 8, 11 on  
page 389 - 391**