


Questions from Homework

$$\frac{-13\pi + 24\pi}{6} = \frac{11\pi}{6} \qquad -14\pi + 16\pi = 2\pi$$

$$\textcircled{5} \quad \csc^2\left(\frac{4\pi}{3}\right) \tan\left(\frac{15\pi}{4}\right) + \cos\left(\frac{-13\pi}{6}\right) - \sin\left(\frac{115\pi}{2}\right) + \cos(-14\pi)$$

$$\csc^2\left(\frac{4\pi}{3}\right) \tan\left(\frac{15\pi}{4}\right) + \cos\left(\frac{11\pi}{6}\right) - \sin\left(\frac{115\pi}{2}\right) + \cos(2\pi)$$



$$\boxed{\csc^2\left(\frac{4\pi}{3}\right)} \boxed{\tan\left(\frac{15\pi}{4}\right)} + \boxed{\cos\left(\frac{11\pi}{6}\right)} - \boxed{\sin\left(\frac{115\pi}{2}\right)} + \boxed{\cos(2\pi)}$$

$$\left(\frac{-2}{\sqrt{3}}\right)^2 (-1) + \left(\frac{\sqrt{3}}{2}\right) - (-1) + (1)$$

$$\left(\frac{4}{3}\right)(-1) + \frac{\sqrt{3}}{2} + 1 + 1$$


$$-\frac{4}{3} + \frac{\sqrt{3}}{2} + \frac{2}{1}$$

$$-\frac{8}{6} + \frac{3\sqrt{3}}{6} + \frac{12}{6}$$

$$\boxed{\frac{4 + 3\sqrt{3}}{6}} \quad \text{or} \quad \frac{3\sqrt{3} + 4}{6}$$

Questions from Homework

6) $\sec 15\pi + \sqrt{2} \sin \frac{39\pi}{4} \sin \frac{2\pi}{2} - \csc^2 \frac{100\pi}{3}$



$\sec 15\pi$ + $\sqrt{2} \sin \frac{39\pi}{4} \sin \frac{2\pi}{2} - \csc^2 \frac{100\pi}{3}$

$(-1) + \sqrt{2} \left(\frac{-1}{\sqrt{2}}\right) (1) - \left(\frac{2}{-\sqrt{3}}\right)^2$

$-1 + \sqrt{2} \left(\frac{-1}{\sqrt{2}}\right) (1) - \frac{4}{3}$

$-1 - \frac{\sqrt{2}}{\sqrt{2}} - \frac{4}{3}$

$-1 - 1 - \frac{4}{3}$

$-\frac{3}{3} - \frac{3}{3} - \frac{4}{3}$


$\frac{-10}{3}$

Questions from Homework

$$\textcircled{8} \quad \frac{\tan\left(-\frac{15\pi}{4}\right) + \sec\left(\frac{301\pi}{3}\right)}{\csc\left(\frac{25\pi}{2}\right) + \cot\left(-\frac{31\pi}{6}\right)}$$

$\xrightarrow{-\frac{15\pi}{4} + \frac{16\pi}{4} = \frac{\pi}{4}}$
 $\xrightarrow{-\frac{31\pi}{6} + \frac{36\pi}{6} = \frac{5\pi}{6}}$

$$\frac{\tan\left(\frac{\pi}{4}\right) + \sec\left(\frac{301\pi}{3}\right)}{\csc\left(\frac{25\pi}{2}\right) + \cot\left(\frac{5\pi}{6}\right)}$$



$$\frac{\tan\left(\frac{\pi}{4}\right) + \sec\left(\frac{301\pi}{3}\right)}{\csc\left(\frac{25\pi}{2}\right) + \cot\left(\frac{5\pi}{6}\right)}$$

$$\frac{1 + 2}{1 + (-\sqrt{3})}$$

$$\frac{3}{(1-\sqrt{3})(1+\sqrt{3})}$$

$$\frac{3+3\sqrt{3}}{1+\sqrt{3}-\sqrt{3}-3}$$

$$\boxed{\frac{3+3\sqrt{3}}{-2}} \quad \text{or} \quad \boxed{\frac{-3-3\sqrt{3}}{2}}$$

Introduction to Trigonometric Equations

trigonometric equation

- an equation involving trigonometric ratios

Focus on...

- algebraically solving first-degree and second-degree trigonometric equations in radians and in degrees
- verifying that a specific value is a solution to a trigonometric equation
- identifying exact and approximate solutions of a trigonometric equation in a restricted domain
- determining the general solution of a trigonometric equation

Did You Know?

In equations, mathematicians often use the notation $\cos^2 \theta$. This means the same as $(\cos \theta)^2$.

Let's start with basic LINEAR trigonometric equations...

...Pre-Calculus 110

Solve: $\sin \theta = 0.9659$, $-360^\circ < \theta < 720^\circ$
(Degrees)

- Reference angle?
- Which quadrants?
- Any co-terminal angles acceptable?

- If the domain is in degrees, give solutions in degrees.
- If the domain is in radians, give solutions in radians.

$\sin \theta = 0.9659$ use positive for $\bar{\theta}$ where is $\sin \theta > 0$ (positive)

$\bar{\theta} = \sin^{-1}(0.9659)$

$\bar{\theta} = 75^\circ$

Q1	Q2
$\theta = \bar{\theta}$	$\theta = 180^\circ - \bar{\theta}$
$\theta = 75^\circ$	$\theta = 180^\circ - 75^\circ = 105^\circ$
$\theta = 75^\circ - 360^\circ = -285^\circ$	$\theta = 105^\circ - 360^\circ = -255^\circ$
$\theta = 75^\circ + 360^\circ = 435^\circ$	$\theta = 105^\circ + 360^\circ = 465^\circ$

Solve: $\sec \theta = \frac{-1.3054}{1}$, $-2\pi \leq \theta \leq 2\pi$ (Approx. Value)
 (Radians) -6.28 6.28

* $\cos \theta = \frac{1}{-1.3054}$ where is $\cos < 0$ (negative)

$\cos \theta = -0.7660$

$\bar{\theta} = \cos^{-1}(0.7660)$

$\bar{\theta} = 0.7$ rads.

use positive for $\bar{\theta}$

Q2	Q3
----	----

$\theta = \pi - \bar{\theta}$

$\theta = \pi + \bar{\theta}$

$\theta = 3.14 - 0.7 = 2.44$ $\theta = 3.14 + 0.7 = 3.84$

$\theta = 2.44 - 6.28 = -3.84$ $\theta = 3.84 - 6.28 = -2.44$

Ex. $\sqrt{2} \cos \theta + 1 = 0, -360^\circ \leq \theta \leq 720^\circ$ (Degrees)

(Exact Value)

$$\frac{\sqrt{2} \cos \theta}{\sqrt{2}} = \frac{-1}{\sqrt{2}}$$

$\frac{S}{A}$
 $\frac{c}{c}$

$$\cos \theta = -\frac{1}{\sqrt{2}}$$

where is $\cos < 0$ (negative)

$$\bar{\theta} = \cos^{-1}\left(\frac{1}{\sqrt{2}}\right)$$

use positive for $\bar{\theta}$

$$\bar{\theta} = 45^\circ$$

Q2

Q3

$$\theta = 180^\circ - \bar{\theta}$$

$$\theta = 180^\circ + \bar{\theta}$$

$$\theta = 180^\circ - 45^\circ = 135^\circ$$

$$\theta = 180^\circ + 45^\circ = 225^\circ$$

$$\theta = 135^\circ - 360^\circ = -225^\circ$$

$$\theta = 225^\circ - 360^\circ = -135^\circ$$

$$\theta = 135^\circ + 360^\circ = 495^\circ$$

$$\theta = 225^\circ + 360^\circ = 585^\circ$$

Ex. $\sin x + 1 = 0$, $-2\pi \leq x \leq 4\pi$ (Radians)

(Exact Value)

$$\sin x = -1$$

$$\bar{x} = \frac{3\pi}{2} \text{ (Unit Circle)}$$

$$x = \frac{3\pi}{2}$$

$x = \frac{3\pi}{2} - \frac{2\pi}{1}$	$x = \frac{3\pi}{2} + \frac{2\pi}{1}$
$x = \frac{3\pi}{2} - \frac{4\pi}{2}$	$x = \frac{3\pi}{2} + \frac{4\pi}{2}$
$x = -\frac{\pi}{2}$	$x = \frac{7\pi}{2}$

Solutions:

$$x = -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}$$

① $\sin \theta = -\frac{\sqrt{3}}{2}$ where is $\sin \theta < 0$ $\frac{S}{T}$ $\frac{A}{E}$

$\bar{\theta} = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$ *use positive*

Q3	Q4
$\theta = 180^\circ + 60^\circ = 240^\circ$	$\theta = 360^\circ - 60^\circ = 300^\circ$
$240^\circ \pm 360^\circ n, n \in \mathbb{N}$	$300^\circ \pm 360^\circ n, n \in \mathbb{N}$

$\bar{\theta} = 60^\circ$

② $\cos \theta = -\frac{1}{2}$ where is $\cos \theta < 0$ $\frac{S}{T}$ $\frac{A}{C}$

$\bar{\theta} = \cos^{-1}\left(\frac{1}{2}\right)$

Q2	Q3
$\theta = 180^\circ - 60^\circ = 120^\circ$	$\theta = 180^\circ + 60^\circ = 240^\circ$
$120^\circ \pm 360^\circ n, n \in \mathbb{N}$	$240^\circ \pm 360^\circ n, n \in \mathbb{N}$

$\bar{\theta} = 60^\circ$

③ $\tan \theta = -\frac{\sqrt{3}}{3}$ where is $\tan \theta < 0$ $\frac{S}{T}$ $\frac{A}{E}$

$\tan \theta = -\frac{1}{\sqrt{3}}$

Q2	Q4
$\theta = 180^\circ - 30^\circ = 150^\circ$	$\theta = 360^\circ - 30^\circ = 330^\circ$
$150^\circ \pm 360^\circ n, n \in \mathbb{N}$	$330^\circ \pm 360^\circ n, n \in \mathbb{N}$

$\bar{\theta} = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$

$\bar{\theta} = 30^\circ$

④ $\tan \theta = \frac{\sqrt{3}}{1}$ where is $\tan \theta > 0$ $\frac{S}{T}$ $\frac{A}{C}$

$\bar{\theta} = \tan^{-1}\left(\frac{\sqrt{3}}{1}\right)$

Q1	Q3
$\theta = 60^\circ$	$\theta = 360^\circ - 60^\circ = 300^\circ$
$60^\circ \pm 360^\circ n, n \in \mathbb{N}$	$300^\circ \pm 360^\circ n, n \in \mathbb{N}$

$\bar{\theta} = 60^\circ$

⑤ $\sin \theta = -1$

* Unit Circle

$\theta = 270^\circ \rightarrow 270^\circ \pm 360^\circ n, n \in \mathbb{N}$

⑥ $5 \sin \theta - 4 = 0$ where is $\sin \theta > 0$ $\frac{S}{T}$ $\frac{A}{C}$

$\sin \theta = \frac{4}{5}$

$\sin \theta = 0.8$ (approx. value)

$\bar{\theta} = \sin^{-1}(0.8)$

$\bar{\theta} = 53.1^\circ$

Q1	Q2
$\theta = 53.1^\circ$	$\theta = 180^\circ - 53.1^\circ = 126.9^\circ$
$53.1^\circ \pm 360^\circ n, n \in \mathbb{N}$	$126.9^\circ \pm 360^\circ n, n \in \mathbb{N}$

Your Turn

Solve each trigonometric equation in the specified domain.

a) $3 \cos \theta - 1 = \cos \theta + 1, -2\pi \leq \theta \leq 2\pi$

b) $4 \sec x + 8 = 0, 0^\circ \leq x < 360^\circ$

Homework

Page 211 #1-5

Check-Up:

Solve:

$$\textcircled{1} \cot \theta = 0.7834, \quad -\frac{\pi}{2} < \theta < -\pi$$

$$\textcircled{2} 3\cos x + 5 = 6, \quad -360^\circ \leq x \leq 720^\circ$$

$$\textcircled{3} 2\csc x (1 - \csc x) = 0, \quad -4\pi < x < 4\pi$$

$$\textcircled{1} \cot \theta = 0.7834, \quad -\frac{\pi}{2} < \theta < -\pi$$

$$\textcircled{2} 3\cos x + 5 = 6, \quad -360^\circ \leq x \leq 720^\circ$$

$$\textcircled{3} 2\csc x (1 - \csc x) = 0, \quad -4\pi < x < 4\pi$$

Let's move onto QUADRATIC trigonometric equations...

...Pre-Calculus 110

- What strategies can we use to solve quadratic equations?
- Quadratic trigonometric equations will ultimately become TWO linear trigonometric equations.

Solve: $2x^2 + x = 1$

Solve: $2\sin^2 x + \sin x = 1, 0 \leq x \leq 4\pi$

$$\text{Ex. } \cos^2 \theta - \frac{1}{2} \cos \theta = 0, -2\pi \leq \theta \leq 4\pi$$

$$\text{Ex. } 6 \sin^2 x - \sin x = 2, -2\pi \leq \theta \leq 4\pi$$

Your Turn

Solve for θ .

$$\cos^2 \theta - \cos \theta - 2 = 0, 0^\circ \leq \theta < 360^\circ$$

Give solutions as exact values where possible. Otherwise, give approximate measures to the nearest thousandth of a degree.

General Solution of a Trigonometric Equation


Solve: $3\cos^2 \theta - \cos \theta = 2; \theta \in \mathbb{R}$

Determine the general solution for $\sin^2 x - 1 = 0$ over the real numbers if x is measured in radians.

Did You Know?

$2n$, where $n \in \mathbb{I}$,
represents all even
integers.

$2n + 1$, where $n \in \mathbb{I}$,
is an expression for all
odd integers.



Determine the general solution for $\cos^2 x - 1 = 0$, where the domain is real numbers measured in degrees.

Practice Problems:

Pages 212 - 214

#11 - 23

Check-Up problem...

Solve:

$$\sin x \sec x + 2 \sin x = 0, \quad x \in R \quad (x \text{ is measured in radians})$$

Unit Review...

What topics have we covered??

Review...

- C4 a)** Determine all solutions for the equation $2 \sin^2 \theta = 1 - \sin \theta$ in the domain $0^\circ \leq \theta < 360^\circ$.
- b)** Are your solutions exact or approximate? Why?
- c)** Show how you can check one of your solutions to verify its correctness.

A grandfather clock shows a time of 7 o'clock. What is the exact radian measure of the angle between the hour hand and the minute hand?

Determine the angular velocity of the minute hand on a clock.

Solve: $6 \sin^2 \theta - 3 \sin \theta = 0$, $0 \leq \theta \leq 360^\circ$

[A] $0^\circ, 30^\circ, 180^\circ, 330^\circ, 360^\circ$

[B] $0^\circ, 30^\circ, 180^\circ, 150^\circ, 360^\circ$

[C] $30^\circ, 90^\circ, 120^\circ, 270^\circ$

[D] $0^\circ, 180^\circ, 210^\circ, 330^\circ, 360^\circ$

If $\csc \theta < 0$ and $\tan \theta > 0$, then which of the following could be a possible measure of angle θ ?

[A] $\frac{11\pi}{6}$

[B] $\frac{4\pi}{3}$

[C] $\frac{3\pi}{4}$

[D] $\frac{\pi}{2}$

What is the principal angle of $-\frac{25\pi}{4}$?


[A] $\frac{3\pi}{4}$

[B] $\frac{\pi}{4}$

[C] $-\frac{\pi}{4}$

[D] $\frac{7\pi}{4}$

If the belt in the pulley system below travels 30 cm, what is the angle of rotation of the smaller pulley?



[A] $\frac{\pi}{9}$ radians

[B] 20°

[C] 20 radians

[D] 5°

Nibbles the hamster is running at 0.02 m/s on an exercise wheel of radius 8 cm. What is the angular velocity of this wheel?

[A] 0.15 rad/minute

[B] 240 rad/minute

[C] 0.25 rad/minute

[D] 15 radians/minute

Solve: $2(1 - \sin \theta)^2 + \sin \theta = 2(3 - 4 \sin^2 \theta)$, $-360^\circ \leq \theta \leq 720^\circ$

Little Johnny has a rock tied to the end of a piece of rope 1.5 m long and he is swinging it around his head in a circular pattern. Mrs. Centripetal, his physics teacher, is watching Johnny out the window of her physics lab and notes that the rock is making 12 revolutions every 48 seconds.

- (a) Determine the angular velocity with which little Johnny is twirling the rope above his head. [2]

- (b) The rock comes flying from the rope 3 minutes after Mrs. Centripetal started to time little Johnny. How far did the rock travel during the 3 minutes? [2]

Attachments

Worksheet - Sketching Angles in Radians.doc