

Correct Homework Sheet

$$\textcircled{2} \quad \frac{1-\cos^2\theta}{\sin^2\theta} = \frac{1-(\cos^2\theta-\sin^2\theta)}{\sin^2\theta}$$

$$\frac{1-\cos^2\theta+\sin^2\theta}{\sin^2\theta}$$

$$\frac{\sin^2\theta+\sin^2\theta}{\sin^2\theta}$$

$$\frac{2\sin^2\theta}{\sin^2\theta}$$

↓

$$\textcircled{3} \quad \frac{\sin(x+y)}{\sin(x-y)} = \cos^2 y - \cos^2 x$$

$$(\sin x \cos y + \cos x \sin y)(\sin x \cos y - \cos x \sin y)$$

$$\frac{\sin^2 x \cos^2 y - \cos^2 x \sin^2 y}{(\cos^2 x \cos^2 y - \cos^2 x)(1-\cos^2 y)}$$

$$\frac{\cos^2 y - \cos^2 x \cos^2 y - \cos^2 x + \cos^2 x \cos^2 y}{\cos^2 y - \cos^2 x}$$

$$\textcircled{5} \quad \tan^4 \theta = \sec^4 \theta (1 - 2\cos^2 \theta + \cos^4 \theta)$$

$$\frac{\sin^4 \theta}{\cos^4 \theta} \quad \frac{\sec^4 \theta (1-\cos^2 \theta)(1-\cos^2 \theta)}{\cos^4 \theta}$$

↓

$$\textcircled{1} \quad \frac{\cos\theta}{1+\sin\theta} + \frac{1+\sin\theta}{\cos\theta} = \boxed{2\sec\theta}$$

$$\frac{\cos^2\theta + (1+\sin\theta)(1+\sin\theta)}{\cos\theta(1+\sin\theta)}$$

$$2\left(\frac{1}{\cos\theta}\right)$$

$$\frac{\cos^2\theta + 1 + 2\sin\theta + \sin^2\theta}{\cos\theta(1+\sin\theta)}$$

$$\frac{2}{\cos\theta}$$

$$\frac{2+2\sin\theta}{\cos\theta(1+\sin\theta)}$$

$$\cancel{\frac{2(1+\sin\theta)}{\cos\theta(1+\sin\theta)}}$$

$$\frac{\partial}{\cos\theta}$$

$$\textcircled{10} \quad \frac{\tan^2\theta}{\tan^2\theta + 1} = \sin^2\theta$$

$$\boxed{\tan^2\theta} \div \sec^2\theta$$


$$\frac{\sin^2\theta}{\cos^2\theta} \div \frac{1}{\cos^2\theta}$$

$$\frac{\sin^2\theta}{\cos^2\theta} \times \cancel{\frac{1}{\cos^2\theta}}$$

$$\sin^2\theta$$

Negative Angles

$$\sin(-4x) = -\sin(4x) \quad \cos(-4x) = \cos(4x)$$



Ex: 7.2




① a) $y = \cos(-4x)$

Ex: 7.3

① d) $y = -\frac{1}{4} \csc(-8x)$

Derivatives of Trigonometric Functions**The Sine Function**

- We recall that the derivative $f'(x)$ of a function $f(x)$ gives the slope of the tangent.
- On the next slide we graph $f(x) = \sin x$ together with $f'(x)$, as determined by the slope of the tangent to the sine curve.
- Note that x is measured in radians.
- The derivative graph resembles the graph of the cosine!



Let's check this using the definition of a derivative...

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x + h) - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\sin x \left(\frac{\cos h - 1}{h} \right) + \cos x \left(\frac{\sin h}{h} \right) \right] \\
 &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h}
 \end{aligned}$$

■ Our calculations have brought us to four limits, two of which are easy:

■ Since x is constant while $h \rightarrow 0$,

$$\lim_{h \rightarrow 0} \sin x = \sin x \text{ and } \lim_{h \rightarrow 0} \cos x = \cos x$$

■ With some work we can also show that

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \text{ and } \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} = 0$$

■ Thus our guess is confirmed:

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\
 &= (\sin x) \cdot 0 + (\cos x) \cdot 1 = \cos x
 \end{aligned}$$

$$f(x) = \sin x$$

$$f'(x) = \cos x$$

Rules to differentiate trigonometric functions:

Given that "u" represents some differentiable function...

$$\frac{d}{du}(\sin u) = \cos u \bullet du$$

$$\frac{d}{du}(\csc u) = -\csc u \cot u \bullet du$$

$$\frac{d}{du}(\cos u) = -\sin u \bullet du$$

$$\frac{d}{du}(\sec u) = \sec u \tan u \bullet du$$

$$\frac{d}{du}(\tan u) = \sec^2 u \bullet du$$

$$\frac{d}{du}(\cot u) = -\csc^2 u \bullet du$$

$$\text{Ex: } f(x) = \tan(5x^2) \quad u=5x^2$$

$$f'(x) = \sec^2(5x^2) \cdot 10x \quad du=10x$$

$$f'(x) = 10x \sec^2(5x^2)$$

Let's Practice...

Differentiate the following:

$$y = \sin 3x \quad u = 3x \quad du = 3$$

$$y' = \cos(3x) \cdot 3$$

$$y' = 3\cos(3x)$$

$$y = \sin(x + 2) \quad u = x + 2 \quad du = 1$$

$$y' = \cos(x + 2) \cdot 1$$

$$y' = \cos(x + 2)$$

$$u = kx + d$$

$$\frac{du}{dx} = k$$

$$y = \sin(kx + d)$$

$$y' = \cos(kx + d) \cdot k$$

$$y' = k\cos(kx + d)$$

Ex #2.

Differentiate:

$$\begin{array}{c}
 u = x^3 \\
 du = 3x^2
 \end{array}
 \quad
 \begin{array}{c}
 y = (\sin x)^3 \\
 u = x \\
 du = 1
 \end{array}
 \quad
 \begin{array}{c}
 y = [\sin(x^2 - 1)]^3 \\
 u = x^2 - 1 \\
 du = 2x
 \end{array}$$

a) $y = \sin(x^3)$

$$\begin{aligned}
 y' &= \cos(x^3) \cdot 3x^2 \\
 y' &= 3x^2 \cos(x^3)
 \end{aligned}$$

b) $y = \sin^3 x$

$$\begin{aligned}
 y' &= 3(\sin x)^2 (\cos x) \cdot 1 \\
 y' &= 3 \sin^2 x \cos x
 \end{aligned}$$

c) $y = \sin^3(x^2 - 1)$

$$\begin{aligned}
 y' &= 3[\sin(x^2 - 1)]^2 (\cos(x^2 - 1)) \cdot 2x \\
 y' &= 6x \sin^2(x^2 - 1) \cos(x^2 - 1)
 \end{aligned}$$

Ex #3.

Differentiate:

Product rule $f'g + fg'$

$$y = (x^2)(\cos x)$$

$$y' = 2x \cos x + x^2(-\sin x)(1)$$

$$y' = 2x \cos x - x^2 \sin x$$

$$y' = x(2 \cos x - x \sin x)$$


Homework

Do Questions 1 and 3 from Exercise 7.2 Page 313

Worksheet on derivatives of trigonometric functions



Negative Angles



Attachments

Derivatives Worksheet.doc