

### Questions from homework

$$\textcircled{3} \text{ a) } f(x) = 2x^3 - 3x^2 \quad -2 \leq x \leq 2$$

$$f'(x) = 6x^2 - 6x$$

$$f'(x) = 6x(x-1)$$

$$(V: x=0, 1)$$

$$f(0) = 0 \quad (0, 0)$$

$$f(1) = -1 \quad (1, -1)$$

$$f(-2) = -16 - 12 = -28 \quad (-2, -28) \text{ abs min}$$

$$f(2) = 16 - 12 = 4 \quad (2, 4) \text{ abs max}$$


## The First Derivative Test

If  $f$  has a local maximum or minimum at  $c$ , then  $c$  must be a critical value of  $f$  (Fermat's Theorem), but not all critical numbers give rise to a maximum or minimum. For instance, recall that 0 is a critical number of the function  $y = x^3$  but this function has no maximum or minimum at a critical number.

One way of solving this is suggested by the figure below.

If  $f$  is increasing to the left of a critical number  $c$  and decreasing to the right of  $c$ , then  $f$  has a local max at  $c$ .

If  $f$  is decreasing to the left of a critical number  $c$  and increasing to the right of  $c$ , then  $f$  has a local min at  $c$ .



## The First Derivative Test


Let  $c$  be a critical number of a continuous function  $f$ .

1. If  $f'(x)$  changes from positive to negative at  $c$ , then  $f$  has a local max at  $c$ .
2. If  $f'(x)$  changes from negative to positive at  $c$ , then  $f$  has a local min at  $c$ .
3. If  $f'(x)$  does not change signs at  $c$ , then  $f$  has no max or min at  $c$ .

→  $f(x) = x^3$


$f'(x) = 3x^2$

CV:  $x = 0$



$f(0) = (0)^3 = 0$

$(0, 0)$  *no max or min*



**Example 1**

Find the local maximum and minimum values of

$$f(x) = x^3 - 3x + 1$$

$$f(-1) = -1 + 3 + 1 = 3$$

$(-1, 3)$  local max

$$f'(x) = 3x^2 - 3$$


$$f'(x) = 3(x^2 - 1)$$

$$f(1) = 1 - 3 + 1 = -1$$

$(1, -1)$  local min

$$f'(x) = 3(x+1)(x-1)$$

$$\text{CV: } x = \pm 1$$



**Example 2**


Find the local maximum and minimum values of  $g(x) = x^4 - 4x^3 - 8x^2 - 1$ . Use this information to sketch the graph of  $g$ .

$$g'(x) = 4x^3 - 12x^2 - 16x$$

$$g'(x) = 4x(x^2 - 3x - 4)$$

$$g'(x) = 4x(x+1)(x-4)$$

$$\text{CV: } x = -1, 0, 4$$



$$g(-1) = 1 + 4 - 8 - 1 = -4$$


$(-1, -4)$  local min

$$g(0) = -1$$

$(0, -1)$  local max

$$g(4) = 256 - 256 - 128 - 1 = -129$$

$(4, -129)$  local min



## The First Derivative Test

(for absolute extreme values)

Let  $c$  be a critical number of a continuous function  $f$ .

1. If  $f'(x)$  is positive for all  $x < c$  and  $f'(x)$  is negative for all  $x > c$ , then  $f(c)$  is the absolute maximum value.
2. If  $f'(x)$  is negative for all  $x < c$  and  $f'(x)$  is positive for all  $x > c$ , then  $f(c)$  is the absolute minimum value.

# Homework

Page 182 # 1-4