

# Transformations of Exponential Functions

## Focus on...

---

- applying translations, stretches, and reflections to the graphs of exponential functions
- representing these transformations in the equations of exponential functions
- solving problems that involve exponential growth or decay

**Link the Ideas**

The graph of a function of the form  $f(x) = a(c)^{b(x-h)} + k$  is obtained by applying transformations to the graph of the base function  $y = c^x$ , where  $c > 0$ .

Parameter	Transformation	Example
$a$	<ul style="list-style-type: none"> <li>Vertical stretch about the x-axis by a factor of <math> a </math></li> <li>For <math>a &lt; 0</math>, reflection in the x-axis</li> <li><math>(x, y) \rightarrow (x, ay)</math></li> </ul>	
$b$	<ul style="list-style-type: none"> <li>Horizontal stretch about the y-axis by a factor of <math>\frac{1}{ b }</math></li> <li>For <math>b &lt; 0</math>, reflection in the y-axis</li> <li><math>(x, y) \rightarrow (\frac{x}{b}, y)</math></li> </ul>	
$k$	<ul style="list-style-type: none"> <li>Vertical translation up or down</li> <li><math>(x, y) \rightarrow (x, y + k)</math></li> </ul>	
$h$	<ul style="list-style-type: none"> <li>Horizontal translation left or right</li> <li><math>(x, y) \rightarrow (x + h, y)</math></li> </ul>	

## Example 1

### Apply Transformations to Sketch a Graph

Consider the base function  $y = 3^x$ . For each transformed function,

- state the parameters and describe the corresponding transformations
- create a table to show what happens to the given points under each transformation

$y = 3^x$
$(-1, \frac{1}{3})$
$(0, 1)$
$(1, 3)$
$(2, 9)$
$(3, 27)$

- sketch the graph of the base function and the transformed function
- describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts

a)  $y = 2(3)^{x-4}$

b)  $y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

**Solution**


a) i) Compare the function  $y = 2(3)^{x-4}$  to  $y = a(c)^{b(x-h)} + k$  to determine the values of the parameters.

- $b = 1$  corresponds to no horizontal stretch.
- $a = 2$  corresponds to a vertical stretch of factor 2. Multiply the  $y$ -coordinates of the points in column 1 by 2.
- $h = 4$  corresponds to a translation of 4 units to the right. Add 4 to the  $x$ -coordinates
- $k = 0$  corresponds to no vertical translation.

ii) Add columns to the table representing the transformations.

$y = 3^x$	$y = 2(3)^{x-4}$
$(-1, \frac{1}{3})$	
(0, 1)	
(1, 3)	
(2, 9)	
(3, 27)	

iii) To sketch the graph, plot the points from column 3 and draw a smooth curve through them.



iv) The domain remains the same:  $\{x \mid x \in \mathbb{R}\}$ .

The range also remains unchanged:  $\{y \mid y > 0, y \in \mathbb{R}\}$ .

The equation of the asymptote remains as  $y = 0$ .

There is still no  $x$ -intercept, but the  $y$ -intercept changes to  $\frac{2}{81}$  or approximately 0.025.

$$b) y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$$

- i) state the parameters and describe the corresponding transformations
- ii) create a table to show what happens to the given points under each transformation
- iii) sketch the graph of the base function and the transformed function
- iv) describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts

$$b) y = \left(-\frac{1}{2}\right) (3)^{\frac{1}{5}x} - 5$$

(i)  $a = -\frac{1}{2} \rightarrow$  vertically stretched by a factor of  $\frac{1}{2}$  and reflected in the  $x$ -axis.

$b = \frac{1}{5} \rightarrow$  horizontally stretched by a factor of 5

$h = 0 \rightarrow$  No horizontal translation

$k = -5 \rightarrow$  translated 5 units down.

$$(x, y) \rightarrow \left[5x, -\frac{1}{2}y - 5\right]$$

(ii)

$$y = 3^x$$


x	y
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9

$$(x, y) \rightarrow [5x, -\frac{1}{2}y - 5]$$


Transformed

x	y
-10	$-\frac{9}{18}$ or $-5.0\bar{5}$
-5	$-\frac{3}{6}$ or $-5.1\bar{6}$
0	$-\frac{1}{2}$ or $-5.5$
5	$-\frac{13}{2}$ or $-6.5$
10	$-\frac{19}{2}$ or $-9.5$

(iii)



(iv)



$y = 3^x$   
 D:  $\{x | x \in \mathbb{R}\}$   
 R:  $\{y | y > 0, y \in \mathbb{R}\}$   
 HA:  $y = 0$   
 x-int: none  
 y-int:  $(0, 1)$

$y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$   
 D:  $\{x | x \in \mathbb{R}\}$   
 R:  $\{y | y < -5, y \in \mathbb{R}\}$   
 HA:  $y = -5$   
 x-int: none  
 y-int:  $(x=0)$

$y = -\frac{1}{2}(3)^{\frac{1}{5}(0)} - 5$

$y = -\frac{1}{2}(3)^0 - 5$

$y = -\frac{1}{2}(1) - 5$

$y = -\frac{1}{2} - \frac{10}{2} = -\frac{11}{2}$  or  $-5.5$

$(0, -5.5)$

**Your Turn** Page 351

Transform the graph of  $y = 4^x$  to sketch the graph of  $y = 4^{-2(x+5)} - 3$ . Describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts.

## Finding Intercepts:

Ex:  $y = 2^{-3(x+4)} - 4$

x int ( $y=0$ )

$$0 = 2^{-3(x+4)} - 4$$

$$4 = 2^{-3(x+4)}$$

$$\cancel{2^2} = \cancel{2}^{-3(x+4)}$$

$$\frac{2}{-3} = \frac{-3(x+4)}{-3}$$

$$-\frac{2}{3} = x+4$$

$$-\frac{2}{3} - \frac{4}{1} = x$$

$$\boxed{-\frac{14}{3} = x} \rightarrow \left(-\frac{14}{3}, 0\right)$$

y int ( $x=0$ )

$$y = 2^{-3(0+4)} - 4$$

$$y = 2^{-12} - 4$$

$$y = \left(\frac{1}{2}\right)^{12} - 4$$

$$y = \frac{1}{4096} - 4$$

$$\boxed{y = \frac{-16383}{4096} = -3.999} \rightarrow (0, -3.999)$$



## Homework

#1-7 and #10 on page 354

