

Transformations of Exponential Functions

Focus on...

- applying translations, stretches, and reflections to the graphs of exponential functions
- representing these transformations in the equations of exponential functions
- solving problems that involve exponential growth or decay

Link the Ideas

The graph of a function of the form $f(x) = a(c)^{b(x-h)} + k$ is obtained by applying transformations to the graph of the base function $y = c^x$, where $c > 0$.

Parameter	Transformation	Example
a	<ul style="list-style-type: none"> Vertical stretch about the x-axis by a factor of a For $a < 0$, reflection in the x-axis $(x, y) \rightarrow (x, ay)$ 	
b	<ul style="list-style-type: none"> Horizontal stretch about the y-axis by a factor of $\frac{1}{ b }$ For $b < 0$, reflection in the y-axis $(x, y) \rightarrow (\frac{x}{b}, y)$ 	
k	<ul style="list-style-type: none"> Vertical translation up or down $(x, y) \rightarrow (x, y + k)$ 	
h	<ul style="list-style-type: none"> Horizontal translation left or right $(x, y) \rightarrow (x + h, y)$ 	

Example 1

Apply Transformations to Sketch a Graph

Consider the base function $y = 3^x$. For each transformed function,

- state the parameters and describe the corresponding transformations
- create a table to show what happens to the given points under each transformation

$y = 3^x$
$(-1, \frac{1}{3})$
(0, 1)
(1, 3)
(2, 9)
(3, 27)


- sketch the graph of the base function and the transformed function
 - describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts
- $y = 2(3)^{x-4}$
 - $y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

Solution

- a) i) Compare the function $y = 2(3)^{x-4}$ to $y = a(c)^{b(x-h)} + k$ to determine the values of the parameters.
- $b = 1$ corresponds to no horizontal stretch.
 - $a = 2$ corresponds to a vertical stretch of factor 2. Multiply the y -coordinates of the points in column 1 by 2.
 - $h = 4$ corresponds to a translation of 4 units to the right. Add 4 to the x -coordinates of the points in column 2.
 - $k = 0$ corresponds to no vertical translation.
- ii) Add columns to the table representing the transformations.

$y = 3^x$	$y = 2(3)^{x-4}$
$(-1, \frac{1}{3})$	$(3, \frac{2}{3})$
$(0, 1)$	$(4, 2)$
$(1, 3)$	$(5, 6)$
$(2, 9)$	$(6, 18)$
$(3, 27)$	$(7, 54)$

- iii) To sketch the graph, plot the points from column 3 and draw a smooth curve through them.



- iv) The domain remains the same: $\{x \mid x \in \mathbb{R}\}$.

The range also remains unchanged: $\{y \mid y > 0, y \in \mathbb{R}\}$.

The equation of the asymptote remains as $y = 0$.

There is still no x -intercept, but the y -intercept changes to $\frac{2}{81}$ or approximately 0.025.

$$\text{b) } y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$$

- i) state the parameters and describe the corresponding transformations
- ii) create a table to show what happens to the given points under each transformation
- iii) sketch the graph of the base function and the transformed function
- iv) describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts

b) $y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

c = base = 3

(i) $a = -\frac{1}{2} \rightarrow$ vertical stretch by a factor of $\frac{1}{2}$ and a vertical reflection in the x-axis

$b = \frac{1}{5} \rightarrow$ horizontal by a factor of 5

$h = 0 \rightarrow$ no horizontal translation

$k = -5 \rightarrow$ vertical translation 5 units down

(ii) $(x, y) \rightarrow [5x, -\frac{1}{2}y - 5]$

$y = 3^x$

x	y
-2	$\frac{1}{9}$
-1	$\frac{1}{3}$
0	1
1	3
2	9

x	y
-10	$-\frac{1}{18} = -5.06$
-5	$-\frac{1}{6} = -5.17$
0	$-\frac{1}{2} = -5.5$
5	$-\frac{13}{2} = -6.5$
10	$-\frac{19}{2} = -9.5$

$$-\frac{1}{2}\left(\frac{1}{9}\right) - 5$$

$$\frac{-1}{18} - \frac{90}{18} = \frac{-91}{18}$$

$$-\frac{1}{2}\left(\frac{1}{3}\right) - 5$$

$$-\frac{1}{6} - \frac{30}{6} = \frac{-31}{6}$$

$$-\frac{1}{2}(1) - 5$$


$$-\frac{1}{2} - \frac{10}{2} = \frac{-11}{2}$$

$$-\frac{1}{2}(3) - 5$$

$$-\frac{3}{2} - \frac{10}{2} = \frac{-13}{2}$$

$$-\frac{1}{2}(9) - 5$$

$$-\frac{9}{2} - \frac{10}{2} = \frac{-19}{2}$$



(iii) D: $\{x | x \in \mathbb{R}\}$ or $(-\infty, \infty)$

R: $\{y | y < -5, y \in \mathbb{R}\}$ or $(-\infty, -5)$

HA: $y = -5$

x int ($y = 0$)

$y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

$0 = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

$5 = -\frac{1}{2}(3)^{\frac{1}{5}x}$
 $-\frac{1}{2} \quad \frac{1}{2}$
 $-\frac{1}{2} \quad -\frac{1}{2}$

$-10 = 3^{\frac{1}{5}x}$

you cannot take the log of a negative number

No x-intercept

(Check graph)

y int ($x = 0$)

$y = -\frac{1}{2}(3)^{\frac{1}{5}x} - 5$

$y = -\frac{1}{2}(3)^{\frac{1}{5}(0)} - 5$

$y = -\frac{1}{2}(3)^0 - 5$

$y = -\frac{1}{2}(1) - 5$

$y = -\frac{1}{2} - \frac{10}{2}$

$y = \frac{-11}{2} = -5.5$

$(0, -5.5)$

(Check graph)

Your Turn

Transform the graph of $y = 4^x$ to sketch the graph of $y = 4^{-2(x+5)} - 3$. Describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts.

$$y = \underline{\underline{1}}(\underline{\underline{4}})^{\underline{\underline{-2(x+5)}}} - \underline{\underline{3}}$$

$c = \text{base} = 4$

$a = 1$

$b = -2$

$h = -5$

$k = -3$

$y = 4^x$

x	y
-2	1/16
-1	1/4
0	1
1	4
2	16

x-int ($y = \underline{0}$)

$$y = 4^{-2(x+5)} - 3$$

$$0 = 4^{-2(x+5)} - 3$$

$$3 = 4^{-2(x+5)}$$

$$\hookrightarrow \frac{\log(3)}{\log(4)} = 0.79248$$

Base exponent

$$4^{0.79248} = 4^{-2(x+5)}$$

$$0.79248 = \frac{-2(x+5)}{-2}$$

$$-0.39624 = x+5$$

$$\boxed{-5.39624 = x}$$

$(-5.39624, 0)$

y-int ($x = \underline{0}$)

$$y = 4^{-2(x+5)} - 3$$

$$y = 4^{-2(0+5)} - 3$$

$$y = 4^{-2(5)} - 3$$

$$y = 4^{-10} - 3$$

$$y = \left(\frac{1}{4}\right)^{10} - 3$$

$$y = \frac{1}{1048576} - \frac{3}{1}$$

$$y = \frac{1}{1048576} - \frac{3145728}{1048576}$$

$$y = \frac{-3145727}{1048576} = \boxed{-2.9}$$

$(0, -2.9)$

Homework

#1-7 and #10 on page 354