

**Questions From Homework**

$$\text{Q3) } \int \frac{x+1}{x^2+2x-6} dx$$

$u = x^2 + 2x - 6$   
 $du = 2x + 2 dx$   
 $\frac{1}{2} du = x+1 dx$

$$= \int \frac{1}{u} \cdot \left( \frac{1}{2} du \right)$$

$$= \frac{1}{2} \int \frac{1}{u} du$$

$$= \frac{1}{2} \ln|u| + C$$

May 26-9:36 AM

**Questions From Homework**

$$\text{Q4) } \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos \theta}{\sin^3 \theta} d\theta = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1}{u^3} du$$

$u = \sin \theta$   
 $du = \cos \theta d\theta$   
 $d\theta = \frac{du}{\cos \theta}$

$$\frac{\theta}{\pi/3} \quad \frac{u}{1}$$

$$\frac{\pi/6}{\pi/3} \quad \frac{1}{1}$$

$$= \frac{1}{2u^2} \Big|_{\frac{1}{2}}^{\frac{1}{1}}$$

$$= -\frac{1}{2u^2} \Big|_{\frac{1}{2}}^{\frac{1}{1}}$$

$$= -\frac{1}{2} \Big|_{\frac{1}{2}}^{\frac{1}{1}}$$

$$= -\frac{1}{2} + \frac{4}{3} = \frac{3}{2}$$

May 22-8:21 AM

Warm Up

Find:  $\int 5x^2 \sin(4x^3 + 1) dx$

$$= -\frac{5}{12} \cos(4x^3 - 1) + C$$
  

$\int x \sqrt{2x^2 - 5} dx$

$$= \frac{(2x^2 - 5)^{3/2}}{6} + C$$
  

$\int \cot x dx$

$$= \ln|\sin x| + C$$

May 23-5:12 PM

**Differential and Integral Calculus 120****Integration by Parts**

$$f(x)g(x) = f'(x)g(x) + f(x)g'(x)$$

May 23-9:36 AM

As we have discussed before, every differentiation rule has a corresponding integration rule.

The rule that corresponds to the Product Rule for differentiation is called the rule for **integration by parts**.

The product rule stated that if  $f$  and  $g$  are differentiable functions, then

$$\frac{d}{dx} [f(x)g(x)] = f'(x)g'(x) + g(x)f'(x)$$

In the notation for indefinite integrals this equation becomes...  $\int [f(x)g'(x)dx + g(x)f'(x)dx] = \int f'(x)g(x)dx$

or  $\int f'(x)g'(x)dx + \int g(x)f'(x)dx = \int f'(x)g(x)dx$

which can be rearranged as:

$$\int f'(x)g(x)dx = f(x)g(x) - \int g(x)f'(x)dx$$

This formula above is called the **formula for integration by parts**.

It is perhaps easier to remember in the following notation... Let  $u = f(x)$  and  $v = g(x)$ . Then the differentials are  $du = f'(x)dx$  and  $dv = g'(x)dx$ .

And by the Substitution Rule, the formula becomes...

$$\int u dv = uv - \int v du$$

**Integration By Parts**

Let's do an example... Find  $\int x \sin x dx$

It helps when you stick to this pattern:

$$u = \underline{x} \quad dv = \underline{e^x dx}$$

$$du = \underline{1 dx} \quad v = \underline{e^x}$$

Again, the goal is to use integration by parts to obtain a simpler integral than the one we started with, so we must decide on what  $u$  and  $dv$  are very carefully!

In general, when deciding on a choice for  $u$  and  $dv$ , we usually try to choose  $u = f(x)$  to be a function that becomes simpler when differentiated... (or at least not more complicated) as long as  $dv = g(x)dx$  can be readily integrated to give  $v$ .

$$\int x \sin x dx = x \cos x - \int \cos x dx$$

$$= x \cos x + \sin x + C$$

May 23-4:34 PM

$$\text{Find: } \int xe^x dx = x e^x - \int e^x dx$$

It helps when you stick to this pattern:

$$u = \underline{x} \quad dv = \underline{e^x dx}$$

$$\int xe^x - e^x + C$$

May 23-5:33 PM

Find:  $\int x \cos(3x) dx = x \left( \frac{1}{3} \sin(3x) \right) - \int \frac{1}{3} \sin(3x) dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{x} \quad dv = \underline{\cos(3x)} dx \\ du &= \underline{1dx} \quad v = \underline{\frac{1}{3} \sin(3x)} \end{aligned}$$

$$\begin{aligned} &= \frac{1}{3} x \sin(3x) - \frac{1}{3} \int \sin(3x) dx \\ &= \frac{1}{3} x \sin(3x) - \frac{1}{3} \left( -\frac{1}{3} \cos(3x) \right) \\ &= \boxed{\frac{1}{3} x \sin(3x) + \frac{1}{9} \cos(3x) + C} \end{aligned}$$

Find:  $\int \ln x dx = x \ln x - \int \frac{1}{x} dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{\ln x} \quad dv = \underline{1} dx \\ du &= \underline{\frac{1}{x} dx} \quad v = \underline{\ln x} \end{aligned}$$

$$\begin{aligned} &= x \ln x - \int 1 dx \\ &= x \ln x - x + C \\ &= \boxed{x \ln x - x + C} \end{aligned}$$

May 23-5:33 PM

May 23-5:33 PM

Find:  $\int x^2 \sin(3x) dx = x^2 \left( \frac{1}{3} \cos(3x) \right) - \int \frac{1}{3} \cos(3x) 2x dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{x^2} \quad dv = \underline{\sin(3x)} dx \\ du &= \underline{2x dx} \quad v = \underline{-\frac{1}{3} \cos(3x)} \end{aligned}$$

$$\begin{aligned} &= -\frac{1}{3} x^2 \cos(3x) - \int \frac{2}{3} x \cos(3x) dx \\ &= -\frac{1}{3} x^2 \cos(3x) + \frac{2}{3} \left[ x \left( \frac{1}{3} \sin(3x) \right) - \int \frac{1}{3} \sin(3x) dx \right] \\ &= -\frac{1}{3} x^2 \cos(3x) + \frac{2}{3} \left[ \frac{1}{3} x \sin(3x) - \frac{1}{3} \int \sin(3x) dx \right] \\ &= -\frac{1}{3} x^2 \cos(3x) + \frac{2}{3} \left[ \frac{1}{3} x \sin(3x) + \frac{1}{9} \cos(3x) \right] \\ &= \boxed{-\frac{1}{3} x^2 \cos(3x) + \frac{2}{9} x \sin(3x) + \frac{2}{27} \cos(3x) + C} \\ &= -\frac{1}{3} x^2 \cos(3x) + \frac{2}{9} x \sin(3x) + \frac{2}{27} \cos(3x) + C \end{aligned}$$

May 23-5:33 PM

Find:  $\int x^2 e^x dx = x^2 e^x - \int e^x 2x dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{x^2} \quad dv = \underline{e^x} dx \\ du &= \underline{2x dx} \quad v = \underline{e^x} \end{aligned}$$

$$\begin{aligned} &= x^2 e^x - 2 \int x e^x dx \\ &= x^2 e^x - 2 \left[ x e^x - \int e^x dx \right] \\ &= x^2 e^x - 2 \left[ x e^x - e^x \right] \\ &= \boxed{x^2 e^x - 2x e^x + 2e^x + C} \\ &= x^2 e^x - 2x e^x + 2e^x + C \end{aligned}$$

May 23-5:33 PM

Find:  $\int x^2 \ln x dx = \frac{1}{3} x^3 \ln x - \int \frac{1}{3} x^3 \cdot \frac{1}{x} dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{\ln x} \quad dv = \underline{x^2 dx} \\ du &= \underline{\frac{1}{x} dx} \quad v = \underline{\frac{1}{3} x^3} \end{aligned}$$

$$\begin{aligned} &= \frac{1}{3} x^3 \ln x - \int \frac{1}{3} x^2 dx \\ &= \frac{1}{3} x^3 \ln x - \frac{1}{3} \left( \frac{1}{3} x^3 \right) \\ &= \frac{1}{3} x^3 \ln x - \frac{1}{9} x^3 \\ &= \boxed{\frac{1}{3} x^3 \ln x - \frac{1}{9} x^3 + C} \end{aligned}$$

May 23-5:33 PM

Find:  $\int e^x \sin x dx = e^x (-\cos x) - \int (-\cos x) e^x dx$

It helps when you stick to this pattern:

$$\begin{aligned} u &= \underline{e^x} \quad dv = \underline{\sin x dx} \\ du &= \underline{e^x dx} \quad v = \underline{-\cos x} \end{aligned}$$

$$\begin{aligned} &= e^x \sin x - \int e^x \cos x dx \\ &= e^x \sin x - e^x \cos x + \int e^x \cos x dx \\ &= e^x \sin x - e^x \cos x + e^x \sin x - \int e^x \sin x dx \\ &2 \int e^x \sin x dx = e^x \sin x - e^x \cos x \\ &\int e^x \sin x dx = \frac{1}{2} (e^x \sin x - e^x \cos x) + C \\ &= \boxed{\frac{1}{2} e^x (\sin x - \cos x) + C} \end{aligned}$$

this one as a little twist,  
because you cannot get to a simpler integral  
- rearrange for double the initial integral and  
divide by two!

May 23-5:33 PM

Find:

$$\int e^x \cos x dx$$

It helps when you  
stick to this pattern:

$$u = \underline{\quad} \quad dv = \underline{\quad}$$

$$du = \underline{\quad} \quad v = \underline{\quad}$$

this one as a little twist,  
because you cannot get to a simpler integral  
- rearrange for double the initial integral and  
divide by two!

$$= \frac{1}{2} e^x (\cos x + \sin x) + C$$

Find:

$$\int \underset{u}{\sin^{-1} x} \underset{dv}{dx}$$

$$= x \sin^{-1} x - \int \underset{(u)}{x \cdot \frac{1}{\sqrt{1-x^2}}} dx$$

It helps when you  
stick to this pattern:

$$u = \underline{\sin^{-1} x} \quad dv = \underline{1 dx}$$

$$du = \frac{1}{\sqrt{1-x^2}} dx \quad v = \underline{x}$$

$$du = \frac{1}{\sqrt{1-x^2}} dx$$

$$v = \underline{x}$$

may require substitution rule as well...

$$u = 1-x^2$$

$$du = -2x dx$$

$$\frac{1}{2} du = x dx$$

$$= x \sin^{-1} x - \int \frac{1}{(u)^{1/2}} \cdot \frac{(-1)}{\partial} du$$

$$= x \sin^{-1} x + \frac{1}{2} \int u^{-1/2} du$$

$$= x \sin^{-1} x + \frac{1}{2} \left( \frac{1}{2} u^{1/2} \right) + C$$

$$= x \sin^{-1} x + \frac{1}{4} u^{1/2} + C$$

$$= x \sin^{-1} x + \sqrt{1-x^2} + C$$

May 23-5:33 PM

May 23-5:33 PM

Homework - Exercise 11.4 - pp. 515 - Q. 1,2,4

May 26-2:03 PM

We've done this one already,  
but let's do it again and evaluate:

$$\int_1^e \ln x dx$$

It helps when you  
stick to this pattern:

$$u = \underline{\quad} \quad dv = \underline{\quad}$$

$$du = \underline{\quad} \quad v = \underline{\quad}$$



May 23-5:33 PM

Find:  $\int_0^{\pi/3} \sin x \ln(\cos x) dx$ 

It helps when you  
stick to this pattern:

$$u = \underline{\quad} \quad dv = \underline{\quad}$$

$$du = \underline{\quad} \quad v = \underline{\quad}$$



## Differential and Integral Calculus 120

Thursday, May 31, 2012

$$\int \text{Trigonometric Integrals!} \int$$

May 23-5:33 PM

Apr 9-10:38 AM

## WARM UP

Find:  $\int x^2 \sin x dx$

It helps when you stick to this pattern:

$$u = \underline{\hspace{2cm}} \quad dv = \underline{\hspace{2cm}}$$

$$du = \underline{\hspace{2cm}} \quad v = \underline{\hspace{2cm}}$$



## Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

recall double angle identities:

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

$$\cos 2\theta = 2 \cos^2 \theta - 1$$

$$\cos 2\theta = 1 - 2 \sin^2 \theta$$

also we will make use of the Half-angle identities:

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

May 23-5:33 PM

May 26-1:18 AM

Let's start off by doing some integration of trigonometric functions that require direct substitution and practice using our identities:

$$\int \sin x \cos x dx$$



$$\int \cos^2 x dx$$



$$\int \sin^3 x dx$$



$$\int \cos^3 x dx$$




$$\int_0^{\pi/2} \cos^3 x \sin^4 x dx$$



May 26-1:23 AM

$$\int \sin^2 x \cos x dx$$



$$\int \sin^5 x \cos^2 x dx$$

$$\int \sin^2 x dx$$

$$\int \sin^2 x \cos^2 x dx$$

$$\int \sin^4 x dx$$

may have to use the half-angle identity twice

Strategy for Evaluating  $\int \sin^n x \cos^m x dx$

(a) If the power of cosine is odd ( $m = 2k + 1$ ), save one cosine factor and use  $\cos^2 x = 1 - \sin^2 x$  to express the remaining factors in terms of sine:

$$\int \sin^n x \cos^{2k+1} x dx = \int \sin^n x (\cos^2 x)^k \cos x dx$$

$$= \int \sin^n x (1 - \sin^2 x)^k \cos x dx$$

Then substitute  $u = \sin x$

(b) If the power of sine is odd ( $n = 2k + 1$ ), save one sine factor and use  $\sin^2 x = 1 - \cos^2 x$  to express the remaining factors in terms of cosine:

$$\int \sin^{2k+1} x \cos^n x dx = \int (\cos^2 x)^k \cos^n x \sin x dx$$

$$= \int (1 - \cos^2 x)^k \cos^n x \sin x dx$$

Then substitute  $u = \cos x$  (Note that if the powers of both sine and cosine are odd, either (a) or (b) can be used)

(c) If the powers of both sine and cosine are even, use the half-angle identities:

$$\sin^2 x = (\sin x)^2 \quad \cos^2 x = (\cos x)^2$$

It is sometimes helpful to use the identity

$$\sin x \cos x = \frac{1}{2} \sin 2x$$

May 26-1:38 AM

Evaluate:

$$\int \tan^6 x \sec^4 x dx$$



$$\int \tan^5 x \sec^7 x dx$$



Strat:

Strategy for Evaluating  $\int \tan^n x \sec^m x dx$

(a) If the power of secant is even ( $m = 2k$ ), save a factor of  $\sec^2 x$  and use  $\sec^2 x = 1 + \tan^2 x$  to express the remaining factors in terms of tan x:

$$\int \tan^n x \sec^{2k} x dx = \int \tan^n x (\sec^2 x)^{k-1} \sec^2 x dx$$

$$= \int \tan^n x (1 + \tan^2 x)^{k-1} \sec^2 x dx$$

Then substitute  $u = \tan x$ .  
(b) If the power of tangent is odd ( $n = 2k + 1$ ), save a factor of  $\sec x \tan x$  and use  $\tan^2 x = \sec^2 x - 1$  to express the remaining factors in terms of sec x:

$$\int \tan^{2k+1} x \sec^m x dx = \int (\tan^2 x)^k \sec^m x \sec x \tan x dx$$

$$= \int (\sec^2 x - 1)^k \sec^{m-1} x \sec x \tan x dx$$

Then substitute  $u = \sec x$ .

We will also need to know the indefinite integrals of  $\tan x$  (already found this one) and  $\sec x$  when integrating these types of functions.

$$\int \tan x dx = \ln |\sec x| + C$$

$$\int \sec x dx = \ln |\sec x + \tan x| + C$$

Find:

$$\int \tan^3 x dx$$



$$\int \sec^3 x dx$$



here, we can use integration by parts and rearrange for double the original integral.

May 26-1:38 AM

May 26-1:38 AM

Homework - Exercise 7.2

# Differential and Integral Calculus 120

Friday, May 28, 2010


$\int$  Trigonometric Substitution!  $\int$

May 30-9:35 PM

Apr 9-10:38 AM

**Exercises 7.2 □ page 482**
**Exercises 7.2 □ page 482**

1.  $\frac{1}{3} \cos^3 x - \frac{1}{5} \cos^5 x + C$     3.  $-\frac{11}{384}$   
 5.  $\frac{1}{2} \sin^5 x - \frac{2}{7} \sin^7 x + \frac{1}{9} \sin^9 x + C$     7.  $\pi/4$   
 9.  $\frac{3}{8}x + \frac{1}{4} \sin 2x + \frac{1}{32} \sin 4x + C$   
 11.  $(3x/2) + \cos 2x - \frac{1}{8} \sin 4x + C$     13.  $(3\pi/4) - 4/192$   
 15.  $\left[ \frac{1}{2} \cos^3 x - \frac{2}{5} \cos x \right] \sqrt{\cos x + C}$   
 17.  $\frac{1}{2} \cos^2 x - \ln |\cos x| + C$     19.  $\ln(1 + \sin x) + C$   
 21.  $\tan x - x + C$     23.  $\tan x + \frac{1}{3} \tan^3 x + C$     25.  $\frac{1}{5}$   
 27.  $\frac{1}{3} \sec^3 x - \sec x + C$     29.  $\frac{38}{15}$   
 31.  $\frac{1}{4} \sec^4 x - \tan^2 x + \ln |\sec x| + C$     33.  $\frac{1}{2} \tan^2 x + C$   
 35.  $\sqrt{3} - (\pi/3)$   
 37.  $-\frac{1}{3} \cot^3 w - \frac{1}{5} \cot^5 w + C$     39.  $\ln |\csc x - \cot x| + C$   
 41.  $\frac{1}{2} [\frac{1}{3} \sin 3x - \frac{1}{7} \sin 7x] + C$     43.  $\frac{1}{4} \sin 2\theta + \frac{1}{24} \sin 12\theta + C$   
 45.  $\frac{1}{2} \sin 2x + C$   
 47.  $-\frac{1}{3} \cos^5 x + \frac{2}{5} \cos^3 x - \cos x + C$



May 30-9:33 PM

**Trigonometric Substitution**

This is an integration technique introduced as a means of evaluating integrals involving the radical forms:

$$\sqrt{a^2 - x^2} \quad \sqrt{a^2 + x^2} \quad \sqrt{x^2 - a^2}$$

| For integrals involving: | let                 | then                               |
|--------------------------|---------------------|------------------------------------|
| $\sqrt{a^2 - x^2}$       | $x = a \sin \theta$ | $\sqrt{a^2 - x^2} = a \cos \theta$ |
| $\sqrt{a^2 + x^2}$       | $x = a \tan \theta$ | $\sqrt{a^2 + x^2} = a \sec \theta$ |
| $\sqrt{x^2 - a^2}$       | $x = a \sec \theta$ | $\sqrt{x^2 - a^2} = a \tan \theta$ |

Just as with algebraic substitution, our objective with trigonometric substitution is to eliminate the radicals in the integrand. There are three substitutions that accomplish this objective for the three types of radicals outlined in the table above.

$$x \neq a$$

To show that the radical is eliminated as indicated in each of the three cases, we need to use the following trigonometric identities:  $\sin^2 \theta + \cos^2 \theta = 1$

|                                     |  |
|-------------------------------------|--|
| $1 - \sin^2 \theta = \cos^2 \theta$ | recall double angle identity:<br>$\sin 2\theta = 2 \sin \theta \cos \theta$                            |
| $1 + \tan^2 \theta = \sec^2 \theta$ | also we will make use of the<br>Half-angle identities:<br>$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$ |
| $\sec^2 \theta - 1 = \tan^2 \theta$ | $\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$   |

We will always manipulate the radical first so that we have the constant...  $a = 1$

It also helps to build a triangle marking angle theta.



May 26-12:10 AM

use the help of a triangle to define theta before substitution

$$\int \sqrt{1-x^2} dx$$

$$\int \sqrt{4-x^2} dx$$

May 26-1:38 AM

use the help of a triangle to define theta before substitution

$$\int \frac{1}{\sqrt{1-x^2}} dx$$

$$\int \frac{1}{x^2 \sqrt{1-x^2}} dx$$

$$\int \frac{dx}{x^2 \sqrt{9-x^2}}$$

May 26-1:38 AM

$$\int \frac{1}{\sqrt{9-49x^2}} dx$$

$$\int \sqrt{16+25x^2} dx$$

May 26-11:53 PM

$$\int \frac{1}{\sqrt{16+36x^2}} dx$$

May 26-11:55 PM

Use trigonometric substitution to find each of the following:

$$\int \sqrt{36-25x^2} dx$$

Jun 1-11:12 PM

Find the area enclosed by the ellipse:

$$\frac{x^2}{25} + \frac{y^2}{4} = 1$$

sketch (find area in first quadrant from x = 0 to x = 5)

y = ?

Homework - Exercise 7.3

## Differential and Integral Calculus 120

Monday, May 31, 2010

Partial Fractions!

May 30-9:35 PM

Apr 9-10:38 AM

**Exercises 7.3 □ page 488**

1.  $\sqrt{x^2 - 9}/(9x) + C$     3.  $\frac{1}{3}(x^2 - 18)\sqrt{x^2 + 9} + C$   
 5.  $\pi/24 + \sqrt{3}/8 - \frac{1}{4}$     7.  $-\sqrt{25 - x^2}/(25x) + C$   
 9.  $(1/\sqrt{3}) \ln |(\sqrt{x^2 + 3} - \sqrt{3})/x| + C$   
 11.  $\frac{1}{4} \sin^{-1}(2x) + \frac{1}{2}x\sqrt{1 - 4x^2} + C$   
 13.  $\sqrt{9x^2 - 4} - 2 \sec^{-1}(3x/2) + C$   
 15.  $(x/\sqrt{a^2 - x^2}) - \sin^{-1}(x/a) + C$     17.  $\sqrt{x^2 - 7} + C$   
 19.  $\ln(1 + \sqrt{2})$     21.  $\frac{64}{1215}$   
 23.  $\frac{1}{2}[\sin^{-1}(x - 1) + (x - 1)\sqrt{2x - x^2}] + C$   
 25.  $\frac{1}{3} \ln |3x + 1 + \sqrt{9x^2 + 6x - 8}| + C$   
 27.  $\frac{1}{2}[\tan^{-1}(x + 1) + (x + 1)/(x^2 + 2x + 2)] + C$   
 29.  $\frac{1}{2}[e^t\sqrt{9 - e^{2t}} + 9 \sin^{-1}(e^t/3)] + C$   
 33.  $3\pi/2$     37. 0.81, 2; 2.10  
 39.  $r\sqrt{R^2 - r^2} + \pi r^2/2 - R^2 \arcsin(r/R)$     41.  $2\pi^2 R r^3$

**WARM UP**

Find the area enclosed by the ellipse:

$$\frac{x^2}{16} + \frac{y^2}{36} = 1$$

sketch (find area in first quadrant from x = 0 to x = 4)


y = ?




May 30-9:36 PM

May 26-2:31 AM

$$\int \frac{1}{x^2\sqrt{x^2+4}} dx$$

recall:  $\cot^2 \theta + 1 = \csc^2 \theta$ 

$$\int \frac{\sqrt{9-x^2}}{x^2} dx$$



May 28-1:35 AM

May 27-12:34 AM

**Integration of Rational Functions by Partial Fractions**

This technique allows us to integrate any rational function (a ratio of polynomials)

$$f(x) = \frac{P(x)}{Q(x)}$$

by expressing it as a sum of simpler fractions, called partial fractions, that we already know how to integrate.

Let's do an example....  $\int \frac{x+5}{x^2+x-2} dx$

- rearrange
- collect & equate like terms
- and use system of equations to solve for A and B

May 26-2:11 AM

Find:  $\int \frac{x+7}{x^2-x-6} dx$



May 30-6:13 PM

Find:  $\int \frac{x^2 + 2x - 1}{2x^3 + 3x^2 - 2x} dx$

Find:  $\int \frac{1}{2x^2 + 5x + 2} dx$

May 30-6:42 PM

May 30-6:38 PM

Find:  $\int \frac{5x^2 + 15x - 36}{x^3 - 9x} dx$

The previous examples dealt with the denominator being a **product of distinct linear factors**. (**CASE I**)

In the following example, we look at the case when the denominator is the product of linear factors, however some are **repeated**.

If we do it the same way as before, solving for A and B using a system of equations doesn't work as all terms in the elimination cancel.

(**CASE II**)

So...by inspecting the denominator... for each factor of the form  $(px + q)^n$ , the partial fraction decomposition must include the following sum of m fractions:

$$\frac{A}{px+q} + \frac{B}{(px+q)^2} + \frac{C}{(px+q)^3} + \dots + \frac{\text{const}}{(px+q)^m}$$

Let's do the example...

$$\int \frac{6x+7}{(x+2)^2} dx$$

multiply by common denominator  
 $(x+2)^2$  ... and solve for A and B!

- then integrate!

May 30-6:42 PM

May 30-6:49 PM

Find:  $\int \frac{5x^2 + 20x + 6}{x^3 + 2x^2 + x} dx$

simplify to...

repeating linear factor

(**CASE III**)

So...by inspecting the denominator Q(x)... for each factor of the form  $(ax^2 + bx + c)$ , which is an **irreducible quadratic factor**, the partial fraction decomposition must take the form:

$$\frac{Ax+B}{ax^2+bx+c}$$

if the irreducible quadratic factor is repeating  $(ax^2 + bx + c)^n$ , the partial fraction decomposition must include the following sum of n fractions:

$$\frac{Ax+B}{(ax^2+bx+c)} + \frac{Cx+D}{(ax^2+bx+c)^2} + \dots + \frac{\text{const}(x) + \text{const}}{(ax^2+bx+c)^n}$$

Here are some examples of how we would do the partial fraction decomposition for some rational functions:

(we will not solve these - they are a little tedious to say the least to find all the coefficients)

$$\frac{x^3 - x + 1}{x^2(x-1)^3} =$$

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^2} =$$

May 30-6:42 PM

May 30-6:42 PM

Find:  $\int \frac{2x^3 - 4x - 8}{(x^2 - x)(x^2 + 4)} dx$

Evaluate:  $\int_0^1 \frac{3x + 4}{x^3 - 2x - 4} dx$

May 30-6:42 PM

May 30-6:42 PM

Don't lose sight of easier methods when possible....

Sometimes partial fractions can be avoided when integrating a rational function.

example, Find:

$$\int \frac{x^2 + 1}{x(x^2 + 3)} dx$$

$$= \int \frac{x^2 + 1}{x^3 + 3x} dx$$

how about straight up substitution...

Find:  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$

May 31-12:18 AM

May 30-6:42 PM

Find:  $\int \frac{-2x + 4}{(x^2 + 1)(x - 1)^2} dx$

Find:  $\int \frac{1 - x + 2x^2 - x^3}{x(x^2 + 1)^2} dx$

May 30-6:42 PM

May 30-6:42 PM

**(CASE IV)**

For the case of a rational function:

$$f(x) = \frac{P(x)}{Q(x)}$$

where the degree of the denominator  $Q(x)$  is less than or equal to the degree of the numerator  $P(x)$ ,

(this is called an improper rational function)

then divide  $P(x)$  by  $Q(x)$  by long division to obtain a quotient  $S(x)$  and the remainder  $R(x)$ .

$$f(x) = \frac{P(x)}{Q(x)} = S(x) + \frac{R(x)}{Q(x)}$$

Example Find:  $\int \frac{x^3 + x}{x - 1} dx$

May 30-6:42 PM

Find:

$$= \int \frac{5x^3 - x^2}{x^2 - 1} dx$$

May 30-6:42 PM

Find:  $\int \frac{2x^5 - 5x}{(x^2 + 2)^2} dx$ 

long division case

$$= \int 2x - \frac{8x^3 + 13x}{(x^2 + 2)^2} dx$$

May 30-6:42 PM

Evaluate:  $\int_3^4 \frac{x+4}{x-2} dx$ 

May 30-6:42 PM

Find:  $\int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx$ Rationalizing Substitutions:

$$\text{Find: } \int \frac{\sqrt{x+4}}{x} dx = 2 \int \frac{u^2}{u^2 - 4} du$$

Let:

- integrate by partial fractions using Case IV technique (long division + Remainder) as the degree of the numerator equals the degree of the denominator

$$= 2 \int \left( 1 + \frac{4}{u^2 - 4} \right) du$$

integrate by partial fractions using Case I technique

May 30-6:42 PM

May 30-6:42 PM

Find:

May 30-6:42 PM