

Page 53

Q 8) $f(x) = x^2 + 2$, $x \leq 0$

$$y = x^2 + 2$$

$$x = y^2 + 2$$

$$x - 2 = y^2$$

$$\pm \sqrt{x-2} = y$$

$$y = -\sqrt{x-2}$$


$$f^{-1}(x) = -\sqrt{x-2}$$

$$f(x) = x^2 + 2$$

x	y
0	2
-1	3
-2	6
-3	11

$$f^{-1}(x) = -\sqrt{x-2}$$

x	y
2	0
3	-1
6	-2
11	-3



D: $\{x | x \leq 0, x \in \mathbb{R}\} \cup (-\infty, 0]$

R: $\{y | y \geq 2, y \in \mathbb{R}\} \cup [2, \infty)$

D: $\{x | x \geq 2, x \in \mathbb{R}\} \cup [2, \infty)$

R: $\{y | y \leq 0, y \in \mathbb{R}\} \cup (-\infty, 0]$

Radical Functions and Transformations

Focus on...

- investigating the function $y = \sqrt{x}$ using a table of values and a graph
- graphing radical functions using transformations
- identifying the domain and range of radical functions

radical function

- a function that involves a radical with a variable in the radicand
- $y = \sqrt{3x}$ and $y = 4\sqrt[3]{5} + x$ are radical functions.

$$y = \sqrt{x}$$

x	y
0	0
1	1
4	2
9	3
16	4

Example 1**Graph Radical Functions Using Tables of Values**

Use a table of values to sketch the graph of each function.
Then, state the domain and range of each function.


a) $y = \sqrt{x}$ b) $y = \sqrt{x - 2}$ c) $y = \sqrt{x} - 3$

- a) For the function $y = \sqrt{x}$, the radicand x must be greater than or equal to zero, $x \geq 0$.

D: $x \geq 0$

x	y
0	0
1	1
4	2
9	3
16	4
25	5

How can you choose values of x that allow you to complete the table without using a calculator?



The graph has an endpoint at $(0, 0)$ and continues up and to the right. The domain is $\{x | x \geq 0, x \in \mathbb{R}\}$. The range is $\{y | y \geq 0, y \in \mathbb{R}\}$.

Ex: $2x + 7 \geq 0$

$2x \geq -7$

$x \geq -\frac{7}{2}$

D: $\{x | x \geq 0, x \in \mathbb{R}\}$
 $[0, \infty)$

R: $\{y | y \geq 0, y \in \mathbb{R}\}$
 $[0, \infty)$

- b) For the function $y = \sqrt{x - 2}$, the value of the radicand must be greater than or equal to zero.

D: $x - 2 \geq 0$

$x \geq 2$

x	y
2	0
3	1
6	2
11	3
18	4
27	5


How is this table related to the table for $y = \sqrt{x}$ in part a)?

How does the graph of $y = \sqrt{x - 2}$ compare to the graph of $y = \sqrt{x}$?

$h = 2 \rightarrow$ translated 2 units right

D: $\{x | x \geq 2, x \in \mathbb{R}\}$
 $[2, \infty)$

R: $\{y | y \geq 0, y \in \mathbb{R}\}$
 $[0, \infty)$



The domain is $\{x | x \geq 2, x \in \mathbb{R}\}$. The range is $\{y | y \geq 0, y \in \mathbb{R}\}$.

- c) The radicand of $y = \sqrt{x} - 3$ must be non-negative.

D: $x \geq 0$


x	y
0	-3
1	-2
4	-1
9	0
16	1
25	2

How does the graph of $y = \sqrt{x} - 3$ compare to the graph of $y = \sqrt{x}$?

$k = -3 \rightarrow$ translated 3 units down

D: $\{x | x \geq 0, x \in \mathbb{R}\}$
 $[0, \infty)$

R: $\{y | y \geq -3, y \in \mathbb{R}\}$
 $[-3, \infty)$



The domain is $\{x | x \geq 0, x \in \mathbb{R}\}$ and the range is $\{y | y \geq -3, y \in \mathbb{R}\}$.

Graphing Radical Functions Using Transformations

You can graph a radical function of the form $y = a\sqrt{b(x - h)} + k$ by transforming the graph of $y = \sqrt{x}$ based on the values of a , b , h , and k . The effects of changing parameters in radical functions are the same as the effects of changing parameters in other types of functions.

- Parameter a results in a vertical stretch of the graph of $y = \sqrt{x}$ by a factor of $|a|$. If $a < 0$, the graph of $y = \sqrt{x}$ is reflected in the x -axis.
- Parameter b results in a horizontal stretch of the graph of $y = \sqrt{x}$ by a factor of $\frac{1}{|b|}$. If $b < 0$, the graph of $y = \sqrt{x}$ is reflected in the y -axis.
- Parameter h determines the horizontal translation. If $h > 0$, the graph of $y = \sqrt{x}$ is translated to the right h units. If $h < 0$, the graph is translated to the left $|h|$ units.
- Parameter k determines the vertical translation. If $k > 0$, the graph of $y = \sqrt{x}$ is translated up k units. If $k < 0$, the graph is translated down $|k|$ units.

Chapter 3

Domain:

$$\{x | x \geq h, x \in \mathbb{R}\} \quad (b > 0)$$

$$\{x | x \leq h, x \in \mathbb{R}\} \quad (b < 0)$$

Range:

$$\{y | y \geq k, y \in \mathbb{R}\} \quad (a > 0)$$

$$\{y | y \leq k, y \in \mathbb{R}\} \quad (a < 0)$$

Example 2**Graph Radical Functions Using Transformations**

Sketch the graph of each function using transformations. Compare the domain and range to those of $y = \sqrt{x}$ and identify any changes.

a) $y = 3\sqrt{-(x - 1)}$ b) $y - 3 = -\sqrt{2x}$

a) $y = \underline{3} \sqrt{\underline{-}(x - \underline{1})}$

$a=3 \rightarrow$ A vertical stretch about the x-axis by a factor of 3.

$b=-1 \rightarrow$ No horizontal stretch about the y-axis and a reflection in the y-axis.

$h=1 \rightarrow$ translated 1 unit right.

$k=0 \rightarrow$ No vertical translation.


$$y = \sqrt{x}$$

x	y
0	0
1	1
4	2
9	3
16	4
25	5

$$(x,y) \rightarrow \left[\frac{1}{-1} x + 1, 3y + 0 \right]$$

$$(x,y) \rightarrow (-x+1, 3y)$$

x	y
1	0
0	3
-3	6
-8	9
-15	12
-24	15



D: $\{x | x \leq 1, x \in \mathbb{R}\}$ ($b = -1$)

$(-\infty, 1]$

R: $\{y | y \geq 0, y \in \mathbb{R}\}$ ($a = 3$)

$[0, \infty)$

b) $y - 3 = -\sqrt{2x}$

$$\begin{array}{rcl} y & = & -\sqrt{2x} + 3 \\ & = & \end{array}$$

$a = -1 \rightarrow$ Vertically reflected in the x-axis

$b = 2 \rightarrow$ horizontally stretched about the y-axis by a factor of $\frac{1}{2}$.

$h=0 \rightarrow$ no horizontal translation.


$k=3 \rightarrow$ vertically translated 3 units up.

$$y = \sqrt{x}$$

$$(x, y) \rightarrow \left[\frac{1}{2}x, -y + 3 \right]$$

x	y
0	0
1	1
4	2
9	3
16	4
25	5

x	y
0	3
$\frac{1}{2}$	2
$\frac{1}{4}$	1
$\frac{1}{9}$	0
$\frac{1}{16}$	-1
$\frac{1}{25}$	-2



D: $\{x | x \geq 0, x \in \mathbb{R}\}$ ($b=2$)
 $[0, \infty)$

R: $\{y | y \leq 3, y \in \mathbb{R}\}$ ($a=-1$)
 $(-\infty, 3]$


Homework

#2-5 on page 72-73


assignment

$$\begin{aligned}y - 4 &= -3\sqrt{-x+2} \\y &= -3\sqrt{-x+2} + 4 \\y &= -3\sqrt{-1(\underline{x}-2)} + 4\end{aligned}$$


**2.1 Radical Functions and Transformations,
pages 72 to 77**




2. a) $a = 7 \rightarrow$ vertical stretch by a factor of 7
 $h = 9 \rightarrow$ horizontal translation 9 units right
 domain $\{x \mid x \geq 9, x \in \mathbb{R}\}$, range $\{y \mid y \geq 0, y \in \mathbb{R}\}$
- b) $b = -1 \rightarrow$ reflected in y -axis
 $k = 8 \rightarrow$ vertical translation up 8 units
 domain $\{x \mid x \leq 0, x \in \mathbb{R}\}$, range $\{y \mid y \geq 8, y \in \mathbb{R}\}$
- c) $a = -1 \rightarrow$ reflected in x -axis
 $b = \frac{1}{5} \rightarrow$ horizontal stretch factor of 5
 domain $\{x \mid x \geq 0, x \in \mathbb{R}\}$, range $\{y \mid y \leq 0, y \in \mathbb{R}\}$
- d) $a = \frac{1}{3} \rightarrow$ vertical stretch factor of $\frac{1}{3}$
 $h = -6 \rightarrow$ horizontal translation 6 units left
 $k = -4 \rightarrow$ vertical translation 4 units down
 domain $\{x \mid x \geq -6, x \in \mathbb{R}\}$,
 range $\{y \mid y \geq -4, y \in \mathbb{R}\}$
3. a) B b) A c) D d) C
4. a) $y = 4\sqrt{x + 6}$ b) $y = \sqrt{8x - 5}$
 c) $y = \sqrt{-(x - 4)} + 11$ or $y = \sqrt{-x + 4} + 11$
 d) $y = -0.25\sqrt{0.1x}$ or $y = -\frac{1}{4}\sqrt{\frac{1}{10}x}$




domain
 $\{x \mid x \leq 0, x \in \mathbb{R}\}$,
range
 $\{y \mid y \geq -3, y \in \mathbb{R}\}$




domain
 $\{x \mid x \geq -1, x \in \mathbb{R}\}$,
range
 $\{y \mid y \geq 0, y \in \mathbb{R}\}$




domain
 $\{x \mid x \geq 2, x \in \mathbb{R}\}$,
range
 $\{y \mid y \leq 0, y \in \mathbb{R}\}$



domain
 $\{x \mid x \leq 2, x \in \mathbb{R}\}$,
range
 $\{y \mid y \leq 1, y \in \mathbb{R}\}$



domain
 $\{x \mid x \geq 0, x \in \mathbb{R}\}$,
range
 $\{y \mid y \geq 4, y \in \mathbb{R}\}$



domain
 $\{x \mid x \leq -2, x \in \mathbb{R}\}$,
range
 $\{y \mid y \geq -1, y \in \mathbb{R}\}$