


Sketch the following and determine a negative angle co-terminal with:

$$(i) \frac{5881\pi}{3}$$

$$\frac{5880\pi}{3}, \frac{5881\pi}{3}, \frac{5882\pi}{3}$$

$$1960\pi$$



Negative coterminal angle:

$$\frac{5881\pi}{3} - \frac{1960\pi}{1}$$


$$\frac{5881\pi}{3} - \frac{5886\pi}{3}$$

$$\frac{-5\pi}{3}$$

Reference Triangles:

Definition 17 The reference angle $\bar{\theta}$ of an angle θ in standard position is the acute angle (between 0 and 90°) the terminal side makes with the x-axis.

The picture below illustrates this concept.




What is the significance of reference angles?

Angles on the Cartesian Plane

$<90^\circ$ or $<\frac{\pi}{2}$ or $<1.57\text{rads}$

- **Reference Angle** - an acute angle formed between the terminal arm and the x-axis.
- **Reference Triangle** - a triangle formed by drawing a perpendicular line from a point on the terminal to the x-axis.




Notice what will happen if the rotation moves into other quadrants?

TRIG RATIOS on the CARTESIAN PLANE	
$\sin \theta = \frac{y}{r} = \frac{\alpha}{h}$	$\csc \theta = \frac{r}{y} = \frac{h}{\alpha}$
$\cos \theta = \frac{x}{r} = \frac{\alpha}{h}$	$\sec \theta = \frac{r}{x} = \frac{h}{\alpha}$
* $\tan \theta = \frac{y}{x} = \frac{\alpha}{\alpha}$	$\cot \theta = \frac{x}{y} = \frac{\alpha}{\alpha}$
"Primary"	"Reciprocal"

TRIG RATIOS IN ALL 4 QUADRANTS

What primary trig ratios are POSITIVE in...



Where is θ if...

$$\csc \theta < 0$$

$$\sin\theta < 0 \text{ & } \tan\theta < 0$$

$$\csc \theta > 0 \text{ & } \cot \theta < 0$$

Homework


If $\underline{\sec \theta = -\sqrt{10}}$ and $\underline{\sin \theta > 0}$, determine the value of $\csc \theta$

$$\sec \theta = -\frac{\sqrt{10}}{1} = \frac{\text{hyp}}{\text{adj}} = \frac{r}{x}$$

$$r = \sqrt{10} \quad (\text{Always } +)$$

$$x = -1$$

- ① Determine what quadrant:
 $\sec \theta < 0 \Rightarrow \sin \theta > 0$
or $\cos \theta < 0 \Rightarrow \sin \theta > 0$




- ④ Find $\csc \theta$:

$$\csc \theta = \frac{r}{y}$$

$$\boxed{\csc \theta = \frac{\sqrt{10}}{3}}$$

- ② Draw a diagram




$$\begin{aligned} ③ \quad x^2 + y^2 &= r^2 \\ (-1)^2 + y^2 &= (\sqrt{10})^2 \\ 1 + y^2 &= 10 \\ y^2 &= 9 \\ y &= \pm 3 \\ y &= 3 \quad (\text{Q2}) \end{aligned}$$

Determine the measure (in radians) of an angle whose terminal arm passes through the ordered pair $(-2\sqrt{3}, -4)$

$$x = -2\sqrt{3}$$

$$y = -4$$



- ① Draw a diagram

- ② Find $\bar{\theta}$

$$\tan \bar{\theta} = \frac{y}{x}$$

$$\tan \bar{\theta} = \frac{-4}{-2\sqrt{3}} = \frac{2}{\sqrt{3}}$$

$$\tan \bar{\theta} = 1.1547$$

$$\bar{\theta} = \tan^{-1}(1.1547)$$

convert calculator
to radians

$$\boxed{\bar{\theta} = 0.86 \text{ rads}}$$

- ③ Find θ



$$\theta = \pi + \bar{\theta}$$

$$\theta = 3.14 + 0.86$$



$$\boxed{\theta = 4 \text{ rads}}$$

$\theta = \pi - \bar{\theta}$	$\theta = \bar{\theta}$
$\theta = 180^\circ - \bar{\theta}$	$\theta = 360^\circ - \bar{\theta}$
$\theta = 180^\circ + \bar{\theta}$	$\theta = 2\pi - \bar{\theta}$
$\theta = \pi + \bar{\theta}$	




In Degrees



In Radians



Special Angles (in radians)



(Multiples of 90°)

- The Unit Circle
- Center is @ $(0,0)$
 - radius is 1 unit


Solving Trig Expressions by Sketching Angles

Ex. Evaluate the $\sin 690^\circ$

Optional

(i) Find principal angle:

$$\theta = 690^\circ - 360^\circ = 330^\circ$$



(ii) Sketch (θ)

(iii) Find $\bar{\theta}$

$$\bar{\theta} = 360^\circ - 330^\circ = 30^\circ$$

(iv) Label


(v) Evaluate $\sin 690^\circ = \frac{-1}{2}$

$$\text{Ex. } \cos \frac{13\pi}{3} = \frac{x}{r} = \frac{\frac{1}{2}}{2} = \frac{1}{4}$$

$\frac{12\pi}{3}, \frac{13\pi}{3}, \frac{14\pi}{3}$

4π

Even



$$\theta = \frac{13\pi}{3} - \frac{12\pi}{3}$$


$$\bar{\theta} = \frac{\pi}{3}$$

Homework

Evaluate each Trig Expression (provide a sketch of each angle)

$$1. \tan \frac{17\pi}{6} = -\frac{1}{\sqrt{3}} \quad 2. \sin \frac{15\pi}{4} = -\frac{1}{\sqrt{2}} \quad 3. \cos\left(-\frac{21\pi}{4}\right) = -\frac{1}{\sqrt{2}}$$


① $\frac{16\pi}{6}, \boxed{\frac{17\pi}{6}}, \frac{18\pi}{6}$
 $\frac{3\pi}{6} (0\text{st})$



$$\tan \frac{17\pi}{6} = \frac{y}{x} = \frac{1}{\sqrt{3}} = -\frac{1}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}}$$

$$\boxed{\tan \frac{17\pi}{6} = -\frac{\sqrt{3}}{3}}$$


② $\frac{14\pi}{4}, \boxed{\frac{15\pi}{4}}, \frac{16\pi}{4}$
 $\frac{4\pi}{4} (\text{Even})$



$$\sin \frac{15\pi}{4} = \frac{y}{r} = \frac{-1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}}$$

$$\boxed{\sin \frac{15\pi}{4} = -\frac{\sqrt{2}}{2}}$$

③ $-\frac{21\pi}{4} + \frac{6\pi}{1}$
 $-\frac{21\pi}{4} + \frac{24\pi}{4}$
 $\frac{3\pi}{4}$




$$\cos\left(-\frac{21\pi}{4}\right) = \cos \frac{3\pi}{4} = \frac{x}{r} = \frac{-1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}}$$

$$\boxed{\cos\left(-\frac{21\pi}{4}\right) = -\frac{\sqrt{2}}{2}}$$

$\frac{2\pi}{4}, \boxed{\frac{3\pi}{4}}, \frac{4\pi}{4}$
 $\frac{\pi}{4} (\text{odd})$

Unit Circle



unit circle

- a circle with radius 1 unit ($r = 1$)
- a circle of radius 1 unit with centre at the origin on the Cartesian plane is known as *the unit circle*

$$\sin \theta = \frac{y}{r} = \underline{y} = y \rightarrow \text{Ex: } \sin 90^\circ = 1$$


$$\cos \theta = \frac{x}{r} = \underline{x} = x \rightarrow \text{Ex: } \cos \pi = -1$$

$$\tan \theta = \frac{y}{x} \rightarrow \text{Ex: } \tan 270^\circ = \frac{-1}{0} = \text{undefined}$$

$$\csc \theta = \frac{1}{y} \rightarrow \text{Ex: } \csc 360^\circ = \frac{1}{0} = \text{undefined}$$

$$\sec \theta = \frac{1}{x} \rightarrow \text{Ex: } \sec 5\pi = \frac{1}{-1} = -1$$

$$\cot \theta = \frac{x}{y} \rightarrow \text{Ex: } \cot \frac{3\pi}{2} = \frac{0}{-1} = 0$$



$$\begin{aligned}x^2 + y^2 &= r^2 \\x^2 + y^2 &= (1)^2\end{aligned}$$

The equation of the unit circle is $x^2 + y^2 = 1$.

$$\underline{r=1}$$

Determine the equation of a circle with centre at the origin and radius 6.

$$\underline{r=6}$$


$$\begin{aligned}x^2 + y^2 &= r^2 \\x^2 + y^2 &= (6)^2 \\x^2 + y^2 &= 36\end{aligned}$$

Problems Involving the Unit Circle:

Determine Coordinates for Points of the Unit Circle

Determine the coordinates for all points on the unit circle that satisfy the conditions given. Draw a diagram in each case.

- the y -coordinate is $-\frac{1}{\sqrt{2}}$ and the point is in quadrant III



Find x

$$\begin{aligned}x^2 + y^2 &= r^2 \\x^2 + \left(-\frac{1}{\sqrt{2}}\right)^2 &= (1)^2 \\x^2 + \frac{1}{2} &= 1\end{aligned}$$

$$x^2 = 1 - \frac{1}{2}$$

$$x^2 = \frac{1}{2}$$

$$x = \pm \frac{1}{\sqrt{2}}$$

$$x = \pm \frac{\sqrt{2}}{2}$$

$$x = -\frac{1}{\sqrt{2}}$$


$$x = -\frac{1}{\sqrt{2}} \quad (\text{Q3})$$

coordinates are $\left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$


Problems Involving the Unit Circle:

If $P(150^\circ)$ is the point at which the terminal arm of an angle θ in standard position intersects the unit circle, determine the exact coordinates of...

$$r=1$$




Scale the diagram so that $r=1$ (unit circle)




coordinates are $\left(\frac{-\sqrt{3}}{2}, \frac{1}{2}\right)$

Special Angles on the Unit Circle:




Let's use $\frac{\pi}{4}$ as our reference angle

Construct reference triangles
for all multiples of $\pi/4$
between 0 and 2π




Unit Circle of Special Angles in Degrees



This is lovely...so what is it used for????

Unit Circle of Special Angles in Radians




Questions from Homework

① c)


$$\frac{10\pi}{6}, \frac{11\pi}{6}, \frac{12\pi}{6}$$

$\frac{2\pi}{6}$



$$\bar{\theta} = \frac{12\pi}{6} - \frac{11\pi}{6}$$

$$\bar{\theta} = \frac{\pi}{6}$$



Coordinates are:
 $(\frac{\sqrt{3}}{2}, -\frac{1}{2})$

Questions from Homework

- ③ If $\csc \theta = -\frac{\sqrt{10}}{2}$ and $\tan \theta > 0$ determine the value of the 5 remaining trig ratios as radicals in simplest form.

Given:

$$\csc \theta = -\frac{\sqrt{10}}{2} = \frac{h}{o} = \frac{r}{y}$$

$$r = \sqrt{10} \quad (r \text{ is always positive})$$

$$y = -2$$

- ④ Find the 5 trig ratios

$$\sin \theta = -\frac{2}{\sqrt{10}} = -\frac{2\sqrt{10}}{10} = -\frac{\sqrt{10}}{5}$$

$$\cos \theta = -\frac{\sqrt{6}}{\sqrt{10}} = -\frac{\sqrt{60}}{10} = -\frac{2\sqrt{15}}{10} = -\frac{\sqrt{15}}{5}$$

$$\tan \theta = -\frac{2}{-\sqrt{6}} = \frac{2\sqrt{6}}{6} = \frac{\sqrt{6}}{3}$$


$$\sec \theta = \frac{\sqrt{10}}{\sqrt{6}} = -\frac{\sqrt{60}}{6} = -\frac{2\sqrt{15}}{6} = -\frac{\sqrt{15}}{3}$$

$$\cot \theta = -\frac{\sqrt{6}}{-2} = \frac{\sqrt{6}}{2}$$

① Determine what quadrant
 $\csc \theta < 0 + \tan \theta > 0$

Quad 3

- ② Draw a diagram:



- ③ Find the missing side:

$$x^2 + y^2 = r^2$$

$$x^2 + (-2)^2 = (\sqrt{10})^2$$

$$x^2 + 4 = 10$$


$$x^2 = 6$$

$$x = \pm \sqrt{6}$$

$$x = -\sqrt{6} \quad (\text{Quad 3})$$

Questions from Homework


$$\textcircled{4} \quad b) \quad \sec \frac{2\pi}{3} = \frac{2}{1} = -2$$



Evaluate without the use of a calculator:


$$\sin \frac{9\pi}{2} - \cos^2 \left(\frac{29\pi}{6} \right) \tan \left(\frac{15\pi}{4} \right)$$

① $\frac{3\pi}{2}$, $\frac{9\pi}{2}$, $\frac{10\pi}{2}$




$$\sin \frac{9\pi}{2} = 1$$

② $\frac{2\pi}{6}$, $\frac{29\pi}{6}$, $\frac{30\pi}{6}$



$$\cos \frac{29\pi}{6} = -\frac{\sqrt{3}}{2}$$

③ $\frac{14\pi}{4}$, $\frac{15\pi}{4}$, $\frac{16\pi}{4}$



$$\tan \frac{15\pi}{4} = -1$$

$$\sin \frac{9\pi}{2} - \cos^2 \left(\frac{29\pi}{6} \right) \tan \left(\frac{15\pi}{4} \right)$$

$$(1) - \left(-\frac{\sqrt{3}}{2} \right)^2 (-1)$$

$$1 - \frac{3}{4} (-1)$$

$$1 + \frac{3}{4}$$

$$\frac{4}{4} + \frac{3}{4}$$

$$\frac{7}{4}$$

Evaluate without the use of a calculator:

$$\cos\left(\frac{16\pi}{3}\right) \tan^2\left(\frac{23\pi}{6}\right) + \csc\left(\frac{11\pi}{2}\right) + \sin^2\left(\frac{27\pi}{4}\right)$$

(Q3) (Q4) *Border of
3 and 4* (Q2)

$$\left(-\frac{1}{2}\right) \left(-\frac{1}{\sqrt{3}}\right)^2 + \left(-\frac{1}{1}\right) + \left(+\frac{1}{\sqrt{2}}\right)^2$$

$$-\frac{1}{2} \left(\frac{1}{3}\right) + (-1) + \frac{1}{2}$$

$$-\frac{1}{6} - 1 + \frac{1}{2}$$

$$\frac{-1}{6} - \frac{6}{6} + \frac{3}{6}$$

$$\frac{-4}{6}$$

- $\frac{2}{3}$

Homework:



Worksheet - Sketching Angles in Radians.doc

Solutions...

$$1. -\frac{5}{3}$$

$$5. \frac{4+3\sqrt{3}}{6}$$

$$2. \frac{-\sqrt{6}}{3}$$

$$6. \frac{-10}{3}$$

$$3. -2-\sqrt{3}$$

$$7. 0$$

$$4. \frac{-5}{3}$$

$$8. \frac{3+3\sqrt{3}}{-2}$$

Attachments

Worksheet - Sketching Angles in Radians.doc