

$$y = af[b(x-h)] + k \quad \text{Ch.1}$$

$$y = a\sqrt{b(x-h)} + k \quad \text{Ch.2}$$

$$y = ac^{b(x-h)} + k \quad \text{Ch.7}$$

$$y = a \log_c [b(x-h)] + k \quad \text{Ch.8}$$

$$y = a \sin[b(x-h)] + k \quad \text{Ch.5}$$

Ch.2 (Radical Functions)

$$\begin{array}{ll} D: \{x | x \geq h, x \in \mathbb{R}\} \text{ if } b > 0 & R: \{y | y \geq k, y \in \mathbb{R}\} \text{ if } a > 0 \\ \{x | x \leq h, x \in \mathbb{R}\} \text{ if } b < 0 & \{y | y \leq k, y \in \mathbb{R}\} \text{ if } a < 0 \end{array}$$

Ch.7 (Exponential Functions)

$$\begin{array}{ll} D: \{x | x \in \mathbb{R}\} & R: \{y | y > k, y \in \mathbb{R}\} \text{ if } a > 0 \\ & \{y | y < k, y \in \mathbb{R}\} \text{ if } a < 0 \end{array}$$

$$\text{HA: } y = k$$

Ch.8 (Logarithmic Functions)

$$\begin{array}{ll} D: \{x | x > h, x \in \mathbb{R}\} \text{ if } b > 0 & R: \{y | y \in \mathbb{R}\} \\ \{x | x < h, x \in \mathbb{R}\} \text{ if } b < 0 & \end{array}$$

$$\text{VA: } x = h$$

Ch.5 (Sinusoidal Functions)

$$\begin{array}{ll} D: \{x | x \in \mathbb{R}\} & R: \{y | \min \leq y \leq \max, y \in \mathbb{R}\} \\ \text{or } \{\theta | \theta \in \mathbb{R}\} & \end{array}$$

Function Operations

To combine two functions, $f(x)$ and $g(x)$, add or subtract as follows:

Sum of Functions

$$h(x) = f(x) + g(x)$$

$$h(x) = (f + g)(x)$$

\uparrow
adding

Difference of Functions

$$h(x) = f(x) - g(x)$$

$$h(x) = (f - g)(x)$$

\uparrow
subtracting

polynomial functions.

- $y = 3x + 5$ \rightarrow D: $\{x | x \in \mathbb{R}\}$

- $y = x^2 - x - 6$ \curvearrowleft D: $\{x | x \in \mathbb{R}\}$

- $y = x^3 + 7x^2 - 5x + 6$ \curvearrowleft D: $\{x | x \in \mathbb{R}\}$

- $y = x^4 - 2x^3 - 1$ \curvearrowleft D: $\{x | x \in \mathbb{R}\}$

Radical Function:

- $f(x) = \sqrt{x+3}$ \curvearrowleft $x+3 \geq 0$
 $x \geq -3$ D: $\{x | x \geq -3, x \in \mathbb{R}\}$

- $y = \sqrt{2x-5}$ \curvearrowleft $2x-5 \geq 0$
 $2x \geq 5$
 $x \geq \frac{5}{2}$ D: $\{x | x \geq \frac{5}{2}, x \in \mathbb{R}\}$

- $g(x) = \sqrt{4-x}$ \curvearrowleft $4-x \geq 0$
 $-x \geq -4$
 $x \leq 4$ D: $\{x | x \leq 4, x \in \mathbb{R}\}$
 $4-x \geq 0$
 $4 \geq x$

exponential function:

- $y = 3^x$ \curvearrowleft D: $\{x | x \in \mathbb{R}\}$

- $y = 3(2)^{x+5} - 1$ \curvearrowleft D: $\{x | x \in \mathbb{R}\}$

logarithmic function:

$$y = \log_3 x$$

\downarrow

D: $\{x | x > 0, x \in \mathbb{R}\}$

Rational function

$$y = \frac{x+1}{x-3}$$

$x-3 \neq 0$
 $x \neq 3$

D: $\{x | x \neq 3, x \in \mathbb{R}\}$


Key Ideas

- You can add two functions, $f(x)$ and $g(x)$, to form the combined function $h(x) = (f + g)(x)$.
- You can subtract two functions, $f(x)$ and $g(x)$, to form the combined function $h(x) = (f - g)(x)$.

 The domain of the combined function formed by the sum or difference of two functions is the domain common to the individual functions. For example,

- Domain of $f(x)$: $\{x \mid x \leq 3, x \in \mathbb{R}\}$
- Domain of $g(x)$: $\{x \mid x \geq -5, x \in \mathbb{R}\}$
- Domain of $h(x)$: $\{x \mid -5 \leq x \leq 3, x \in \mathbb{R}\}$

- The range of a combined function can be determined using its graph.
- To sketch the graph of a sum or difference of two functions given their graphs, add or subtract the y -coordinates at each point.



Domain of $f(x)$ $(-\infty, 3]$

Domain of $g(x)$ $[-5, \infty)$

Domain of $h(x)$ $[-5, 3]$

Example 1**Determine the Sum of Two Functions**

Consider the functions $f(x) = 2x + 1$ and $g(x) = x^3$.


- Determine the equation of the function $h(x) = (f + g)(x)$. **(Add)**
- Sketch the graphs of $f(x)$, $g(x)$, and $h(x)$ on the same set of coordinate axes.
- State the domain and range of $h(x)$.
- Determine the values of $f(x)$, $g(x)$, and $h(x)$ when $x = 4$.

a)
$$\begin{aligned} h(x) &= (f+g)(x) \\ h(x) &= \underline{f(x)} + \underline{g(x)} \\ h(x) &= 2x+1 + x^3 \\ h(x) &= x^3 + 2x+1 \end{aligned}$$

Domain:	$f(x) = 2x+1$	\leftrightarrow	Domain: $g(x) = x^3$
$\{x x \in \mathbb{R}\}$			$\{x x \in \mathbb{R}\}$
$(-\infty, \infty)$			$(-\infty, \infty)$

b)
$$\begin{aligned} f(x) &= 2x+1 & g(x) &= x^3 & h(x) &= x^3 + 2x+1 \end{aligned}$$

$\begin{array}{ c c } \hline x & y \\ \hline -2 & -3 \\ -1 & -1 \\ 0 & 1 \\ 1 & 3 \\ 2 & 5 \\ \hline \end{array}$	$\begin{array}{ c c } \hline x & y \\ \hline -2 & -8 \\ -1 & -1 \\ 0 & 0 \\ 1 & 1 \\ 2 & 8 \\ \hline \end{array}$	$\begin{array}{ c c } \hline x & y \\ \hline -2 & -1 \\ -1 & 0 \\ 0 & 1 \\ 1 & 4 \\ 2 & 9 \\ \hline \end{array}$
---	---	--




How are the y -coordinates of points on the graph of $h(x)$ related to those on the graphs of $f(x)$ and $g(x)$?

c)
$$\boxed{h(x) = x^3 + 2x + 1}$$

 D: $\{x | x \in \mathbb{R}\}$ or $(-\infty, \infty)$
 R: $\{y | y \geq 0, y \in \mathbb{R}\}$ or $[0, \infty)$

d) When $x = 4$

$f(4) = 2(4) + 1$ $f(4) = 9$	$g(4) = (4)^3$ $g(4) = 64$	$h(4) = (4)^3 + 2(4) + 1$ $h(4) = 64 + 8 + 1$ $h(4) = 73$ or $9 + 64 = \underline{\underline{73}}$
---------------------------------	-------------------------------	---



How are the y -coordinates of points on the graph of $h(x)$ related to those on the graphs of $f(x)$ and $g(x)$?

- c) The function $f(x) = 2x + 1$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $g(x) = x^2$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $h(x) = (f + g)(x)$ has domain $\{x \mid x \in \mathbb{R}\}$, which consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$.
 The range of $h(x)$ is $\{y \mid y \geq 0, y \in \mathbb{R}\}$.

- d) Substitute $x = 4$ into $f(x)$, $g(x)$, and $h(x)$.

$$\begin{array}{lll} f(x) = 2x + 1 & g(x) = x^2 & h(x) = x^2 + 2x + 1 \\ f(4) = 2(4) + 1 & g(4) = 4^2 & h(4) = 4^2 + 2(4) \\ f(4) = 8 + 1 & g(4) = 16 & h(4) = 16 + 8 + 1 \\ f(4) = 9 & & h(4) = 25 \end{array}$$

Example 2**Determine the Difference of Two Functions**

Consider the functions $f(x) = \sqrt{x-1}$ and $g(x) = x-2$.

- Determine the equation of the function $h(x) = (f - g)(x)$. (subtract)
- Sketch the graphs of $f(x)$, $g(x)$, and $h(x)$ on the same set of coordinate axes.
- State the domain of $h(x)$.
- Use the graph to approximate the range of $h(x)$.

a)
$$\begin{aligned} h(x) &= (f - g)(x) \\ h(x) &= f(x) - g(x) \\ h(x) &= \sqrt{x-1} - (x-2) \\ h(x) &= \sqrt{x-1} - x + 2 \end{aligned}$$

Domain:
 $f(x) = \sqrt{x-1}$ (\curvearrowright)
 $x-1 \geq 0$
 $x \geq 1$
 $\{x | x \geq 1, x \in \mathbb{R}\}$
or $[1, \infty)$

Domain:
 $g(x) = x-2$ (\curvearrowleft)
 $\{x | x \in \mathbb{R}\}$
or $(-\infty, \infty)$


b) Method 1: Use Paper and Pencil

For the function $f(x) = \sqrt{x-1}$, the value of the radicand must be greater than or equal to zero: $x-1 \geq 0$ or $x \geq 1$.

x	$f(x) = \sqrt{x-1}$	$g(x) = x-2$	$h(x) = \sqrt{x-1} - x + 2$
-2	undefined	-4	undefined
-1	undefined	-3	undefined
0	undefined	-2	undefined
1	0	-1	1
2	1	0	1
5	2	3	-1
10	3	8	-5

Why is the function $h(x)$ undefined when $x < 1$?

How could you use the values in the columns for $f(x)$ and $g(x)$ to determine the values in the column for $h(x)$?




How could you use the y-coordinates of points on the graphs of $f(x)$ and $g(x)$ to create the graph of $h(x)$?

c)
$$h(x) = \sqrt{x-1} - x + 2$$

D: $\{x | x \geq 1, x \in \mathbb{R}\}$ or $[1, \infty)$


b)
$$h(x) = \sqrt{x-1} - x + 2$$

R: $\{y | y \leq 1.2, y \in \mathbb{R}\}$ or $(-\infty, 1.2]$ see graph



How could you use the y-coordinates of points on the graphs of $f(x)$ and $g(x)$ to create the graph of $h(x)$?

- c) The function $f(x) = \sqrt{x - 1}$ has domain $\{x \mid x \geq 1, x \in \mathbb{R}\}$.
 The function $g(x) = x - 2$ has domain $\{x \mid x \in \mathbb{R}\}$.
 The function $h(x) = (f - g)(x)$ has domain $\{x \mid x \geq 1, x \in \mathbb{R}\}$, which consists of all values that are in both the domain of $f(x)$ and the domain of $g(x)$.



What values of x belong to the domains of both $f(x)$ and $g(x)$?

- d) From the graph, the range of $h(x)$ appears to be approximately $\{y \mid y \leq 1.2, y \in \mathbb{R}\}$.




How can you use a graphing calculator to verify the range?

Homework



finish #1-11 on page 483-484

10.1 Sums and Differences of Functions, pages 483 to 487

1. a) $h(x) = |x - 3| + 4$ b) $h(x) = 2x - 3$
 c) $h(x) = 2x^2 + 3x + 2$ d) $h(x) = x^2 + 5x + 4$
2. a) $h(x) = 5x + 2$ b) $h(x) = -3x^2 - 4x + 9$
 c) $h(x) = -x^2 - 3x + 12$ d) $h(x) = \cos x - 4$
3. a) $h(x) = x^2 - 6x + 1$; $h(2) = -7$
 b) $m(x) = -x^2 - 6x + 1$; $m(1) = -6$
 c) $p(x) = x^2 + 6x - 1$; $p(1) = 6$
4. a) $y = 3x^2 + 2 + \sqrt{x+4}$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 b) $y = 4x - 2 - \sqrt{x+4}$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 c) $y = \sqrt{x+4} - 4x + 2$; domain $\{x \mid x \geq -4, x \in \mathbb{R}\}$
 d) $y = 3x^2 + 4x$; domain $\{x \mid x \in \mathbb{R}\}$



6. a) 8 b) 6 c) 7
 d) not in the domain
7. a) B b) C c) A



9. a) $y = 3x^2 + 11x + 1$
c) $y = 3x^2 + 3x + 1$

b) $y = 3x^2 - 3x + 3$
d) $y = 3x^2 - 11x + 3$

10. a) $g(x) = x^2$

b) $g(x) = \sqrt{x+7}$

c) $g(x) = -3x + 1$

d) $g(x) = 3x^2 - x - 4$

11. a) $g(x) = x^2 - 1$

b) $g(x) = -\sqrt{x-4}$

c) $g(x) = 8x - 9$

d) $g(x) = 2x^2 - 11x - 6$