1.5

Proofs That Are Not Valid

GOAL

Identify errors in proofs.

EXPLORE...

• Consider the following statement: There are tthree errorss in this sentence. Is the statement valid?

SAMPLE ANSWER

There are only two spelling errors in the statement, not three, so the statement is invalid. If the statement is invalid, however, the statement itself is an error, making a total of three errors in the statement. Because the statement contains three errors, it is valid. But a statement cannot be both valid and invalid.

APPLY the Math

EXAMPLE 1

Using reasoning to determine the validity of an argument

Athletes do not compete in both the Summer and Winter Olympics. Hayley Wickenheiser has represented Canada four times at the Winter Olympics. Therefore, Hayley Wickenheiser has not participated in the Summer Olympics.

Tia read these statements and knew that there was an error. Identify the error in the reasoning.



Hayley Wickenheiser has represented Canada four times at the Winter Olympics.

Tia's Solution

2010

in statement

Athletes do not compete in both the Summer and Winter Olympics.

Hayley Wickenheiser has represented Canada four times at the Winter Olympics.

Therefore, Hayley Wickenheiser has not participated in the Summer Olympics.

I did some research and found that 18 athletes have competed in both Games. This statement is not valid.

This statement is true. She has played on the national hockey team.

The conclusion was false because the first statement was false. Hayley played for Canada in the softball competition in the 2000 Summer Olympics.

EXAMPLE 1

Using reasoning to determine the validity of an argument

ecros in

Athletes do not compete in both the Summer and Winter Olympics. Hayley Wickenheiser has represented Canada four times at the Winter Olympics. Therefore, Hayley Wickenheiser has not participated in the Summer Olympics.

Tia read these statements and knew that there was an error. Identify the error in the reasoning.



Hayley Wickenheiser has represented Canada four times at the Winter Olympics.

Your Turn

Zack is a high school student. All high school students dislike cooking. Therefore, Zack dislikes cooking. Where is the error in the reasoning?

Communication | Tip

Stereotypes are generalizations based on culture, gender, religion, or race. There are always counterexamples to stereotypes, so conclusions based on stereotypes are not valid.

Answer

The error is in the second statement. Not all high school students dislike cooking.



EXAMPLE 2

Using reasoning to determine the validity of a proof

Bev claims he can prove that 3 = 4.

Bev's Proof

Suppose that: $\underline{a} + \underline{b} = \underline{c}$

This statement can be written as: 4a - 3a + 4b - 3b = 4c - 3cAfter reorganizing, it becomes: 4a + 4b - 4c = 3a + 3b - 3c

Using the distributive property, Dividing both sides by (a + b - c),

$$4(a + b - c) = 3(a + b - c)$$


Show that Bev has written an **invalid proof**.

Pru's Solution

Suppose that:

$$a + b = c$$

Bev's **premise** was made at the beginning of the proof. Since variables can be used to represent any numbers, this part of the proof is valid.



V

Bev substituted 4a - 3a for a since 4a - 3a = a. Bev substituted 4b - 3b for b since 4b - 3b = b.

Bev substituted 4c - 3c for c since 4c - 3c = c.

$$4a + 4b - 4c = 3a + 3b - 3c$$

I reorganized the equation and I came up with the same result that Bev did when he reorganized. Simplifying would take me back to the premise. This part of the proof is valid.

$$4(a + b - c) = 3(a + b - c)$$

Since each side of the equation has the same coefficient for all the terms, factoring both sides is a valid step.

$$\frac{4(a+b-c)}{(a+b-c)} = \frac{3(a+b-c)}{(a+b-c)}$$

This step appears to be valid, but when I looked at the divisor, I identified the flaw.

$$a+b=c$$

$$a+b-c=c-c$$

$$a+b-c=0$$
Alvide

When I rearranged the premise, I determined that the divisor equalled zero.

Dividing both sides of the equation by a + b - c is not valid. Division by zero is undefined.

invalid proof

A proof that contains an error in reasoning or that contains invalid assumptions.

premise

A statement assumed to be true.

Using reasoning to determine the validity of a proof

Bev claims he can prove that 3 = 4.

Bev's Proof

Suppose that: a + b = c

This statement can be written as: 4a - 3a + 4b - 3b = 4c - 3c

After reorganizing, it becomes: 4a + 4b - 4c = 3a + 3b - 3c

Using the distributive property, 4(a + b - c) = 3(a + b - c)

Dividing both sides by (a + b - c), 4 = 3

Show that Bev has written an **invalid proof**.

Your Turn

How could this type of false proof be used to suggest that 65 = 64?



Answer

Suppose that a + b = c.

The statement can be written as	65a - 64a + 65b - 64b = 65c - 64c
After reorganizing, it becomes	65a + 65b - 65c = 64a + 64b - 64c
Using the distributive property,	65(a+b-c) = 64(a+b-c)
Dividing both sides by $(a + b - c)$,	65 = 64



Using reasoning to determine the validity of a proof

Liz claims she has proved that -5 = 5.

Liz's Proof

I assumed that -5 = 5.

Then I squared both sides: $(-5)^2 = 5^2$

I got a true statement: 25 = 25

This means that my assumption, -5 = 5, must be correct.

Where is the error in Liz's proof?

Simon's Solution

perror in line 1

I assumed that -5 = 5.

Liz started off with the false assumption that the two numbers were equal.

Then I squared both sides: $(-5)^2 = 5^2$ I got a true statement: 25 = 25 Everything that comes after the false assumption doesn't matter because the reasoning is built on the false assumption.

Even though 25 = 25, the underlying premise is not true.

-5≠5 -----

Liz's conclusion is built on a false assumption, and the conclusion she reaches is the same as her assumption.

If an assumption is not true, ---then any argument that was built on the assumption is not valid. **Circular reasoning** has resulted from these steps. Starting with an error and then ending by saying that the error has been proved is arguing in a circle.

circular reasoning

An argument that is incorrect because it makes use of the conclusion to be proved.

Using reasoning to determine the validity of a proof

Liz claims she has proved that -5 = 5.

Liz's Proof

I assumed that -5 = 5.

Then I squared both sides: $(-5)^2 = 5^2$

I got a true statement: 25 = 25

This means that my assumption, -5 = 5, must be correct.

Where is the error in Liz's proof?

Your Turn

How is an error in a premise like a counterexample?



Answer

An error in a premise is like a counterexample because a single error invalidates the argument, just as a single counterexample makes a conjecture invalid.

EXAMPLE 4 Using reasoning to determine the validity of a proof

Hossai is trying to prove the following number trick:

Choose any number. Add 3. Double it. Add 4. Divide by 2. Take away the number you started with.

Each time Hossai tries the trick, she ends up with 5. Her proof, however, does not give the same result.

Hossai's Proof

n	Choose any number.
n + 3	Add 3.
2n + 6	Double it. 🗸
2n + 10	Add 4. 🗸
2n + 5	Divide by 2.
n + 5	Take away the number you started with.

X

Where is the error in Hossai's proof?

Sheri's Solution

$$\begin{array}{ccc}
1 & \longrightarrow & 5 \\
10 & \longrightarrow & 5
\end{array}$$
 inductive...

I tried the number trick twice, for the number 1 and the number 10. Both times, I ended up with 5. The math trick worked for Hossai and for me, so the error must be in Hossai's proof.

n

n + 3

2n + 6

2n + 10

2n + 5

I corrected the mistake:

$$\frac{2n+10}{2} = n+5$$

$$n+5-n=5$$

The variable n can represent any number. This step is valid.

Adding 3 to *n* is correctly represented.

Doubling a quantity is multiplying by 2. This step is valid. Its implification is correct as well.

Adding 4 to the expression is correctly represented, and the simplification is correct.

The entire expression should be divided by 2, not just the constant. This step is where the mistake occurred.

I completed Hossai's proof by subtracting *n*. I showed that the answer will be 5 for any number.

EXAMPLE 4 Using reasoning to determine the validity of a proof

Hossai is trying to prove the following number trick:

Choose any number. Add 3. Double it. Add 4. Divide by 2. Take away the number you started with.

Each time Hossai tries the trick, she ends up with 5. Her proof, however, does not give the same result.

Hossai's Proof

Choose any number.

n+3 Add 3. 2n+6 Double it. 2n+10 Add 4. 2n+5 Divide by 2.

n + 5 Take away the number you started with.

Where is the error in Hossai's proof?

Your Turn

Is there a number that will not work in Hossai's number trick? Explain.



Answer

Hossai's number trick will work for every number because the proof uses *n* as any number and results in the number 5.

proof which means no counterexamples exist

In Summary

Key Idea

 A single error in reasoning will break down the logical argument of a deductive proof. This will result in an invalid conclusion, or a conclusion that is not supported by the proof.

Need to Know

- Division by zero always creates an error in a proof, leading to an invalid conclusion.
- Circular reasoning must be avoided. Be careful not to assume a result that follows from what you are trying to prove.
- The reason you are writing a proof is so that others can read and understand it. After you write a proof, have someone else who has not seen your proof read it. If this person gets confused, your proof may need to be clarified.

Assignment: pages 42-44

Questions: 1, 2, 3, 5, 6ab, 7

SOLUTIONS => 1.5 Proofs That Are Not Valid

- 1. Determine the error in each example of deductive reasoning.
- a) All runners train on a daily basis. Gabriel is a runner. Therefore, Gabriel trains daily
 - ⇒ The statement "all runners train on a daily basis" is invalid.
- b) All squares have four right angles. Quadrilateral PQRS has four right angles. Therefore, PQRS is a square.
 - ⇒ The reasoning leading to the conclusion is invalid. Rectangles also have four right angles
- 2. According to this proof, Proof 5=7. Identify the error. 1=1+1

From => The first line 2(1) = 2(1+1)of the proof 2+3 = 4+3is invalid. 5 = 7

3.	Micker	y says he	can	prove	that	2=0.	Here
	is his	proof.					

Let both a and b be equal to 1.

$$a = b$$

$$a^2 = b^2 \checkmark$$

$$a^2 - b^2 = 0$$

$$(a-b)(a+b) = 0$$

$$\frac{(a-b)(a+b)}{(a-b)} = \frac{0}{(a-b)}$$

$$1(a+b)=0$$

$$a+b=0$$

$$1 + 1 = 0$$

$$2 = 0$$

Transitive property

Squaring both sides

Subtracting b^2 from both sides

Factoring a difference of squares

Dividing both sides by a - b

Simplifying

Substitution

Explain whether each statement in Mickey's proof is valid.

Line 5 is not valid since Mickey is dividing by (a-b) and a-b=0.

	ed a moth trick in which she nded with 4. When Ali tried to trick, however, it did not work.
Ali's Proof	
n	I used <i>n</i> to represent any number.
2n 🗸	Multiply by 2.
2n+8	Add 8.
2n+4 ×	Divide by 2.
n+4	Subtract your starting number.
	he error in Ali's proof, and explain asoning is incorrect.

6. Connie tried this number trick:
 Write down the number of your street address. Multiply by 2. Add the number of days in a week. Multiply by 50. Add your age. Subtract the number of days in a year. Add 15.
Connie's result was a number in which the tens and ones digits were her age and the rest of the digits were the number from her street address. She tried to prove why this works, but her final expression did not make sense.
a) Try this number trick to see if you get the same result as Connie. Street Address: 14 Age: 16 n. 14. x2 28 +7 35 x50 1750. +16 1766 -365 1401 +15 1416

Let n represent any house number.

2n

Multiply by 2.

2n + 7

Add the number of days in a week.

100n + 350

Multiply by 50.

Let a represent any age.

100n + 350 + a

Add your age.

100n + 350 + a - 360 Subtract the number of days in a year.

100n + a + 5

Add 15.

b) Determine the errors in her proof, and then correct them.

⇒ There is an error in line 6 of Connie's proof. She subtracted 360 instead of 365 for the number of days in the year.

Correction.

Line 6: 100n+350+a-365 Line 7: 100n+a-15+15 4 100n + a

```
7. According to this proof, 2=1. Determine
   the error in reasoning.
    Let a = b.
         a^2 = ab
                         Multiply by a.
      a^2 + a^2 = a^2 + ab
                          Add a^2.
         2a^2 = a^2 + ab
                          Simplify.
    2a^2 - 2ab = a^2 + ab - 2ab
                          Subtract 2ab.
    2a^2 - 2ab = a^2 - ab
                          Simplify.
    2(a^2 - ab) = 1(a^2 - ab)
                          Factor.
          2 = 1 \times
                          Divide by (a^2 - ab).
    Frror => In the final line, there is a
                division by O.
              Ly a2-ab
                = a(a-b)
                = a(0) (since a = b)
```

PM11-1s5.gsp

1s5e1 finalt.mp4

1s5e2 finalt.mp4

1s5e3 finalt.mp4

1s5e4 finalt.mp4

1s5e5 finalt.mp4