





UNIT 1

UNDERLYING STRUCTURE OF MATTER

THE ATOMIC NUCLEUS



DISCOVERY OF THE NUCLEUS





PRACTICE QUESTIONS


- See Learning Target Guide
- 

SUBATOMIC PARTICLES

Particle	Charge	Mass (g)	Location
Electron (e⁻)	-1	9.11 x 10⁻²⁸	Electron cloud
Proton (p⁺)	+1	1.67 x 10⁻²⁴	Nucleus
Neutron (n⁰)	0	1.67 x 10⁻²⁴	Nucleus

COMPLETE SYMBOLS

- Contain the symbol of the element, the mass number and the atomic number.



Information from Symbols

- Find each of these:

a) number of protons

b) number of
neutrons

c) number of
electrons


d) Atomic number

e) Mass Number



ISOTOPES

- Atoms that have the same number of protons, but a different number of neutrons. i.e. the same atomic number but different mass numbers.
- We can also put the mass number *after* the name of the element:
 - carbon-12
 - carbon-14
 - uranium-235



ATOMIC MASS

- How heavy is an atom of oxygen?
 - It depends, because there are different *kinds* of oxygen atoms.
- We are more concerned with the average atomic mass.
- This is based on the abundance (percentage) of each variety of that element in nature.
 - We don't use grams for this mass because the numbers would be too small.

MEASURING ATOMIC MASS

- Instead of grams, the unit we use is the Atomic Mass Unit (amu)
- It is defined as one-twelfth the mass of a carbon-12 atom.
- Each isotope has its own atomic mass, thus we determine the average from percent abundance.

ATOMIC MASSES

Atomic mass is the average of all the naturally occurring isotopes of that element.

Isotope	Symbol	Composition of the nucleus	% in nature
Carbon-12	^{12}C	6 protons 6 neutrons	98.89%
Carbon-13	^{13}C	6 protons 7 neutrons	1.11%
Carbon-14	^{14}C	6 protons 8 neutrons	<0.01%


Carbon = 12.011

CALCULATING ATOMIC MASS


- The two most abundant isotopes of carbon are carbon-12 (mass 12.00 amu) and carbon-13 (mass 13.00 amu). Their relative abundances are 98.9% and 1.10%, respectively. Calculate the atomic mass of carbon.

ANOTHER EXAMPLE

- Using the information below, calculate the approximate atomic mass of silicon.




92.2%



4.7%



3.1%



trace

APPROXIMATING RELATIVE ABUNDANCE


- Copper, Cu, forms naturally with 34 and 36 neutrons. If its average atomic mass is 63.546, calculate the relative abundance found naturally.



REVIEW QUESTIONS

- See Learning Target Guide
- 

ELECTRONS IN ATOMS – CHAPTER 5



SECTION 5.1


MODELS OF THE ATOM

- OBJECTIVES:

- Identify the inadequacies in the Rutherford atomic model.
- Identify the new proposal in the Bohr model of the atom.
- Describe the energies and positions of electrons according to the quantum mechanical model.
- Describe how the shapes of orbitals related to different sublevels differ.

ERNEST RUTHERFORD'S MODEL

- Discovered dense positive piece at the center of the atom- “nucleus”
- Electrons would surround and move around it, like planets around the sun
- Atom is mostly empty space
- It did not explain the *chemical properties* of the elements – a better description of the **electron behavior** was needed



NIELS BOHR'S MODEL

- Why don't the electrons fall into the nucleus?
- Move like planets around the sun.
 - In specific circular paths, or orbits, at different levels.
 - An amount of fixed energy separates one level from another.

THE BOHR MODEL OF THE ATOM




Niels Bohr

I pictured the electrons orbiting the nucleus much like planets orbiting the sun. However, electrons are found in **specific** circular paths around the nucleus, and can **jump** from *one level to another*.



QUANTUM ENERGY LEVELS



BOHR'S MODEL

- Energy level of an electron
 - analogous to the rungs of a ladder
- The electron cannot exist between energy levels, just like you can't stand between rungs on a ladder
- A quantum of energy is the amount of energy required to move an electron from one energy level to another

THE QUANTUM MECHANICAL MODEL

- Energy is “quantized” - It comes in chunks.
- A quantum is the amount of energy needed to move from one energy level to another.
- Since the energy of an atom is never “in between” there must be a quantum leap in energy.
- In 1926, **Erwin Schrodinger** derived an equation that described the energy and position of the electrons in an atom

SCHRODINGER'S WAVE EQUATION



Erwin Schrodinger

$$-\frac{\hbar^2}{8\pi^2m} \frac{d^2\psi}{dx^2} + V\psi = E\psi$$


Equation for the probability of a single electron being found along a single axis (x-axis)

THE QUANTUM MECHANICAL MODEL

- Things that are very small *behave differently* from things big enough to see.
- The *quantum mechanical model* is a mathematical solution
- It is not like anything you can see.


THE QUANTUM MECHANICAL MODEL

- Has energy levels for electrons.
- Orbits are not circular.
- It can only tell us the probability of finding an electron a certain distance from the nucleus.



THE QUANTUM MECHANICAL MODEL


- The atom is found inside a blurry “electron cloud”
- An area where there is a *chance* of finding an electron.
- Think of fan blades




ATOMIC ORBITALS

- Principal Quantum Number (n) = the energy level of the electron: 1, 2, 3, etc.
- Within each energy level, the complex math of Schrodinger's equation describes several shapes.
- These are called atomic orbitals - regions where there is a high probability of finding an electron.
- Sublevels- like theater seats arranged in sections: letters s, p, d, and f


ORBITAL SHAPES




s orbital




p_x orbital




p_y orbital




p_z orbital




d_{xy}




d_{xz}



d_{yz}




















$d_{x^2-y^2}$



d_{z^2}

F- ORBITALS






PRINCIPAL QUANTUM NUMBER

Generally symbolized by “n”, it denotes the shell (energy level) in which the electron is located.

Maximum number of electrons that can fit in an energy level is:

$$2n^2$$

How many e⁻ in level 2? 3?



SUMMARY

	# of shapes (orbitals)	Maximum electrons	Starts at energy level
s	1	2	1
p	3	6	2
d	5	10	3
f	7	14	4

BY ENERGY LEVEL

- First Energy Level

- Has only s orbital

- only **2** electrons

- $1s^2$

- Second Energy Level

- Has s and p orbitals available

- 2 in s, 6 in p

- $2s^2 2p^6$

- 8 total electrons

BY ENERGY LEVEL

- Third energy level
- Has s, p, and d orbitals
- 2 in s, 6 in p, and 10 in d
- $3s^2 3p^6 3d^{10}$
- 18 total electrons

- Fourth energy level
- Has s, p, d, and f orbitals
- 2 in s, 6 in p, 10 in d, and 14 in f
- $4s^2 4p^6 4d^{10} 4f^{14}$
- 32 total electrons



REVIEW QUESTIONS

Refer to Learning Target Guide



SECTION 5.2

ELECTRON ARRANGEMENT IN ATOMS

- OBJECTIVES:

- Describe how to write the *electron configuration* for an atom.
- Explain why the actual electron configurations for some elements *differ* from those predicted by the Aufbau principle.

BY ENERGY LEVEL

- The orbitals do not fill up in a neat order.
- The energy levels overlap
- Lowest energy fill first.


QUANTUM NUMBERS

Each electron in an atom has a unique set of 4 quantum numbers which describe it.

- 1) Principal quantum number
- 2) Angular momentum quantum number
- 3) Magnetic quantum number
- 4) Spin quantum number

ELECTRON CONFIGURATIONS...


- ...are the way electrons are arranged in various orbitals around the nuclei of atoms. *Three rules tell us how:*
 - 1) Aufbau principle - electrons enter the lowest energy first.
 - This causes difficulties because of the overlap of orbitals of different energies – follow the diagram!




aufbau diagram - page 133

Aufbau is German for "building up"


ORBITAL SHAPES




s orbital




p_x orbital




p_y orbital




p_z orbital




d_{xy}




d_{xz}



d_{yz}



$d_{x^2-y^2}$



d_{z^2}

n=1	1s²				2e ⁻
n=2	2s²	2p⁶			8e ⁻
n=3	3s²	3p⁶	3d¹⁰		18e ⁻
n=4	4s²	4p⁶	4d¹⁰	4f¹⁴	32e ⁻
n=5	5s²	5p⁶	5d¹⁰	5f¹⁴	...
n=6	6s²	6p⁶	6d¹⁰
n=7	7s²	7p⁶

↙ = filling order

RULE 2: PAULI EXCLUSION PRINCIPLE



Wolfgang Pauli


No two electrons in an atom can have the same four quantum numbers.


To show the different direction of spin, a pair in the same orbital is written as:




ELECTRON CONFIGURATIONS

- 3) Hund's Rule- When electrons occupy orbitals of equal energy, they don't pair up until they have to.
- Let's write the electron configuration for Phosphorus
 - We need to account for all 15 electrons in phosphorus







- The next electrons go into the 2s orbital
- only 11 more...



- The next electrons go into the 2p orbital
- only 5 more...



- The next electrons go into the 3s orbital
- only 3 more...



The background is a solid teal color with a subtle gradient. In the four corners, there are decorative white line-art patterns resembling circuit traces or neural network connections, with small circles at the end of the lines.

PRACTICE QUESTIONS

- Refer to Learning Target Guide

ORBITALS FILL IN AN ORDER

- Lowest energy to higher energy.
- Adding electrons can change the energy of the orbital. Full orbitals are the absolute best situation.
- However, half filled orbitals have a lower energy, and are next best
 - Makes them more stable.
 - Changes the filling order

WRITE THE ELECTRON CONFIGURATIONS FOR THESE ELEMENTS:

- Titanium - 22 electrons



- Vanadium - 23 electrons




- Chromium - 24 electrons



❖ **But this is not what happens!!**

CHROMIUM IS ACTUALLY:

- $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^5$

- Why?
- This gives us two **half filled orbitals** (the others are all still full)
- Half full is slightly lower in energy.
- The same principal applies to copper.

COPPER'S ELECTRON CONFIGURATION

- Copper has 29 electrons so we expect:
 $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^9$
- But the *actual configuration* is:
- $1s^2 2s^2 2p^6 3s^2 3p^6 \underline{4s^1 3d^{10}}$
- This change gives one more filled orbital and one that is half filled.
- Remember these exceptions: **d^4** , **d^9**

IRREGULAR CONFIGURATIONS OF CHROMIUM AND COPPER

Chromium steals a 4s electron to **make** its 3d sublevel **HALF FULL**

Copper steals a 4s electron to **FILL** its 3d sublevel

K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
$4s^1$	$4s^2$	$3d^1$	$3d^2$	$3d^3$	$4s^1 3d^5$	$3d^5$	$3d^6$	$3d^7$	$3d^8$	$4s^1 3d^{10}$	$3d^{10}$	$4p^1$	$4p^2$	$4p^3$	$4p^4$	$4p^5$	$4p^6$

ELECTRON CONFIGURATION IN GROUPS

- Noble gases
 - Elements in group 8A.
 - The highest energy levels are completely filled with electrons.
 - That leads to them being relatively inert.
- Representative Elements
 - Groups 1A – 7A.
 - Group number is the number of electrons in the highest energy level.

1	1	H Hydrogen 1.00794
3	2 1	Li Lithium 6.941
11	2 8 1	Na Sodium 22.98976928
19	2 8 8 1	K Potassium 39.0983
37	2 8 18 8 1	Rb Rubidium 85.4678
55	2 8 18 18 8 1	Cs Caesium 132.9054519
87	2 8 18 32 18 8 1	Fr Francium (223)

4	2 2	Be Beryllium 9.012182
12	2 8 2	Mg Magnesium 24.3050
20	2 8 8 2	Ca Calcium 40.078
38	2 8 18 8 2	Sr Strontium 87.62
56	2 8 18 18 8 2	Ba Barium 137.327
88	2 8 18 32 18 8 2	Ra Radium (226)

5	2 3	B Boron 10.811
13	2 8 3	Al Aluminium 26.9815386
31	2 8 18 3	Ga Gallium 69.723
49	2 8 18 18 3	In Indium 114.818
81	2 8 18 32 18 3	Tl Thallium 204.3833
113	2 8 18 32 18 3	Uut Ununtrium (284)

6	2 4	C Carbon 12.0107
14	2 8 4	Si Silicon 28.0855
32	2 8 18 4	Ge Germanium 72.64
50	2 8 18 18 4	Sn Tin 118.710
82	2 8 18 32 18 4	Pb Lead 207.2
114	2 8 18 32 18 4	Uuq Ununquadium (289)

7	2 5	N Nitrogen 14.0067
15	2 8 5	P Phosphorus 30.973762
33	2 8 18 5	As Arsenic 74.92160
51	2 8 18 18 5	Sb Antimony 121.760
83	2 8 18 32 18 5	Bi Bismuth 208.98040
115	2 8 18 32 18 5	Uup Ununpentium (288)

8	2 6	O Oxygen 15.9994
16	2 8 6	S Sulfur 32.065
34	2 8 18 6	Se Selenium 78.96
52	2 8 18 18 6	Te Tellurium 127.60
84	2 8 18 32 18 6	Po Polonium (208.9824)
116	2 8 18 32 18 6	Uuh Ununhexium (292)

9	2 7	F Fluorine 18.9984032
17	2 8 7	Cl Chlorine 35.453
35	2 8 18 7	Br Bromine 79.904
53	2 8 18 18 7	I Iodine 126.90447
85	2 8 18 32 18 7	At Astatine (209.9871)
117	2 8 18 32 18 7	Uus Ununseptium

2	2	He Helium 4.002602
10	2 8	Ne Neon 20.1797
18	2 8 8	Ar Argon 39.948
36	2 8 18 8	Kr Krypton 83.798
54	2 8 18 18 8	Xe Xenon 131.293
86	2 8 18 32 18 8	Rn Radon (222.0176)
118	2 8 18 32 18 8	Uuo Ununoctium (294)



REVIEW QUESTIONS

- Refer to Learning Target Guide
- 