

Questions from homework

$$\textcircled{3} \text{ a) } f(x) = 2x^3 - 3x^2 \quad -2 \leq x \leq 2$$

$$f'(x) = 6x^2 - 6x$$

$$f'(x) = 6x(x-1)$$

$$(V: x=0, 1)$$

$$f(0) = 0 \quad (0, 0)$$

$$f(1) = -1 \quad (1, -1)$$

$$f(-2) = -16 - 12 = -28 \quad (-2, -28) \text{ abs min}$$

$$f(2) = 16 - 12 = 4 \quad (2, 4) \text{ abs max}$$

Questions from homework

$$\textcircled{4} \quad g) \quad g(x) = x^2 + \frac{16}{x} \quad 1 \leq x \leq 4$$

$$g(x) = x^2 + 16x^{-1}$$

$$g'(x) = 2x - 16x^{-2}$$

$$g'(x) = 2x - \frac{16}{x^2}$$


$$g'(x) = \frac{2x^3 - 16}{x^2}$$

$$\begin{array}{l|l} \text{cv: } 2x^3 - 16 = 0 & x^2 = 0 \\ 2x^3 = 16 & x = 0 \\ x^3 = 8 & \\ x = 2 & \end{array} \quad \begin{array}{l} x = 0 \text{ is not in} \\ \text{the domain} \end{array}$$

$f(2) = (2)^2 + \frac{16}{2}$	$f(1) = (1)^2 + \frac{16}{1}$	$f(4) = (4)^2 + \frac{16}{4}$
$f(2) = 4 + 8$	$f(1) = 1 + 16$	$f(4) = 16 + 4$
$f(2) = 12$	$f(1) = 17$	$f(4) = 20$
$(2, 12)$ abs min	$(1, 17)$	$(4, 20)$ abs max

The First Derivative Test

If f has a local maximum or minimum at c , then c must be a critical value of f (Fermat's Theorem), but not all critical numbers give rise to a maximum or minimum. For instance, recall that 0 is a critical number of the function $y = x^3$ but this function has no maximum or minimum at a critical number.





$y = x^3$
 $y' = 3x^2$
 CV: $0 = 3x^2$
 $0 = x^2$
 $0 = x$

One way of solving this is suggested by the figure below.

If f is increasing to the left of a critical number c and decreasing to the right of c , then f has a local max at c .

If f is decreasing to the left of a critical number c and increasing to the right of c , then f has a local min at c .



The First Derivative Test

Let c be a critical number of a continuous function f .

1. If $f'(x)$ changes from positive to negative at c , then f has a local max at c .
2. If $f'(x)$ changes from negative to positive at c , then f has a local min at c .
3. If $f'(x)$ does not change signs at c , then f has no max or min at c .

Example 1

Find the local maximum and minimum values of
 $f(x) = x^3 - 3x + 1$


$$f'(x) = 3x^2 - 3$$

$$f'(x) = 3(x^2 - 1)$$

$$f'(x) = 3(x-1)(x+1)$$

$$0 = 3(x-1)(x+1)$$

$$\text{CV: } x-1=0 \mid x+1=0 \\ x=1 \mid x=-1$$



Increasing on $(-\infty, -1) + (1, \infty)$

Decreasing on $(-1, 1)$

max @ $x = -1$

$$f(x) = x^3 - 3x + 1$$

$$f(-1) = (-1)^3 - 3(-1) + 1$$

$$f(-1) = -1 + 3 + 1$$

$$f(-1) = 3$$

$$\boxed{(-1, 3)}$$

min @ $x = 1$

$$f(x) = x^3 - 3x + 1$$

$$f(1) = (1)^3 - 3(1) + 1$$

$$f(1) = 1 - 3 + 1$$

$$f(1) = -1$$

$$\boxed{(1, -1)}$$

/

Example 2

Find the local maximum and minimum values of $g(x) = x^4 - 4x^3 - 8x^2 - 1$. Use this information to sketch the graph of g .

$$g'(x) = 4x^3 - 12x^2 - 16x$$

$$g'(x) = 4x(x^2 - 3x - 4)$$

$$g'(x) = 4x(x-4)(x+1)$$

$$0 = 4x(x-4)(x+1)$$

$$\text{cv: } 4x=0 \mid x-4=0 \mid x+1=0$$

$$x=0 \mid x=4 \mid x=-1$$

$$x = -1, 0, 4$$

$$\text{min @ } x = -1$$

$$g(x) = x^4 - 4x^3 - 8x^2 - 1$$

$$g(-1) = (-1)^4 - 4(-1)^3 - 8(-1)^2 - 1$$

$$g(-1) = 1 + 4 - 8 - 1$$

$$g(-1) = -4$$

$$(-1, -4)$$


$$\text{max @ } x = 0$$

$$g(x) = x^4 - 4x^3 - 8x^2 - 1$$

$$g(0) = (0)^4 - 4(0)^3 - 8(0)^2 - 1$$

$$g(0) = -1$$

$$(0, -1)$$



Increasing on $(-1, 0)$ and $(4, \infty)$

Decreasing on $(-\infty, -1)$ and $(0, 4)$

$$\text{min @ } x = 4$$


$$g(x) = x^4 - 4x^3 - 8x^2 - 1$$

$$g(4) = (4)^4 - 4(4)^3 - 8(4)^2 - 1$$

$$g(4) = 256 - 256 - 128 - 1$$

$$g(4) = -129$$

$$(4, -129)$$



The First Derivative Test

(for absolute extreme values)

Let c be a critical number of a continuous function f .

1. If $f'(x)$ is positive for all $x < c$ and $f'(x)$ is negative for all $x > c$, then $f(c)$ is the absolute maximum value.
2. If $f'(x)$ is negative for all $x < c$ and $f'(x)$ is positive for all $x > c$, then $f(c)$ is the absolute minimum value.

Homework

Page 182 # 1-4