

Questions from Homework

$$\textcircled{17} \quad k) \quad f(x) = \frac{x^2 + 5x + 4}{x^2 + 6x + 8} = \frac{(x+1)\cancel{(x+4)}}{(x+2)\cancel{(x+4)}} = \frac{x+1}{x+2}$$

VA: $x+2=0$ Infinite discontinuity @ $x=-2$

$x = -2$ $\lim_{x \rightarrow -2^-} \frac{f(x)}{f(x)} = +\infty$ $\lim_{x \rightarrow -2^+} \frac{f(x)}{f(x)} = -\infty$

($x=-2, 1$) ($x=-1, 1$)

HA: $\lim_{x \rightarrow \infty} \frac{x^2 + 5x + 4}{x^2 + 6x + 8}$ $y = 1$

$$= \lim_{x \rightarrow \infty} \frac{\frac{x^2}{x^2} + \frac{5x}{x^2} + \frac{4}{x^2}}{\frac{x^2}{x^2} + \frac{6x}{x^2} + \frac{8}{x^2}}$$

$$= \lim_{x \rightarrow \infty} \frac{1 + \frac{5}{x} + \frac{4}{x^2}}{1 + \frac{6}{x} + \frac{8}{x^2}}$$

$$= \frac{1}{1}$$

$$= 1$$

Point of Discontinuity/Hole:

$$x+4=0$$

$x = -4$

removable discontinuity

$$f(x) = \frac{x+1}{x+2}$$

$$f(-4) = \frac{-4+1}{-4+2}$$

$$f(-4) = \frac{-3}{-2}$$

$$f(-4) = \frac{3}{2}$$

→ $(-4, \frac{3}{2})$ or $(-4, 1.5)$

Questions from Homework

$$\textcircled{11} \text{ b) } f(x) = \frac{x^2 + 2x - 3}{x^2 + 3x - 4} = \frac{\cancel{(x-1)}(x+3)}{\cancel{(x-1)}(x+4)} = \frac{x+3}{x+4}$$

HA: $\lim_{x \rightarrow \infty} \frac{x^2 + 2x - 3}{x^2 + 3x - 4} = \frac{1}{1}$

$$y = 1$$

VA: $x + 4 = 0$

$$x = -4$$

$$\lim_{x \rightarrow -4^-} \frac{(-)}{(-)} = +\infty$$

$(x = -4.1)$

$$\lim_{x \rightarrow -4^+} \frac{(-)}{(+)} = -\infty$$

$(x = -3.9)$

Hole: $x - 1 = 0$

$$x = 1$$

sub in $x = 1$

$$f(1) = \frac{1+3}{1+4} = \frac{4}{5} = 0.8$$

$$(1, 0.8)$$

Questions from Homework

$$\textcircled{17} \text{ j) } f(x) = \frac{6x^3 - 30x^2 - 84x}{2x^3 + 3x^2 + x} = \frac{6x(x^2 - 5x - 14)}{x(\underline{2x+1})(x+2)} = \frac{6x(x-7)(x+2)}{x(\underline{x+1})(x+2)}$$

$$= \frac{6x(x-7)(x+2)}{x(2x+1)(x+1)}$$

① HA:

$$\lim_{x \rightarrow \infty} \frac{6x^3 - 30x^2 - 84x}{2x^3 + 3x^2 + x} = \frac{6}{2} = 3$$

$$y = 3$$

② VA:

$$(2x+1)(x+1) = 0$$

$$2x+1=0$$

$$2x = -1$$

$$x = -\frac{1}{2}$$

$$x+1=0$$

$$x = -1$$

③ Point of Discontinuity (Hole)

$$x = 0$$

$$\frac{6(-7)(2)}{(1)(1)} = -84$$

$$(0, -84)$$

Questions from Homework

$$\textcircled{17} \text{ c) } f(x) = \frac{x^3 + 3x^2 - 10x}{x^2 + 9x + 8} = \frac{x(x^2 + 3x - 10)}{(x+1)(x+8)} = \frac{x(x-5)(x+2)}{(x+1)(x+8)}$$

VA:

$$(x+1)(x+8) = 0$$

$$\begin{array}{l|l} x+1=0 & x+8=0 \\ \hline \boxed{x=-1} & \boxed{x=-8} \end{array}$$

HA:

$$\lim_{x \rightarrow \infty} \frac{x^3 + 3x^2 - 10x}{x^2 + 9x + 8} = \text{DNE}$$

There is no HA.

Holes:

None

Curve Sketching

In this chapter we look at further aspects of curves such as vertical and horizontal asymptotes, concavity, and inflections points. Then we use them, together with intervals of increase and decrease and maximum and minimum values, to develop a procedure for curve sketching.

Slant Asymptotes

For rational functions, slant asymptotes occur when the degree of the numerator is one more than the degree of the denominator and can be found by division. (Ignore the remainder)

- Similar to horizontal asymptotes
- you can cut through a slant asymptote.

Example

Find the slant asymptote of the curve $y = \frac{2x^3 - 3x^2 + x - 3}{x^2 + 1}$


$$\begin{array}{r}
 \underline{x^2 + 1} \overline{) 2x^3 - 3x^2 + x - 3} \\
 \underline{-(2x^3 + 2x)} \\
 -3x^2 - x - 3 \\
 \underline{-(-3x^2 - 3)} \\
 -x
 \end{array}$$

SA: $y = 2x - 3$

$m = \frac{2}{1}$

$b = -3$
(y-int)

-x
remainder



when $x = 7$

$$y = \frac{543}{50}$$

$$y = 10.86$$

$$2x^3 - 3x^2 + x - 3 - (2x^3 + 2x)$$
$$\cancel{2x^3} - 3x^2 + \underline{x} - 3 - \cancel{2x^3} - \underline{2x}$$
$$-3x^2 - \underline{x} - 3$$

Example

Find the slant asymptote of the curve


$$\begin{array}{r} \underline{x-1} \overline{\overset{-x}{-x^2+x+1}} \\ \underline{-(-x^2+x)} \\ \hline \text{remainder} \end{array}$$

$$y = \frac{1+x-x^2}{x-1}$$

$$\text{SA: } y = -x$$

$$m = -\frac{1}{1}$$

$$b = 0$$



$$\textcircled{a} \text{ VA: } x-1=0$$

$$\boxed{x=1}$$

$$\lim_{x \rightarrow 1^-} \frac{(+)}{(-)} = -\infty$$

(x=0.9)

$$\lim_{x \rightarrow 1^+} \frac{(+)}{(+)} = +\infty$$

(x=1.1)

Homework