


Questions from Homework

8. If seafood is not kept frozen (below 0 °C), it will spoil due to bacterial growth. The relative rate of spoilage increases with temperature according to the model $R = 100(2.7)^{\frac{T}{5}}$, where T is the temperature, in degrees Celsius, and R is the relative spoilage rate.

- Sketch a graph of the relative spoilage rate R versus the temperature T from 0 °C to 25 °C.
- Use your graph to predict the temperature at which the relative spoilage rate doubles to 200.
- What is the relative spoilage rate at 15 °C?
- If the maximum acceptable relative spoilage rate is 500, what is the maximum storage temperature?



Assignment:

y-int ($x=0$)

$$y = 4^{-2(x+5)} - 3$$

$$y = 4^{-2(0+5)} - 3$$

$$y = 4^{-2(5)} - 3$$

$$y = 4^{-10} - 3$$

$$y = \left(\frac{1}{4}\right)^{10} - 3$$

$$y = \frac{1}{1048576} - \frac{3}{1}$$

$$y = \frac{1}{1048576} - \frac{3145728}{1048576}$$

$$y = \frac{-3145727}{1048576}$$

$$y = -2.\bar{9}$$

$$(0, -2.\bar{9})$$

x-int ($y=0$)

$$y = 4^{-2(x+5)} - 3$$

$$0 = 4^{-2(x+5)} - 3$$

$$3 = 4^{-2(x+5)}$$

$$\sqrt[0.7925]{4} = 4^{-2(x+5)}$$

$$\frac{0.7925}{-2} = \frac{-2(x+5)}{-2}$$

$$-0.3962 = x+5$$

$$-5.3962 = x$$

$$(-5.4, 0)$$

9. A bacterial culture starts with 2000 bacteria and doubles every 0.75 h. After how many hours will the bacteria count be 32 000?

Given:

$$\text{Initial Amount} = 2000$$

$$\text{Base} = 2$$

$$\text{exp} = \frac{x}{0.75}$$

$$A = 32000$$

$$y = 2000(2)^{\frac{x}{0.75}}$$

$$A = 2000(2)^{\frac{t}{0.75}}$$

$$\frac{32000}{2000} = \frac{2000(2)^{\frac{t}{0.75}}}{2000}$$

$$\frac{\log 16}{\log 2} = 4 \quad 16 = 2^{\frac{t}{0.75}}$$

$$\cancel{2^4} = \cancel{2}^{\frac{t}{0.75}}$$

$$0.75 \quad 4 = \frac{t}{0.75} \cdot 0.75$$

$$\boxed{3h = t}$$

$$16 = 2^{t/0.75}$$

$$16 = (16)^{0.25 t/0.75}$$

$$(0.75) 1 = \frac{0.25t}{0.75} (0.75)$$

$$\frac{0.75}{0.25} = \frac{0.25t}{0.25}$$

$$3 = t$$

Given: $y = -3(2)^{2x+2} + 4$
 $y = -3(2)^{2(x+1)} + 4$


- i) state the parameters and describe the corresponding transformations
- ii) create a table to show what happens to the given points under each transformation
- iii) sketch the graph of the base function and the transformed function
- iv) describe the effects on the domain, range, equation of the horizontal asymptote, and intercepts

(i) $y = -3(2)^{2(x+1)} + 4$ $c = \text{base} = 2$

$a = -3 \rightarrow$ a vertical stretch by a factor of 3 and a reflection in the x axis
 $b = 2 \rightarrow$ a horizontal stretch by a factor of $\frac{1}{2}$
 $h = -1 \rightarrow$ 1 unit left
 $k = 4 \rightarrow$ 4 units up

(ii) $(x, y) \rightarrow (\frac{1}{2}x - 1, -3y + 4)$

$y = 2^x$			
x	y	x	y
-2	$\frac{1}{4}$	-2	$\frac{1}{4} = 3.25$
-1	$\frac{1}{2}$	-1.5	$\frac{1}{2} = 2.5$
0	1	-1	1
1	2	-0.5	2
2	4	0	8



(iv) D: $\{x | x \in \mathbb{R}\}$ or $(-\infty, \infty)$
 R: $\{y | y < 4, y \in \mathbb{R}\}$ or $(-\infty, 4)$
 HA: $y = 4$

x int ($y=0$)
 $y = -3(2)^{2(x+1)} + 4$
 $0 = -3(2)^{2(x+1)} + 4$

$-4 = -3(2)^{2(x+1)}$
 $\frac{-4}{-3} = \frac{-3(2)^{2(x+1)}}{-3}$
 $1.\bar{3} = (2)^{2(x+1)}$
 $(2)^{0.42} = (2)^{2(x+1)}$ $\frac{\log 1.\bar{3}}{\log 2} = 0.42$

$0.42 = 2(x+1)$

$0.21 = x+1$

$-0.79 = x$

$(-0.79, 0)$

check graph for these points

y int ($x=0$)
 $y = -3(2)^{2(x+1)} + 4$
 $y = -3(2)^{2(0+1)} + 4$

$y = -3(2)^2 + 4$

$y = -3(2)^2 + 4$

$y = -3(4) + 4$

$y = -12 + 4$

$y = -8$

$(0, -8)$

Ex: Exponential Equation

$$64^x = \left(\frac{1}{8}\right)^{x+1} (\sqrt{32})$$

$$64^x = \left(\frac{1}{8}\right)^{x+1} \cdot (32)^{\frac{1}{2}}$$

$$\frac{\log 64}{\log 2} = \underline{\underline{6}}$$

$$\frac{\log\left(\frac{1}{8}\right)}{\log 2} = -3$$

$$\frac{\log 32}{\log 2} = 5$$

$$(2^6)^x = (2^{-3})^{x+1} (2^5)^{\frac{1}{2}}$$

$$2^{6x} = 2^{-3x-3} \cdot 2^{\frac{5}{2}}$$

$$2^{6x} = 2^{-3x-3+\frac{5}{2}}$$

$$2^{6x} = 2^{-3x-\frac{6}{2}+\frac{5}{2}}$$

$$\cancel{2}^{6x} = \cancel{2}^{-3x-\frac{1}{2}}$$

$$6x = -3x - \frac{1}{2}$$

$$\frac{9x}{9} = -\frac{1}{2} \div 9$$

$$x = -\frac{1}{2} \cdot \frac{1}{9}$$

$$x = -\frac{1}{18}$$

Exponential Equation

$$81^{x+2} = \left(\frac{1}{27}\right)^{3x} \cdot (729)^{x-1}$$

$$\frac{\log\left(\frac{1}{27}\right)}{\log(3)} = -3$$

$$(3^4)^{x+2} = (3^{-3})^{3x} \cdot (3^9)^{x-1}$$

$$3^{4x+8} = 3^{-9x} \cdot 3^{6x-6}$$

$$3^{4x+8} = 3^{-9x+6x-6}$$

~~$$3^{4x+8} = 3^{-3x-6}$$~~

$$4x+8 = -3x-6$$

$$4x+3x = -6-8$$

$$\frac{7x}{7} = \frac{-14}{7}$$

$$x = -2$$

Base 2: a) $\left(\frac{1}{4}\right)$ and $\sqrt{64}$

$\left(\frac{1}{4}\right)$ and 8

$$\frac{\log\left(\frac{1}{4}\right)}{\log 2} = -2 \quad \left(2\right)^{-2} \quad \text{and} \quad \left(2\right)^3 \quad \frac{\log 8}{\log 2} = 3$$

$$\begin{aligned} &\sqrt{64} \\ &(\underline{64})^{\frac{1}{2}} \\ &(\underline{2^6})^{\frac{1}{2}} \\ &2^3 \end{aligned}$$

$$\begin{aligned} &\underline{4}^6 \quad \text{or} \quad 4096 \\ &(\underline{2^2})^6 \\ &2^{12} \end{aligned}$$

$$\frac{\log 4096}{\log 2} = \underline{12}$$

↑
base exp

$$\left(\frac{1}{125}\right)^{2x} = 5^{3x+2} \cdot \sqrt{3125}$$

$$\left(\frac{1}{125}\right)^{2x} = \underline{(5)}^{3x+2} \cdot \underline{(3125)}^{\frac{1}{2}}$$

$$\left(5^3\right)^{2x} = \left(5\right)^{3x+2} \left(5^5\right)^{\frac{1}{2}}$$

$$5^{-6x} = 5^{3x+2} \cdot 5^{5/2}$$

$$5^{-6x} = 5^{3x+2 + \frac{5}{2}}$$

$$\cancel{5}^{-6x} = \cancel{5}^{3x + \frac{9}{2}}$$

$$-6x = \textcircled{3x} + \frac{9}{2}$$

$$\frac{\cancel{-9}x}{\cancel{-9}} = \frac{9}{2} \div -9$$

$$x = \frac{9}{2} \cdot \frac{-1}{9}$$

$$x = -\frac{9}{18} = -\frac{1}{2}$$

Homework

Chapter 7 Review pg. 366-367 (Do all questions)

For $y = c^x$

D: $\{x \mid x \in \mathbb{R}\}$

R: $\{y \mid y > \underline{0}, y \in \mathbb{R}\}$

x int: none

y int: $(0, 1)$

HA: $y = \underline{0}$

For $y = ac^{b(x-h)} + \underline{k}$

D: $\{x \mid x \in \mathbb{R}\}$

R: $\{y \mid y > \underline{k}, y \in \mathbb{R}\}$ (if $a < 0$ then $y < k$)

x int: sub 0 in for y

y int: sub 0 in for x

HA: $y = \underline{k}$


⑥ $\underline{2} = 1.07^x$

$(\underline{1.07})^{10.2} = \underline{1.07}^x$

$10.2 = x$

$\frac{\log 2}{\log 1.07} = \frac{10.2}{\text{Base}} \text{ exp}$

③



decreasing

x	y
-2	16
-1	4
0	1

$\frac{1}{4}$
 $\frac{1}{4}$

$y = \left(\frac{1}{4}\right)^x$

$$\textcircled{11} \text{ b) } 2^{x-2} = \underline{3}^{x+1}$$

$$2^{x-2} = (2)^{1.58x+1}$$

$$2^{x-2} = 2^{1.58x+1.58}$$

$$x-2 = 1.58x + 1.58$$

$$-2 - 1.58 = 1.58x - x$$

$$\underline{-3.58} = \underline{0.58x}$$

0.58 0.58

$$\boxed{-6.17 = x}$$

$$\frac{\log 3}{\log 2} = 1.58$$

Base exp

⑫ Ni-65 half-life of 2.5 h

$$\text{Base} = \frac{1}{2}$$

$$\text{exp} = \frac{t}{2.5}$$

$$A_F = A_0 \left(\frac{1}{2} \right)^{\frac{t}{2.5}}$$

Initial Amount = A_0

$$y = a(c)^{b(x-h)} + k$$
$$(x, y) \rightarrow \left[\frac{1}{b}x + h, ay + k \right]$$
$$y = \text{Initial Amount}(\text{Base})^{\text{exp.}}$$

Finding a common base

$$\frac{\log(\text{have})}{\log(\text{want})} = \text{exp.}$$

Base

Ex:

$$\sqrt{27} \cdot 9^x = 81^{5x-4} \div \left(\frac{1}{3}\right)^x$$

$$27^{1/2} \cdot 9^x = 81^{5x-4} \div \left(\frac{1}{3}\right)^x$$

$$(3^3)^{1/2} \cdot (3^2)^x = (3^4)^{5x-4} \div (3^{-1})^x$$

$$3^{3/2} \cdot 3^{2x} = 3^{20x-16} \div 3^{-x} \quad \begin{array}{l} 20x-16 - (-x) \\ 20x-16+x \end{array}$$

$$\cancel{3}^{2x+3/2} = \cancel{3}^{21x-16}$$

$$2x+3/2 = 21x-16$$

$$\frac{3}{2} + \frac{16}{1} = 21x - 2x$$

$$\frac{3}{2} + \frac{32}{2} = 19x$$

$$\frac{35}{2} = \frac{19x}{19}$$

$$\frac{35}{2} \times \frac{1}{19} = x$$

$$\boxed{\frac{35}{38} = x}$$





7.1 Characteristics of Exponential Functions, pages 334-345

1. Match each item in set A with its graph from set B.

Set A

- a) The population of a country, in millions, grows at a rate of 1.5% per year. *(Increasing) B*
- b) $y = 10^x$ *(Increasing) D*
- c) Tungsten-187 is a radioactive isotope that has a half-life of 1 day. *(Decreasing) A*
- d) $y = 0.2^x$ *(Decreasing) C*

Set B



③ (x, y)

-2, 16	>	4
-1, 4	>	4
0, 1	>	4

$y = c^x$ (Find c)

$y = (\frac{1}{4})^x$

$$16 \left| \begin{array}{c} (\frac{1}{4})^{-2} \\ 4^2 \\ 16 \end{array} \right. \checkmark$$

⑤ $y = -2(4)^{3(x-1)} + 2$ $a = -2$ $b = 3$ $h = 1$ $k = 2$

$y = 4^x$ $(x, y) \rightarrow (\frac{1}{3}x + 1, -2y + 2)$

x	y
-2	1/16
-1	1/4
0	1
1	4
2	16

x	y
1/3	15/8
2/3	3/2
1	0
4/3	-6
5/3	-20

$$\textcircled{9} \text{ c) } \left(\sqrt[3]{216} \right)^5$$

$$(216^{1/3})^5$$

$$(216)^{5/3}$$

$$\frac{\log 216}{\log 6} = 3$$

$$(6^3)^{5/3}$$

$$6^5$$

$$y = c^x$$

$$16 = c^{-2}$$

$$16 = \left(\frac{1}{c}\right)^2$$

$$16 = \frac{1}{c^2}$$

$$16c^2 = 1$$

$$c^2 = \frac{1}{16}$$

$$c = \frac{1}{4}$$

$$y = a(c)^{b(x-h)} + k$$

$$b) \quad y = -2(4)^{3(x-1)} + 2$$

a) $a = -2 \rightarrow$ vertical stretch by a factor of 2.
vertical reflection in x-axis

$b = 3 \rightarrow$ horizontal stretch by a factor of $\frac{1}{3}$

$h = 1 \rightarrow$ translate 1 unit right

$k = 2 \rightarrow$ translate 2 units up

b) horizontal stretch $b = 3 \quad (x, y) \rightarrow (\frac{1}{3}x, y)$

c) $(x, y) \rightarrow [\frac{1}{3}x + 1, -2y + 2]$

$y = 4^x$		$-2(4) + 2$
		$-8 + 2$
		-6
$x \mid y$	$x \mid y$	
-2	$\frac{1}{3}$	$\frac{1}{8}$
-1	$\frac{2}{3}$	$\frac{3}{8}$
0	1	0
1	$\frac{4}{3}$	-6
2	$\frac{5}{3}$	-20

graph

b) $D: \{x \mid x \in \mathbb{R}\}$

$R: \{y \mid y < 2, y \in \mathbb{R}\}$

$HA: y = 2$

$x \text{ int } (y=0)$
 $y = -2(4)^{3(x-1)} + 2$
 $0 = -2(4)^{3(x-1)} + 2$

$-2 = -2(4)^{3x-3}$
 $\frac{-2}{-2} = \frac{-2(4)^{3x-3}}{-2}$

$1 = 4^{3x-3}$

$4^0 = 4^{3x-3}$

$0 = 3x - 3$

$\frac{3}{3} = \frac{3x}{3}$

$1 = x$

$y \text{ int } (x=0)$
 $y = -2(4)^{3(x-1)} + 2$
 $y = -2(4)^{3(0-1)} + 2$
 $y = -2(4)^{-3} + 2$
 $y = -2\left(\frac{1}{64}\right) + 2$
 $y = \frac{-2}{64} + 2$
 $y = \frac{-1}{32} + \frac{64}{32} = \frac{63}{32}$

Test

Open Response:

- Question like your assignment (Transforming exponential functions)
- Word problem → $y = (\text{Initial Amount})(\text{Base})^{\text{time it takes...}}$
- exponential equation.

