

Correct Homework Sheet

$$\begin{aligned} \textcircled{2} \quad \frac{1 - \cos 2\theta}{\sin^2 \theta} &= 2 \\ \frac{1 - (\cos^2 \theta - \sin^2 \theta)}{\sin^2 \theta} & \\ \frac{1 - \cos^2 \theta + \sin^2 \theta}{\sin^2 \theta} & \\ \frac{\sin^2 \theta + \sin^2 \theta}{\sin^2 \theta} & \\ \frac{2\sin^2 \theta}{\sin^2 \theta} & \\ 2 & \end{aligned}$$

$$\begin{aligned} \textcircled{3} \quad \sin(x+y)\sin(x-y) &= \cos^2 y - \cos^2 x \\ (\sin x \cos y + \cos x \sin y)(\sin x \cos y - \cos x \sin y) & \\ \sin^2 x \cos^2 y - \cos^2 x \sin^2 y & \\ (1 - \cos^2 x) \cos^2 y - \cos^2 x (1 - \cos^2 y) & \\ \cos^2 y - \cos^2 x \cos^2 y - \cos^2 x + \cos^2 x \cos^2 y & \\ \cos^2 y - \cos^2 x & \end{aligned}$$

$$\begin{aligned} \textcircled{4} \quad \tan^4 \theta &= \sec^4 \theta (1 - 2\cos^2 \theta + \cos^4 \theta) \\ \frac{\sin^4 \theta}{\cos^4 \theta} & \quad \frac{1}{\cos^4 \theta} (1 - 2\cos^2 \theta + \cos^4 \theta) \\ \frac{(\sin^2 \theta)(\sin^2 \theta)}{\cos^4 \theta} & \quad \boxed{\frac{1 - 2\cos^2 \theta + \cos^4 \theta}{\cos^4 \theta}} \\ \frac{(1 - \cos^2 \theta)(1 - \cos^2 \theta)}{\cos^4 \theta} & \\ \boxed{\frac{1 - 2\cos^2 \theta + \cos^4 \theta}{\cos^4 \theta}} & \end{aligned}$$

$$\textcircled{1} \quad \frac{\cos\theta}{1+\sin\theta} + \frac{1+\sin\theta}{\cos\theta} = 2\sec\theta$$

$$\frac{\cos^2\theta + (1+\sin\theta)(1+\sin\theta)}{\cos\theta(1+\sin\theta)}$$

$$2\left(\frac{1}{\cos\theta}\right)$$

$$\frac{\cos^2\theta + 1 + 2\sin\theta + \sin^2\theta}{\cos\theta(1+\sin\theta)}$$

$$\frac{2}{\cos\theta}$$

$$\frac{2+2\sin\theta}{\cos\theta(1+\sin\theta)}$$

$$\frac{2(1+\sin\theta)}{\cos\theta(1+\sin\theta)}$$

$$\frac{2}{\cos\theta}$$

$$(16) \quad \frac{\sin^4 \theta - \cos^4 \theta}{\sin^2 \theta \cos^2 \theta - \cos^4 \theta} = \frac{\csc^2 \theta}{\cot^2 \theta}$$

$$\frac{(\sin^2 \theta + \cos^2 \theta)(\sin^2 \theta - \cos^2 \theta)}{\cos^2 \theta (\sin^2 \theta - \cos^2 \theta)}$$

$$\frac{1}{\cos^2 \theta}$$

$$\frac{1}{\sin^2 \theta} \div \frac{\cos^2 \theta}{\sin^2 \theta}$$


$$\frac{1}{\sin^2 \theta} \times \frac{\sin^2 \theta}{\cos^2 \theta}$$

$$\frac{1}{\cos^2 \theta}$$

Derivatives of Trigonometric Functions

The Sine Function

- We recall that the derivative $f'(x)$ of a function $f(x)$ gives the slope of the tangent.
- On the next slide we graph $f(x) = \sin x$ together with $f'(x)$, as determined by the slope of the tangent to the sine curve.
 - Note that x is measured in radians.
- The derivative graph resembles the graph of the cosine!



x	$f'(x)$
0	1
$\frac{\pi}{2}$	0
π	-1
$\frac{3\pi}{2}$	0
2π	1

Let's check this using the definition of a derivative...

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h} \right] \\
 &= \lim_{h \rightarrow 0} \left[\sin x \left(\frac{\cos h - 1}{h} \right) + \cos x \left(\frac{\sin h}{h} \right) \right] \\
 &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h}
 \end{aligned}$$

■ Our calculations have brought us to four limits, two of which are easy:

■ Since x is constant while $h \rightarrow 0$,

$$\lim_{h \rightarrow 0} \sin x = \sin x \quad \text{and} \quad \lim_{h \rightarrow 0} \cos x = \cos x$$

■ With some work we can also show that

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \quad \text{and} \quad \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} = 0$$

■ Thus our guess is confirmed:

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \lim_{h \rightarrow 0} \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\
 &= (\sin x) \cdot 0 + (\cos x) \cdot 1 = \cos x
 \end{aligned}$$

Rules to differentiate trigonometric functions:

$$y = \sin x \quad \begin{array}{l} u = x \\ du = 1 \end{array}$$

$$\frac{dy}{dx} = \cos x \cdot 1$$

Given that "u" represents some differentiable function...

$$\frac{dy}{dx} = \cos x$$

$$\frac{d}{du}(\sin u) = \cos u \cdot du$$

$$\frac{d}{du}(\csc u) = -\csc u \cot u \cdot du$$

$$\frac{d}{du}(\cos u) = -\sin u \cdot du$$

$$\frac{d}{du}(\sec u) = \sec u \tan u \cdot du$$

$$\frac{d}{du}(\tan u) = \sec^2 u \cdot du$$

$$\frac{d}{du}(\cot u) = -\csc^2 u \cdot du$$

Let's Practice...

Differentiate the following:

$$y = \sin 3x$$

$$u = \underline{3x}$$

$$du = \underline{3}$$

$$y' = \cos u \cdot du$$

$$y' = \cos(\underline{3x}) \cdot \underline{3}$$

$$\boxed{y' = 3\cos(3x)}$$

$$y = \sin(x + 2)$$

$$u = x + 2$$

$$du = 1$$

$$\boxed{y' = \cos(x + 2)}$$

$$y = \sin(kx + d)$$

$$u = kx + d$$

$$du = k$$

$$\boxed{\frac{dy}{dx} = k\cos(kx + d)}$$

Ex #2.

Differentiate:

a) $y = \sin(x^3)$

$u = x^3$

$du = 3x^2$

$y' = \cos(x^3) \cdot 3x^2$

$y' = 3x^2 \cos(x^3)$

$u = x$

$du = 1$

$y = (\sin x)^3$

b) $y = \sin^3 x$

$\frac{dy}{dx} = 3(\sin x)^2 (\cos x \cdot 1)$

$\frac{dy}{dx} = 3\sin^2 x \cos x$

c) $y = \sin^3(x^2 - 1)$

$y = [\sin(x^2 - 1)]^3$

$u = x^2 - 1$

$du = 2x$

$y' = 3[\sin(x^2 - 1)]^2 \cdot \cos(x^2 - 1) \cdot 2x$

$y' = 6x \sin^2(x^2 - 1) \cos(x^2 - 1)$

Ex #3.

Differentiate:

$$y = (x^2)(\cos x) \quad \leftarrow \begin{array}{l} \text{Product Rule} \\ u = x \\ du = 1 \end{array}$$

$$\frac{dy}{dx} = 2x \cos x + x^2 (-\sin x)(1)$$

$$\frac{dy}{dx} = 2x \cos x - x^2 \sin x$$


$$\frac{dy}{dx} = x(2 \cos x - x \sin x)$$

Homework

Do Questions 1 and 3 from Exercise 7.2 Page 313

 Worksheet on derivatives of trigonometric functions

Negative Angles



Exercise 7.2

① a) $y = \cos(-4x)$ $u = -4x$
 $du = -4$

$y' = -\sin u \cdot du$
 $y' = -\sin(-4x) \cdot -4$
 $y' = 4\sin(-4x)$
 $y' = -4\sin(4x)$

|

$y = \cos(4x)$ $u = 4x$
 $du = 4$

$y' = -\sin u \cdot du$
 $y' = -\sin(4x) \cdot 4$
 $y' = -4\sin(4x)$

Attachments

Derivatives Worksheet.doc